



November 2019

CNOOC UGANDA LTD.

# Biodiversity Impact Assessment for the Proposed Kingfisher Field Development Area and Feeder Pipeline, Hoima & Kikuube District, Uganda

**Submitted to:**  
CNOOC Uganda Ltd.

Report

**Report Number:** 1776816\_D.0

**Distribution:**





## Table of Contents

<b>1.0 INTRODUCTION</b>	<b>1</b>
<b>2.0 TERMS OF REFERENCE</b>	<b>2</b>
2.1 Objectives	2
2.2 Scope of Work	3
2.2.1 Terrestrial Biodiversity	3
2.2.2 Aquatic Biodiversity	3
2.2.3 Overall Biodiversity Value	3
<b>3.0 METHODOLOGY</b>	<b>4</b>
3.1 Key Issues	4
3.1.1 Delineation of the Study Area	4
3.1.2 Spatial Boundaries	5
3.2 Baseline	1
3.2.1 Terrestrial Biodiversity	1
3.2.1.1 Vegetation Communities and Flora Species	1
3.2.1.2 Invertebrates	2
3.2.1.3 Amphibians and Reptiles	3
3.2.1.4 Birds	4
3.2.1.5 Mammals	4
3.2.2 Aquatic Biodiversity	4
3.2.2.1 Water Quality	5
3.2.2.2 Phytoplankton and Macrophytes	5
3.2.2.3 Zooplankton	6
3.2.2.4 Macro-invertebrates	6
3.2.2.5 Fish	7
3.2.3 Overall Biodiversity Value	7
3.2.3.1 Ecosystems and Habitats of Concern	8
3.2.3.1.1 Ecosystem Integrity	8
3.2.3.2 Species of Concern	9
3.2.3.3 IFC Priority Habitat	11
3.2.3.3.1 Natural and Modified Habitat	11
3.2.3.3.2 Critical Habitat	11





3.3	Impact Assessment .....	12
3.3.1	Key Questions and Indicators .....	12
3.3.2	Impact Assessment.....	13
3.4	Limitations .....	14
<b>4.0</b>	<b>RELEVANT LEGISLATION.....</b>	<b>14</b>
4.1	Uganda’s Policy, Legal and Institutional Framework on Biodiversity .....	14
4.1.1	The Constitution of the Republic of Uganda (1995) .....	14
4.1.2	Uganda Wildlife Bill (2017).....	15
4.1.2.1	Uganda Wildlife Act (1996).....	15
4.1.2.2	Uganda Wildlife Act Cap 200 of 2000 .....	15
4.1.2.3	Uganda Wildlife Policy (1999, 2014).....	15
4.1.2.4	Uganda Forestry Policy (2001) and the National Forestry and Tree Planting Act (2003) .....	16
4.1.2.5	The Land Act (1995).....	17
4.1.2.6	Uganda National Land Policy (2013) .....	17
4.1.3	Uganda National Biodiversity Strategy and Action Plan (2015-2025) .....	17
4.2	Conventions and International Agreements .....	18
4.3	International Guidance.....	18
4.3.1	International Finance Corporation’s Performance Standards.....	18
4.3.2	Regional Frameworks .....	20
<b>5.0</b>	<b>KEY ISSUES RELATING TO BIODIVERSITY .....</b>	<b>20</b>
<b>6.0</b>	<b>BASELINE ENVIRONMENT.....</b>	<b>21</b>
6.1	Terrestrial Biodiversity .....	21
6.1.1	Vegetation Communities and Flora Species .....	21
6.1.1.1	Vegetation Communities.....	21
6.1.1.1.1	Wooded Grassland.....	1
6.1.1.1.2	Woodland .....	2
6.1.1.1.3	Thicket-Grassland Mosaic.....	3
6.1.1.1.4	Open Grassland .....	4
6.1.1.1.5	Modified Habitats.....	4
6.1.1.2	Flora Species.....	5
6.1.2	Invertebrates .....	6
6.1.2.1	Butterflies.....	6
6.1.2.2	Dragonflies and Damselflies .....	7





6.1.3	Reptiles and Amphibians .....	7
6.1.3.1	Amphibians .....	8
6.1.3.2	Reptiles.....	8
6.1.4	Birds.....	9
6.1.5	Mammals .....	11
6.2	Aquatic Biodiversity .....	12
6.2.1	Water Quality .....	12
6.2.2	Phytoplankton .....	14
6.2.3	Zooplankton .....	16
6.2.4	Macro-invertebrates .....	17
6.2.5	Fish .....	19
<b>7.0</b>	<b>OVERALL BIODIVERSITY VALUE.....</b>	<b>21</b>
7.1	Ecosystems and Habitats of Concern.....	21
7.1.1	Near-Shore Habitats of Lake Albert .....	22
7.1.2	Escarpment Vegetation Corridors .....	25
7.1.3	Wetlands and Riparian Habitats.....	29
7.1.4	Bugoma Central Forest Reserve.....	31
7.2	Species of Concern .....	33
7.2.1	Threatened, Range-Restricted/Endemic and Statutory Species .....	33
7.2.2	Species of Economic and/or Cultural Importance .....	33
7.2.3	Species listed under CITES .....	34
7.2.4	Evolutionarily Distinct Species .....	34
7.2.5	Invasive and Potentially Invasive Species.....	35
<b>8.0</b>	<b>NATURAL AND MODIFIED HABITAT WITHIN THE CHAA.....</b>	<b>37</b>
<b>9.0</b>	<b>CRITICAL HABITAT WITHIN THE CHAA.....</b>	<b>38</b>
9.1	Assessment of Potential Triggers of Critical Habitat .....	38
9.1.1	Species Potentially Triggering Criterion 1, 2 and 3 Critical Habitat .....	38
9.1.1.1	Mud Snail ( <i>Gabbiella candida</i> ).....	38
9.1.1.2	Snail ( <i>Bellamya rubicunda</i> ).....	39
9.1.1.3	Madagascar Pond Heron ( <i>Ardeola idae</i> ) .....	40
9.1.1.4	Grey Crowned Crane ( <i>Balearica regulorum</i> ) .....	41
9.1.1.5	White-backed Vulture ( <i>Gyps africanus</i> ) .....	42
9.1.1.6	Hooded Vulture ( <i>Necrosyrtes monachus</i> ).....	44





9.1.1.7	Nahan’s Francolin ( <i>Ptilopachus nahani</i> ) .....	45
9.1.1.8	Eastern Chimpanzee ( <i>Pan troglodytes schweinfurthii</i> ) .....	46
9.1.1.9	Migratory and Congregatory Species .....	47
9.1.2	Ecosystems Potentially Triggering Criterion 4 Critical Habitat: Highly Threatened and/or Unique Ecosystems.....	53
9.1.2.1	Bugoma Central Forest Reserve .....	54
9.1.3	Species Potentially Triggering Criterion 5 Critical Habitat (Evolutionarily Distinct Species) .....	54
9.1.3.1	Shoebill ( <i>Balaeniceps rex</i> ) .....	54
9.1.3.2	Secretarybird ( <i>Sagittarius serpentarius</i> ) .....	55
9.1.4	Species/Ecosystems triggering other Qualitative Critical Habitat Criteria .....	56
9.2	Confirmed Triggers of Critical Habitat in the CHAA .....	56
<b>10.0</b>	<b>IMPACT ASSESSMENT – PRODUCTION FACILITY.....</b>	<b>60</b>
10.1	Construction and Decommissioning Phase Impacts.....	61
10.1.1	What impact could the construction/decommissioning of the Project have on habitats and ecosystem integrity? .....	62
10.1.1.1	What impact could the construction/decommissioning of the Project have on the near-shore aquatic habitats of Lake Albert? .....	63
10.1.1.2	What impact could the construction/decommissioning of the Project have on the wetlands in the Buhuka Flats region of the CHAA?.....	69
10.1.1.3	What impact could the construction/decommissioning of the Escarpment Road have on the escarpment vegetation corridors?.....	81
10.1.1.4	What impact could the construction/decommissioning of the Project have on the Bugoma Central Forest Reserve?.....	85
10.1.2	What impact could the construction/decommissioning of the Project have on species of concern? .....	88
10.1.2.1	What impact could the construction/decommissioning of the Production Facility have on the Mud Snail ( <i>Gabbiella candida</i> )?.....	88
10.1.2.2	What impact could the construction/decommissioning of the Production Facility have on Grey Crowned Crane?.....	92
10.1.2.3	What impact could the construction/decommissioning of the Production Facility have on Nahan’s Francolin?.....	96
10.1.2.4	What impact could the construction/decommissioning of the Production Facility have on Eastern Chimpanzee? .....	100
10.2	Operation Phase Impacts .....	105
10.2.1	What impact could the operation of the Project have on habitats and ecosystem integrity? .....	105
10.2.1.1	What impact could the operation of the Project have on the near-shore environment of Lake Albert? .....	106





10.2.1.2	What impact could the operation of the Project have on the wetlands in the Buhuka Flats region of the CHAA?.....	111
10.2.1.3	What impact could the operation of the Project have on the escarpment vegetation corridors?.....	115
10.2.1.4	What impact could the operation of the Project have on the Bugoma Central Forest Reserve? .....	118
10.2.2	What impact could the operation of the Project have on species of concern? .....	120
10.2.2.1	What impact could the operation of the Project have on the Mud Snail?.....	121
10.2.2.2	What impact could the operation of the Project have on Grey Crowned Crane? .....	123
10.2.2.3	What impact could the operation of the Project have on Nahan’s Francolin?.....	126
10.2.2.4	What impact could the operation of the Project have on Eastern Chimpanzee? .....	130
<b>11.0</b>	<b>IMPACT ASSESSMENT – FEEDER PIPELINE .....</b>	<b>133</b>
11.1	Construction Phase Impacts.....	135
11.1.1	What impact could the construction of the Feeder Pipeline have on habitats and ecosystem integrity? .....	136
11.1.1.1	What impact could the construction of the Feeder Pipeline have on the near-shore aquatic habitats of Lake Albert? .....	136
11.1.1.2	What impact could the construction of the Feeder Pipeline have on the escarpment vegetation corridors?.....	139
11.1.1.3	What impact could the construction of the Feeder Pipeline have on wetlands and riparian habitat in the CHAA? .....	143
11.1.1.4	What impact could the construction of the Feeder Pipeline have on Bugoma CFR?.....	148
11.1.2	What impact could the construction of the Feeder Pipeline have on species of concern? .....	151
11.1.2.1	What impact could the construction of the Feeder Pipeline have on the Mud Snail? .....	151
11.1.2.2	What impact could the construction of the Feeder Pipeline have on Grey Crowned Crane?.....	152
11.1.2.3	What impact could the construction of the Feeder Pipeline have on Nahan’s Francolin?.....	156
11.1.2.4	What impact could the construction of the Feeder Pipeline have on Eastern Chimpanzee? .....	160
11.2	Operation Phase Impacts .....	163
11.2.1	What impact could the operation of the Feeder Pipeline have on habitats and ecosystem integrity? .....	163
11.2.1.1	What impact could the operation of the Feeder Pipeline have on the escarpment vegetation corridors?.....	163
11.2.1.2	What impact could the operation of the Feeder Pipeline have on wetlands and riparian habitat in the CHAA? .....	166
11.2.1.3	What impact could the operation of the Feeder Pipeline have on Bugoma CFR? .....	169
11.2.2	What impact could the operation of the Feeder Pipeline have on species of concern? .....	171
11.2.2.1	What impact could the operation of the Feeder Pipeline have on Grey Crowned Crane? .....	171
11.2.2.2	What impact could the operation of the Feeder Pipeline have on Nahan’s Francolin?.....	174





11.2.2.3 What impact could the operation of the Feeder Pipeline have on Eastern Chimpanzee? ..... 176

**12.0 RECOMMENDATIONS FOR MITIGATION AND MONITORING ..... 179**

12.1 Mitigation Hierarchy ..... 179

12.2 Site-Specific Mitigation Measures and Schedule for Completion ..... 181

12.2.1 Pre-construction Phase ..... 181

12.2.2 Construction and Decommissioning Phase ..... 182

12.2.3 Operation Phase ..... 195

**13.0 OFFSETTING STRATEGY ..... 204**

13.1 Construction and Operation Phase Impacts on the Near-Shore Environment of Lake Albert ..... 204

13.2 Construction and Operation Phase Impacts on Wetlands in the CHAA ..... 204

13.3 Construction and Operation Phase Impacts on the Mud Snail *G. candida* ..... 205

13.4 Construction and Operation Phase Impacts on Bugoma CFR ..... 206

13.5 Construction and Operation Phase Impacts on Nahan’s Francolin ..... 207

13.6 Construction and Operation Phase Impacts on Chimpanzee ..... 207

**14.0 REFERENCES ..... 210**

**TABLES**

Table 1: Key questions and indicators ..... 12

Table 2: Extent of escarpment corridor vegetation communities within the CHAA ..... 26

Table 3: Natural and modified habitat in the CHAA ..... 37

Table 4: Triggers of critical habitat in the CHAA ..... 57

Table 5: Biodiversity Valued Components for Impact Assessment ..... 60

Table 6: Area of vegetation communities at baseline, disturbance and loss due to Production Facility, and net change (% loss) in CHAA ..... 62

Table 7: Potential impacts in the construction phase to near-shore habitats ..... 69

Table 8: Wetland areas directly affected by the construction of the production facility ..... 70

Table 9: Potential impacts in the construction phase to the wetlands of the CHAA ..... 81

Table 10: Permanent loss of escarpment vegetation due to construction of escarpment road ..... 82

Table 11: Potential impacts in the construction phase to the vegetation communities of the escarpment ..... 84

Table 12: Potential impacts in the construction phase to the Bugoma Central Forest Reserve ..... 88

Table 13: Potential impacts in the construction phase to the Mud Snail ..... 91

Table 15: Potential impacts in the construction phase to the Nahan’s Francolin ..... 99

Table 16: Potential impacts in the construction phase to the Eastern Chimpanzee ..... 105

Table 17: Potential impacts in the operation phase to near-shore habitats ..... 110

Table 18: Potential impacts in the operation phase to the wetlands of the CHAA ..... 114





Table 19: Potential impacts in the operation phase to the vegetation communities of the escarpment.....	118
Table 20: Potential impacts in the operation phase to the Bugoma Central Forest Reserve .....	120
Table 21: Potential impacts in the operation phase to the Mud Snail.....	122
Table 22: Potential Production Facility operation phase impacts to Grey Crowned Crane .....	126
Table 23: Potential impacts in the operational phase to the Nahan’s Francolin .....	129
Table 24: Potential impacts in the operation phase to the Eastern Chimpanzee .....	133
Table 25: Biodiversity Valued Components for Feeder Pipeline Impact Assessment .....	134
Table 26: Potential impacts in the construction phase to near-shore habitats.....	138
Table 27: Permanent loss of escarpment corridor vegetation due to construction of Feeder pipeline .....	139
Table 28: Potential impacts in the construction phase to the vegetation communities of the escarpment .....	142
Table 29: Area of wetland and riparian habitat impacted along the 30m wide pipeline construction right of way	143
Table 30: Potential impacts of construction of the Feeder pipeline on the wetlands of the CHAA .....	148
Table 32: Potential impacts in the construction phase to the Mud Snail .....	152
Table 35: Potential impacts in the construction phase to the Eastern Chimpanzee .....	162
Table 36: Potential impacts in the operation phase of the Feeder Pipeline to the vegetation communities of the escarpment .....	166
Table 37: Potential impacts of operation of the Feeder pipeline on the wetlands of the CHAA.....	169
Table 39: Potential operation phase impacts to Grey Crowned Crane.....	173
Table 40: Potential impacts in the operational phase to the Nahan’s Francolin .....	176
Table 41: Potential impacts in the operation phase to the Eastern Chimpanzee .....	178
Table 42: Mitigation and management measures for biodiversity, during the construction and decommissioning phases of the Project.....	187
Table 43: Mitigation and management measures for biodiversity, during the operation phase of the Project .....	199
Table 44: Post-offset residual risk: construction phase impacts on near-shore environment of L. Albert.....	204
Table 45: Post-offset residual risk: construction phase impacts on wetlands in the CHAA .....	205
Table 46: Post-offset residual risk: construction phase impacts on Mudsnailed in the CHAA .....	206
Table 47: Post offset residual risk: Construction and operation phase impacts on Bugoma CFR.....	207
Table 48: Post offset residual risk: construction and operation phase impacts on Nahan’s Francolin .....	207
Table 49: Post offset residual risk: construction and operation phase impacts on Eastern Chimpanzee .....	208
Table 50: Possible offset options as part of the mitigation strategy for the Project .....	209

## FIGURES

Figure 1: Biodiversity LSA .....	1
Figure 2: Biodiversity CHAA.....	2
Figure 3: Biodiversity LSA and CHAA .....	3
Figure 4: Land cover and land use in the CHAA .....	1
Figure 5: Vegetation communities of the CHAA (focus on Buhuka Flats) .....	1







Figure 6: Vegetation communities of the CHAA .....	1
Figure 7: Typical wooded grassland in the CHAA .....	1
Figure 8: Typical woodland along a watercourse in the CHAA.....	2
Figure 9: Typical Thicket-grassland mosaic community in the LSA .....	3
Figure 10: Typical grassland community of the LSA .....	4
Figure 11: Escarpment vegetation communities .....	26
Figure 12: 'Chutes' used for transport of thatching grass harvested from the escarpment.....	28
Figure 13: Cattle watering at Kamansinig crossing, Buhuka Flats (Nov 2017).....	31
Figure 14: Invasive species recorded on the Buhuka Flats .....	36
Figure 15: Natural and Modified habitat in the CHAA.....	37
Figure 16: Critical habitat in the CHAA.....	59
Figure 17: Rivers and catchments on Buhuka Flats .....	65
Figure 18: Wetlands in the Buhuka Flats directly impacted by construction/decommissioning of the Production Facility .....	71
Figure 19: Impact of a poorly-designed road crossing on wetland in Buhuka flats.....	73
Figure 20: Direction of storm flows from construction sites .....	75
Figure 21: Proposed Government road upgrades in the region .....	86
Figure 22: The P1 road through Bugoma Central Forest Reserve .....	87

## APPENDICES

### APPENDIX A

Document Limitations

### APPENDIX B

Critical Habitat Approach and Method of Assessment

### APPENDIX C

Terrestrial Ecology Baseline Study

### APPENDIX D

Aquatic Ecology Baseline Study

### APPENDIX E

Land Cover Classification

### APPENDIX F

Impact Assessment Approach and Methods

### APPENDIX G

Critical Habitat Screening and Appraisal





### 1.0 INTRODUCTION

CNOOC Uganda Limited (“CNOOC”) has identified an opportunity to develop the Kingfisher Field Development area (KFDA) on the eastern shore of Lake Albert, Kikuube District, with a feeder pipeline connecting the KFDA to the refinery in Kabaale, Hoima District, all in Uganda. In accordance with Ugandan law, it is necessary for CNOOC to determine the potential environmental and social impacts of the project, and to demonstrate how these will be mitigated and managed. This chapter of the Environmental and Social Impact Assessment (ESIA) presents an assessment of the potential impacts of the Kingfisher Field Development Area (KFDA) (the Project) on biodiversity, and sets out recommendations for their avoidance and reduction, where necessary. This impact assessment has been developed with reference to the baseline terrestrial and aquatic biodiversity surveys, completed between February and March 2014, May and June 2014 and October and November 2014. The baseline was required to enable an appropriate assessment of the Project’s potential impacts on biodiversity. The detailed results of the baseline surveys are included in the Appendices and summarised in Section 6.0.

Uganda is a signatory to the Convention on Biological Diversity (CBD); as such there is a governmental requirement to implement policies to protect biodiversity at all its different levels, including ecosystems, species and genes. That protection is embodied in the obligations the country imposes on entities operating within its borders. Nevertheless, as mentioned, biodiversity as a concept is very broad, and is typically defined as the variety of life at different levels of biological organisation and all the ecological and biological processes through which they are connected (for example, see Hill et al. 2005; Secretariat of the CBD 2006). In line with that definition of biodiversity, this impact assessment focuses on those different levels, as defined below.

Ecosystems are a dynamic complex of plants, animals, micro-organisms, and their non-living environment, interacting as a functional unit. Ecosystems can vary greatly in size, and in the biotic and abiotic elements of which they are comprised. However, ecosystems usually encompass specific, defined spaces. Ecosystems are distinct from communities in that the term “community” typically only refers to coexisting biotic populations, whereas ecosystems can include abiotic (that is, non-living) components and an array of environmental processes (Begon et al. 1990). Species can be defined as groups of morphologically similar organisms that have descended from a common ancestor, with common genetic make-up, and which produce fertile offspring only amongst themselves (Begon et al. 1990). Species are the basic components of ecological communities and are the most recognisable units of biodiversity. Efforts to conserve biodiversity often focus at the species level, and the efforts to conserve species diversity go some way to include the genetic component of biodiversity.

A fourth level of biodiversity has been derived fairly recently in the scientific and wider literature. This is the concept of ecosystem services (see Millennium Ecosystem Assessment 2005). Ecosystem services are the benefits to people generated by a functioning natural environment, the recognition of which has become increasingly important. The assessment of ecosystem services is primarily covered under the International Finance Corporation’s (IFC) Performance Standard 6 (PS6), although it is also covered under PS1, PS3, PS4, PS5, PS7 and PS8. Nevertheless, for the intents of this impact assessment, the assessment of ecosystem services has been given its own report in recognition that the assessment is a cross-over discipline covering social, biological and physical disciplines, as reflected in its assessment requirement across multiple performance standards. As such, the concept of ecosystem services, as a component of biodiversity under the IFC PS6, is not discussed further in this document.

In light of the above, and for the purposes of this assessment, “biodiversity” encompasses terrestrial and aquatic ecosystems (at the habitat, species and genetic level), and is in line with the definitions set out by Uganda’s *National Environment Act 1995*, the CBD (Secretariat of the CBD 2006), and the IFC’s PS6 (2012). Further clarification is set out in Section 2.0. Ecosystems services, although recognised as being part of, or dependent upon biodiversity, and covered under the IFC’s Performance Standard 6, are not covered in this chapter. Instead, the Consultant is devoted to describing the baseline and impact assessment for ecosystem services.

This specialist study report includes the following sections:

- Section 2.0 describes the terms of reference for the report.





- Section 3.0 presents the methods used for the study that entail examining the study objectives, the approach employed and the limitations encountered.
- Section 4.0 sets out the legislative background applicable to the study.
- Section 5.0 Summarises the key issues in relation to biodiversity.
- Section 6.0 summarises the results of the baseline studies.
- Section 7.0 assesses the impacts to biodiversity arising from the CPF, wells and associated infrastructure.
- Section 8.0 assesses the impacts to biodiversity arising from the Feeder Pipeline.
- Section 9.0 recommends mitigation and management measures.
- Section 10.0 provides recommendations for offsetting.
- Section 11.0 includes a complete list of references consulted.

This assessment report is a preliminary version produced for client review.

## 2.0 TERMS OF REFERENCE

Historically, the biodiversity of the area has been assessed, in part, as part of the Environmental Impact Assessments (EIA) for various oil-related projects and developments over the last decade. However, those assessments tended to focus only on the footprint of the particular project components; for example, drill pads, access roads, etc. In the wider Albertine-Graben region, broad-scale landscape assessments have been completed (for example, NEMA 2010, AECOM 2013, MEMD 2013; TBC & FFI, 2017). Nevertheless, no comprehensive biodiversity impact assessment has been completed that covers the full scope of the developments on Buhuka Flats, the escarpment and Feeder pipeline corridor, which is the focus of this ESIA.

In determining the requirements of the biodiversity assessment, reference was made to the appropriate Ugandan legislation and guidance, as well as international standards and guidance. National policy and international standards pertaining to the Project are detailed in Section 4.0.

The biodiversity impact assessment concentrates on assessing changes in ecosystems, habitat and ecosystem function, changes in populations of species, including species of conservation concern, invasive species and species of high value to people.

### 2.1 Objectives

The aim of this biodiversity assessment was to collect scientifically defensible, high quality data of sufficient breadth that could be used to characterise the baseline conditions of the area and assess how the Project could affect that biodiversity. This was undertaken in consideration of Uganda's *Wildlife Policy 1999* and *National Biodiversity Strategy and Action Plan* (NEMA 2002), and with reference to the IFC PS6, which seeks to protect biodiversity and ecosystem services from the adverse impacts of project activities, and support its conservation and sustainable use. Consequently, the objectives of the biodiversity impact assessment, as reflected in the Scoping Report, were to:

- Characterise the ecological integrity of the terrestrial and aquatic (including wetland) ecosystems in the Project's area of influence and ascertain seasonal variation.
- Identify sensitive or unique habitats and species (as protected under Ugandan legislation and international obligations), which could suffer irreplaceable loss due to the Project.
- Identify species of concern that could trigger critical habitat (as defined by IFC PS6).
- Identify populations and trends of exotic and invasive species in the Project's area of influence.
- Identify and describe potential sources of risk and impact associated with the development that could affect biodiversity of the Project's area of influence.





- Identify the potential direct, indirect and cumulative effects (ref Chapter 17 Cumulative Impact Assessment) on biodiversity associated with the Project.
- Recommend suitable mitigation measures where applicable.
- Develop a monitoring programme and action plan for the biodiversity affected by the Project's development.

## 2.2 Scope of Work

In order to address the above objectives, and in line with the KFDA Scoping Report, a description and regional contextualisation of the baseline terrestrial and aquatic ecology was undertaken. Using available regional ecological data and dedicated baseline studies, an assessment of the effects on the biodiversity of the Project's area of influence (i.e. the Local Study Area, and the Critical Habitat Area of Analysis – ref. Sections 3.1.1 & 3.1.2) was conducted to meet the requirements of IFC PS6.

The scopes for the baseline terrestrial and aquatic ecology, and overall biodiversity effects assessment are presented in the following sections.

### 2.2.1 Terrestrial Biodiversity

The baseline terrestrial biodiversity studies focussed on describing the seasonal variation (that is, the two wet seasons and a dry season) of:

- Vegetation communities and habitats within the Project's area of influence, including structure, condition, species composition, representativeness, irreplaceability and vulnerability.
- Populations of vertebrates, and selected invertebrates in the Project's area of influence, including their representativeness, irreplaceability and vulnerability.
- Current drivers of change in the terrestrial ecosystems of the Project's area of influence, including populations of pest and invasive species.

### 2.2.2 Aquatic Biodiversity

The baseline aquatic biodiversity studies focussed on describing the seasonal variation (that is, the two wet seasons and a dry season) of:

- Abiotic factors (that is, physical and chemical characteristics of the water quality) influencing the aquatic habitats and ecosystems supported in Lake Albert, wetlands on the Buhuka Flats, and watercourses draining the escarpment and flats.
- Aquatic habitats and ecosystems within the Project's area of influence, including structure, condition, species composition, representativeness, irreplaceability and vulnerability.
- Biotic components of the various aquatic habitats and ecosystems in the Project's area of influence; in particular, macrophytes, phytoplankton, zooplankton, macro-invertebrates and fish, and the condition of the populations of these groups, their representativeness, irreplaceability and vulnerability.
- Current drivers of change in the aquatic ecosystems of the Project's area of influence, including populations of pest and invasive species.

### 2.2.3 Overall Biodiversity Value

The effects that the Project could have on the terrestrial and aquatic ecosystems, habitats, and species in the Project's area of influence were identified and assessed at the scale of the overall biodiversity, that is, a landscape ecology and ecosystems approach, viz., Secretariat of the CBD (2006), as embodied in IFC PS6 (IFC 2012a), the International Petroleum Industry Environmental Conservation Association's (IPIECA) (IPIECA 2005, 2007, 2010), and the Energy and Biodiversity Initiative's (EBI) (EBI 2006) guidance documents. Furthermore, in line with the requirements of IFC PS6 (IFC 2012a, b), the consideration of the effects of the



Project on the biodiversity were based on the findings of the baseline terrestrial and aquatic ecology assessments and focussed on:

- The identification of modified and natural habitat within the Project's Local Study Area (LSA), and the implications for no net loss of biodiversity.
- The identification of species of concern occurring within the Critical Habitat Area of Analysis (CHAA) surrounding the Project, and the potential for these to trigger critical habitat.
- The identification of protected areas, and other internationally recognised areas within and surrounding the Project's area of influence, and the potential for the Project to affect these.
- Assessing the potential effects of the Project on the functions and processes of the ecosystems of the Project's area of influence.

### 3.0 METHODOLOGY

This section presents the methods used to identify and assess the potential effects and impacts to the biodiversity values of the Project's area of influence. This followed a six-part process, broadly following those outlined in Treweek (1999) and Secretariat for the CBD (2006):

- 1) Identify key issues.
- 2) Delineate study areas.
- 3) Identify the timeframe for the assessment.
- 4) Describe the baseline (including current direct and indirect drivers of change to ecosystem processes and functions, composition and structure).
- 5) Identify valued components, key questions, and indicators.
- 6) Conduct the impact assessment:
  - a) identify Project interactions with the environment;
  - b) consider environmental design features and mitigation;
  - c) assess effects and classify direct and indirect impacts;
  - d) describe the confidence in the impact predictions; and
  - e) determine follow-up and monitoring activities.

The impact assessment shows clearly to the reader all of the steps taken to arrive at the overall impact level score for any key issue or question. Hence, a reader should be able to use these same tools to repeat the analysis if they desired to do so.

The methods used for each of these steps are presented below.

### 3.1 Key Issues

Key issues in relation to the biodiversity within the immediate footprint of the Project, and the wider region surrounding the Project footprint, were identified through stakeholder consultation, review of background biodiversity and environmental reports (that is: RPS (2006); AWE (2008a, b, c, 2013a, b, 2014a, b); AECOM (2012, 2013); EA CL (2013, 2014), NEMA (1996, 2002, 2010), MEMD (2013), and TBC & FFI (2017)), published ecological and social literature, consideration of the IFC's Performance Standards (IFC 2012a), and applying the expertise of the biodiversity impact assessment team.

#### 3.1.1 Delineation of the Study Area

As with any environmental impact assessment, the spatial and temporal boundaries for the analysis need to be set. Described below are those bounds.





### 3.1.2 Spatial Boundaries

The spatial boundaries within which potential effects arising from the Project may have on biodiversity were set. For this assessment, and in order to satisfy IFC requirements, two areas of influence were considered in relation to assessing the potential effects on biodiversity:

- a) Biodiversity local study area.
- b) Critical habitat area of analysis.

For the assessment of local impacts, the area should be large enough to analyse and mitigate efficiently the potential effects from the project on the receiving environment, but not too large as to dilute or confound the potential project-related effects with other human-induced and natural influences.

Described below are how the spatial bounds for each of those areas were determined.

#### a) Local Study Area

- The assessment of impacts within the local area of the Project, or biodiversity local study area (LSA), was based on the spatial extent of a Project's footprint and an associated buffer zone that includes potential immediate, direct effects on the receiving environment. It was derived as a focus for the development of a baseline case where potential direct effects were predicted to occur.
- The LSA incorporates: 20 production well pads 1, 2 and 3 and 4 and associated 11 water injection wells, Central Processing Facility (CPF); permanent camp; Bugoma drilling camp; contractor's camp; the airstrip which is being converted to a laydown area; all in-field roads and flow lines; material yards for production and drilling; borrow pit and spoil area A, B and C; crusher plant and bitumen container storage; jetty; new water intake and pump station; materials yard and stockpile area on top of the escarpment; temporary camp on top of the escarpment; the Feeder pipeline easement to Kabaale.
- A 1 km buffer was incorporated around the infrastructure in order to capture all potential direct effects, including those from noise, dust, changes to surface water quality (that is, Lake Albert, streams and wetlands).
- The LSA is depicted in Figure 1.

#### b) Critical Habitat Area of Analysis (CHAA)

- Critical habitat is present in many places globally, but is only relevant to a development project where the project may affect that habitat (both directly and indirectly) (IFC 2012a). Importantly, the determination of critical habitat is independent of the specifics of the proposed project footprint, and is present under baseline conditions and is not defined by the size of the project footprint, or other project effects.
- For the area, a first step in defining critical habitat was to identify an ecologically-relevant area of analysis surrounding, and including, the anticipated extent of the Project's influence, including broader or regional effects from the Project, in association with other anthropogenic activities (such as other projects) and natural factors (ref Chapter 17 Cumulative Impact Assessment). These include indirect, induced and cumulative effects. The CHAA was defined as that area.
- The boundaries of the CHAA were devised cognisant of the need for an area where the ecological and land management issues have more in common with each other than they do with those in adjacent areas, and constitutes a sensible ecological and political boundary within which critical habitat can be defined (IFC 2012b, paragraph 65).
- This area was also used as the geographical extent to screen biodiversity features to be assessed for critical habitat based on discrete management units (DMU). Critical habitat was therefore identified and mapped at the CHAA-scale, which was inclusive of the LSA.
- The screening was initially undertaken at a desktop level using the following attributes:



- Presence, abundance, and distribution within, or relevance to, the area associated with the CHAA.
- Potential for interaction with the area and proposed project development.
- Conservation status or concern; in particular, IUCN-listed Critically Endangered and Endangered species, range restricted and endemic species, congregatory and migratory species, as well as, nationally listed threatened and priority species.
- Ecological and/or socio-economic value.
- Identified importance to interested public, government agencies, the scientific community, NGOs and/or CNOOC.
- secondary data sources were used, especially for the CHAA away from the LSA; these included:
  - SPOT6 imagery for the determination of land cover, land use, natural and modified habitats.
  - Global Biodiversity Information Facility (GBIF) (GBIF 2014; GBIF 2017).
  - Integrated Biodiversity Assessment Tool (IBAT), including the available data on Red List species, Key Biodiversity Areas (KBA), Endemic Bird Areas (EBA), Important Bird Areas (IBA), protected areas, wetland areas (IUCN 2014b).
  - Catchments and hydrology.
  - Soils and geology mapping.
  - Existing infrastructure and disturbance.
  - Proposed Project infrastructure.
- A biodiversity constraints/sensitivity map of the wider area was then generated, which became the CHAA. This map also formed the basis to identify modified and natural habitats (as per IFC 2012b), focus the assessment of the valued components, and guide field surveys.
- For the intents of this biodiversity impact assessment, the CHAA encompasses: the Buhuka Flats; the catchments of: the Masika River, the two unnamed watercourses to the south of the Masika River, the Kamansinig River, and the four unnamed watercourses to the north of the Kamansinig River; the pipeline corridor, extending to the eastern boundary of the natural vegetation on the escarpment (as derived from SPOT6 imagery); and the Bugoma Central Forest Reserve (CFR).
  - DMUs are defined as an area with a clearly demarcated boundary within which the biological communities and/or management issues have more in common with each other than they do with those in adjacent areas (IFC 2012b).
  - DMUs forming the CHAA included: the Buhuka Flats; the catchments of the Masika River, the two unnamed watercourses to the south of the Masika River, the Kamansinig River, and the four unnamed watercourses to the north of the Kamansinig River; the Bugoma Central Forest Reserve; the subsistence agricultural areas between the escarpment and the Bugoma Central Forest Reserve.
- Figure 2 depicts the CHAA.



# BIODIVERSITY IMPACT ASSESSMENT




Figure 1: Biodiversity LSA







# BIODIVERSITY IMPACT ASSESSMENT




Figure 2: Biodiversity CHAA





# BIODIVERSITY IMPACT ASSESSMENT




Figure 3: Biodiversity LSA and CHAA





### 3.2 Baseline

A baseline of the terrestrial and aquatic ecology of the LSA was determined through a desktop assessment and comprehensive, seasonal surveys. These assessments and surveys were then used as the basis for the description of the biodiversity of the CHAA. The approach and methods for those baseline surveys and derived studies are presented below. The detailed baseline study reports for the terrestrial and aquatic ecology are presented in Appendix C.

#### 3.2.1 Terrestrial Biodiversity

In addition to a detailed review of existing literature and databases, multi-season surveys of the terrestrial ecology of fixed sampling locations within the LSA were completed to describe the baseline conditions. The field surveys focused on the areas of the CPF, construction camp, spoil area, well pads, contractors' camp, parking lot, the existing camp, permanent camp, the airstrip, the borrow pit, and the pipeline route.

Seasonal variation in terrestrial ecology was assessed through multi-season surveys. The first field surveys captured the dry season, and occurred from 25 February 2014 to 8 March 2014. The second set captured the first wet season, and occurred from 28 May 2014 to 23 June 2014. The third set captured the second wet season, and occurred from 28 October to 20 November 2014.

The surveys focussed on:

- vegetation communities and flora species;
- invertebrates;
- reptiles and amphibians;
- birds; and
- mammals.

The methods of survey for each of these groups are summarised below, and presented in detail in the Appendices.

##### 3.2.1.1 *Vegetation Communities and Flora Species*

A desktop search and review was made of available literature about the vegetation and flora of the CHAA, in particular the vegetation communities of the Albertine Graben, the escarpment and beyond, and the conservation status of species. Data sources included: Langdale-Brown et al. (1964); Plumptre et al. (2003, 2007); Kalema and Beentje (2012); and the IUCN's RedList (IUCN 2017). The data thus obtained were used to identify sampling and survey sites on satellite imagery for later ground-truthing during the field surveys. These data were also used to inform the assessment of the probability of species of concern occurring in the CHAA (see Section 3.3.3.2 Species of Concern).

Two-hundred-and-seventy-five sampling plots were identified for survey; in particular: on the Buhuka Flats, 13 line-transects were run, with 150 sample plots; on the escarpment, seven transects were run, with 40 sample plots; and along the pipeline route, 23 transects were run with 153 plots.

The precise locations of each of those plots are displayed in the baseline reports in Appendix C.

Areas of natural and modified vegetation were surveyed in the wet and dry seasons, with focus given to the main vegetation communities on the Buhuka Flats, the escarpment and along the pipeline route. Along the pipeline route, only relic patches of vegetation were selected for survey that were deemed natural or near-natural, because most of the area along the route is under cultivation and settlement. Sampling was undertaken based on the gradsects approach (after Gillison and Brewer 1985), taking into consideration the variation in such aspects as slope, observable indicators of soil moisture and soil type, as well as the plant community assemblages. All species of plants present were identified and recorded, and their relative abundance was assessed using the DAFOR scale (D = dominant, A = abundant, F = frequent, O = occasional, R = rare) (after Kent and Coker 1992). Azonal habitats believed to be unique within a given area were also sampled. These included such areas as shallow depressions, old termite mounds, drainage channels, and





others. This sampling regime was chosen with a view of capturing as wide a range of the vegetation types and species in the area as possible (Gillison and Brewer 1985, Økland 1990, Austin and Heylingers 1989 in Wessels et al. 1998, de Blois et al. 2002).

The general vegetation type in each of the selected sites was characterised. This characterisation was based on the floristic and landscape features observed in the different habitat types. Dominant species of plants in the woody and herbaceous layers were identified and used for this purpose. The general terrain and proximity to important features, such as the lake or streams were noted. From this sampling, species of conservation concern and invasive species occurring in the area were identified and the geographical coordinates of their areas of occurrence recorded.

Specimens that could not be identified in the field were collected as vouchers for subsequent identification and deposition in the Makerere University Herbarium (MHU).

### 3.2.1.2 Invertebrates

Insects and other invertebrates dominate the terrestrial and aquatic ecosystems in terms of species richness, individual abundances and biomass (Wilson 1985, Stork 1988, Gaston, 1991). Their temporal and spatial distributions span the ranges occupied by many vertebrate and plant species, including finer-grained patch sizes and geographical distributions, more complex seasonal and successional sequences and patch dynamics with more rapid turnover (Gaston and Lawton 1988). Insects are also highly susceptible to the adverse effects of disturbance and land use change; this makes them useful as indicators of ecosystem change (Terborgh 1992).

Although insects and other invertebrates dominate the terrestrial and aquatic ecosystems, including the savannah and forest systems of the CHAA, and they are useful indicators, there are often severe limitations in terms of sampling and assessing the insect populations in biodiversity impact assessments. Of primary concern are the taxonomic restrictions for most invertebrate groups; specifically, the lack of a complete catalogue of knowledge of species, and the paucity of experts able to identify invertebrate taxa.

Although all species occurring in an area of interest are a component of overall ecological value, it is neither practicable, nor necessary, to assess potential effects of a project on every species that might be affected. This is particularly the case for most invertebrates, and lower plants (that is, bryophytes and pteridophytes), where, as mentioned, the taxonomy and ecology is often poorly established. As such, and in line with global conservation priority-setting, for this biodiversity impact assessment, vertebrates were largely used as a surrogate for all animal species, and vascular plants as a surrogate for all plants (Secretariat of the CBD, 2006). The selection was based upon a higher level of knowledge (ecology and conservation status) of these surrogates, and adopts the hypothesis that conditions which support restricted range vertebrates and/or vascular plants are likely to also support rare species of other taxonomic groups. Although this is an approximation of the likely situation, it provides manageable and meaningful conclusions.

Nevertheless, Uganda and the Albertine Graben are fortunate in that detailed taxonomy is available for some invertebrate groups. In particular, the butterflies (Order: Lepidoptera) and the dragonflies and damselflies (Order: Odonata).

Butterflies are known sensitive indicators of environmental change associated with natural and human-induced disturbances. Their populations are influenced by changes in local climatic conditions and the availability of host plants for larval and adult stages (Ehrlich et al. 1972, Thomas et al. 1998). Frequently disturbed environments are considered unstable and unpredictable and, as a result, have low species diversity, whereas less disturbed, more stable environments are expected to promote high species diversity (Odum 1985).

Dragonflies and damselflies utilise both aquatic and terrestrial habitats, and hence these groups can contribute greatly to the evaluation of environmental quality (Miller and Miller 2003). They are known to be very sensitive to structural habitat quality, and are used as indicator groups to evaluate landscape degradation.

Given their established taxonomy (*viz.*, Kielland (1990), Larsen (1991, 2005), Davenport (1996, 2003), Carder and Tindimubona (2002), Clausnitzer (2002), Miller and Miller (2003), and Picker et al. (2004), Molleman (2012), and the resources available at [www.africa-dragonfly.net](http://www.africa-dragonfly.net)), a baseline of butterflies, dragonflies and



damselflies was established for the LSA. The IUCN's RedList (IUCN 2014a) was consulted to establish a list of potential species with conservation significance, which informed the analysis of probability for these species to occur in the CHAA (see Section 3.3.3.2 Species of Concern).

Butterfly, dragonfly and damselfly surveys were carried out at 12 sites across the LSA in suitable habitat, including: locations corresponding to the various pieces of Project-related infrastructure on the Buhuka Flats, encompassing the stream and seasonally flooded wetland located near the airstrip, and the well pads; the lake shore of Lake Albert and the Bugoma Lagoon area; the lower reaches of the Masika River and the associated wetland; the escarpment; and along the pipeline route to the refinery area and Kabaale (which consists mainly of cultivated land, fairly degraded seasonal wetlands, riverine vegetation along Hohwa River and pockets of natural woodlands) (Appendix C, Table 2.1).

For butterflies, at each sampling site, survey methods employed included time- and distance-constrained sweep netting and baited traps (after Samways et al. 2010). Eighteen traps, baited with fermenting banana, were set along transect lines at each sampling location and left in place for two days. All specimens captured in the sweep nets and traps were identified in the field and released; only specimens with difficult identification were collected for further processing at Makerere University. Each of the butterfly species was assigned to one of the ecological categories as described by Davenport (1996); that is, forest-dependent species (F), forest edge/woodland species (f), open-habitat species (O), widespread species (W), migratory species (M), and wetland species (S).

Adult dragonflies and damselflies were sampled at each sampling site using time- and distance-constrained sweep netting (after Samways et al. 2010). In most instances, only mature males were sampled to minimise impacts on breeding populations. Familiar local species were recorded by observation only or by catch-and-release after confirmation. Voucher specimens were collected and preserved for further laboratory identification. Particular attention was given to the local habitat where the species were found.

### 3.2.1.3 Amphibians and Reptiles

A desktop search and review was made of available literature about the reptiles and amphibians of the CHAA. Data and information sources included: Plumptre et al. (2003, 2005, 2007); RPS (2006); AWE (2008a, b, c, 2013a, b, 2014a, b); AECOM (2012); EACL (2013, 2014), IUCN (2017); and GBIF (2017). The data thus obtained were used to identify sampling and survey sites on satellite imagery for later investigation during the field surveys. These data were also used to inform the assessment of probability of species of concern occurring in the CHAA (see Section 3.3.3.2 Species of Concern).

Nineteen survey sites were selected across the LSA as a representative sample of habitats that could be affected by the Project development (Appendix C, Tab. 3.1, Fig. 3.1). Eight of these were surveyed for amphibians and reptiles, five exclusively for amphibian fauna, and six exclusively for reptilian fauna. A control site within Bugoma Forest Reserve was selected, because several sites surveyed along the pipeline route, outside the forest, were considered to be analogous with the habitats present within the forest, yet were severely degraded and under heavy cultivation. The reptile and amphibian species composition of the Bugoma Forest Reserve was expected to be close to the original composition, while that along most of the pipeline and refinery areas could constitute a mixture of a few forest and grassland generalists.

Visual encounter and opportunistic survey approaches (after Heyer et al. 1994, and McDiarmid et al. 2012) were the main methods employed at each sampling site during the survey. These methods are well-tested and robust methods for surveying reptiles and amphibians. Visual encounter surveys are time constrained and are effective for most amphibians in most habitats. The data gathered using this method provides information on species richness of a habitat, with the best results for amphibians achieved in the evening between 7 pm and 9 pm, when most amphibian species expected to occur in the CHAA (refer to Channing and Howell 2006) would be active. Opportunistic surveys recorded those species outside of the systematic sampling locations and times.

Species estimators (as species accumulation curves) were used to calculate the possible maximum number of species that could occur in the LSA. Four estimators: Chao 1, Chao 2, Jackknife 1, and Jackknife 2 were used (after Gotelli and Colwell 2011).



### 3.2.1.4 Birds

Birds can represent a significant component of the biodiversity of an area, and they are ecologically versatile, representing herbivores, carnivores and omnivores, as such, they have been shown to be effective indicators of general biodiversity (Sutherland et al. 2004). They also lend themselves well to the identification of conservation priorities (Pain et al. 2005), and are a good indicator group for monitoring (Pearson 1994, Pearson and Carroll 1998).

A desktop search and review was made of available literature about the birds of the CHAA. Data and information sources included: Bennun and Njoroge (1996); Byaruhanga et al. (2001); Plumptre et al. (2003, 2007); RPS (2006); AWE (2008a, b, c, 2013a, b, 2014a, b); AECOM (2012); EACL (2013, 2014), IUCN (2017); and GBIF (2014; 2017). The data thus obtained were used to identify sampling and survey sites on satellite imagery for later investigation during the field surveys. These data were also used to inform the assessment of the probability that species of concern could occur in the CHAA (see Section 3.3.3.2 Species of Concern).

Birds, being highly mobile, tend to reflect the nature of larger areas rather than points within the landscape, except in instances where there are key nesting or roosting sites. Land birds and their habitats were generally surveyed along 2 km transects, each of ten 200 m sections (NatureUganda 2010). Each transect was predominantly within a single habitat, and collectively covered each of the main habitats in the LSA. Waterbirds and waders were recorded at fixed time-constrained, point-count sites (after Gregory et al. 2005). Data were recorded in a standard format, as used by Nature Uganda (2010) for the national bird monitoring programme.

All birds were identified by sight (or sound) in the field, with taxonomy following Stevenson and Fanshawe (2002).

### 3.2.1.5 Mammals

A desktop search and review was made of available literature about the mammal fauna of the CHAA. Data and information sources included: Kityo et al. (2003), Plumptre et al. (2003, 2005, 2007); RPS (2006); AWE (2008a, b, c, 2013a, b, 2014a, b); AECOM (2012); EACL (2013, 2014), IUCN (2017); and GBIF (2017). The data thus obtained were used to identify sampling and survey sites on satellite imagery for later investigation during the field surveys. These data were also used to inform the assessment of the probability that species of concern could occur in the CHAA (see Section 3.3.3.2 Species of Concern).

Standard survey methods, as described in Isabirye-Basuta and Kasenene (1987), Wilson et al. (1996), and Claustinitzer & Kityo (2001), were employed to sample the mammal fauna and their habitats in the LSA. Due to the paucity of large mammals in the LSA due to human disturbance (for example, see AWE (2008a, b, c, 2013a, b, 2014a, b)), intensive surveying was only conducted for small mammals.

In particular, rodents and insectivores (e.g. shrews) were surveyed using Sherman traps deployed in 11 trap lines of 40 traps each on the Buhuka Flats, and 80 traps each in the Bukona area. The trap lines were open for three to five days in the Buhuka Flats, two days in the Bukona area, and baited with a standard bait mixture (after Claustinitzer and Kityo (2001), Isabirye-Basuta and Kasenene (1987)). Traps were re-baited every evening, while checking, recovering and processing any captured animals completed in the morning. Bats were surveyed using mist nets, harp traps and acoustic methods (using the AnaBat II and SM2 bat detectors). Surveys started at dusk and continued until 10 pm on nights when this was possible. Surveys for large mammals were largely opportunistic, that is, tracks and signs (spoor or faecal material), observation, and informant interviews with local people. Informant interviews with local people were the main methods used along the pipeline route.

## 3.2.2 Aquatic Biodiversity

In addition to a detailed review of existing literature and databases, multi-season surveys of the aquatic ecology of fixed sampling locations within the LSA were completed to describe the baseline conditions. The field surveys focused on the near-shore zone of Lake Albert opposite the well pads (that is, Pad 1, Pad 2, Pad 3, Pad 4-2 and Pad 5, Appendix D, Figure 1), and the general Kingfisher Field Development area (that is, the jetty, the CPF, construction camp, spoil area, contractors' camp, parking lot, the existing camp, permanent camp, the airstrip, the borrow pit). Watercourses and wetlands of the Buhuka Flats were sampled, including



the Masika River (upper, mid and lower reaches and wetland), the Kamansinig River<sup>1</sup> (upper, mid (airfield wetland) and lower reaches, including the lagoon), and Well Pad 2 stream<sup>2</sup>.

Seasonal variation in aquatic ecology was assessed through multi-season surveys. As mentioned above, the region of the CHAA experiences two wet seasons (refer to Vol 4a Surface Water Specialist Report). The first field surveys captured the dry season, and occurred from 23 February to 8 March 2014. The second set of field surveys captured the wet season, and occurred from 23 to 28 May 2014. The third set of surveys captured the second wet season, and occurred from 15 to 20 November 2014.

The surveys focussed on Lake Albert and the watercourses and wetlands, and included:

- water quality;
- phytoplankton;
- zooplankton;
- macro-invertebrates; and
- fish.

The precise methods of survey for each of these groups are summarised below, and presented in detail in Appendix D.

### 3.2.2.1 *Water Quality*

Data for studies on lake water quality were collected at two fixed sites on each of five transects of 2 km length (from the shore lake-ward) opposite each of the five well pads. Samples from each transect were collected at about 10 m from the shoreline (inshore) and at the end of the transect, 2 km from shore (offshore).

In the Bugoma Lagoon, the samples were taken about 10 m from the shore and at a point approximately midway across the lagoon. Within the other wetlands associated with Kamansinig River, Masika River and Well Pad 2 stream, samples were collected in the water column.

For each transect and sampled site, the following physical characteristics were recorded: shoreline topography; soil type; vegetation cover; water depth, nature of bottom sediments and GPS location.

Water quality samples were collected using a 5 L van Dorn sampler from a depth of ~50 cm. In-situ and laboratory physical and chemical parameters were recorded at each sampling location. In-situ parameters included: dissolved oxygen ( $\text{mg.L}^{-1}$ ), temperature ( $^{\circ}\text{C}$ ), pH and conductivity ( $\mu\text{S.cm}^{-1}$ ) as measured using a Hach HQ40d Multiprobe. Laboratory-determined parameters included: ammonia-nitrogen and nitrate-nitrogen; ortho-phosphate; total phosphorus (TP); total nitrogen (TN); soluble reactive silica; chlorophyll a; and faecal coliform.

### 3.2.2.2 *Phytoplankton and Macrophytes*

Data for studies on phytoplankton were collected at two fixed sites on each of the five transects of 2 km length (from the shore lake-ward) opposite each of the five well pads. Samples from each transect were collected at about 10 m from the shoreline (inshore) and at the end of the transect, 2 km from shore (offshore).

In the Masika and Kamansinig Rivers, phytoplankton samples were collected mid-stream in the upper, mid and lower reaches. In the Bugoma Lagoon, which is in the lower reaches of the Kamansinig River, the samples were taken about 10 m from the shore and at a point approximately midway across the lagoon. Macrophytes were assessed in conjunction with the vegetation and flora assessments, and focused on major vegetation formations and the extent of surface water flows.

<sup>1</sup> This watercourse is referred to as the Airfield Stream in the baseline studies.

<sup>2</sup> This stream is not formally named on any maps; it drains off the escarpment and flows to the immediate south the





Twenty millilitres of water were sampled at each location from a depth of ~50 cm. Each sample was fixed with Lugol's solution (Utermöhl 1958), and stored away from light (Wetzel and Likens 2000). The sedimentation method of Utermöhl (1958) was used to count the phytoplankton under an inverted microscope, while taxonomic identification was made with the help of Komarek and Anagnostidis (1999) and John et al. (2002).

### 3.2.2.3 Zooplankton

Data for studies on zooplankton were collected at two fixed sites on each of five transects of 2 km length (from the shore lake-ward) opposite each of the five well pads. Samples from each transect were collected at about 10 m from the shoreline (inshore) and at the end of the transect, 2 km from shore (offshore).

In the Masika and Kamansinig Rivers, samples were collected mid-stream in the upper, mid and lower reaches. In the Bugoma Lagoon, which is in the lower reaches of the Kamansinig River, the samples were taken about 10 m from the shore and at a point approximately midway across the lagoon.

Vertical zooplankton hauls were taken from ~50 cm above the bottom sediments to the surface using a conical net of 0.25 m mouth opening and 60 µm mesh. Three hauls were taken to make a composite sample for each site, which was preserved with 4% sugar-formalin solution. In the laboratory, samples were identified based on published keys (*viz.*, Rutner-Kolisko 1974, Brooks 1957, Pennak 1953, Sars 1985).

### 3.2.2.4 Macro-invertebrates

A desktop search and review was made of available literature about the macro-invertebrate fauna of the CHAA. Data and information sources included: AECOM (2012), IUCN (2017); and GBIF (2014, 2017). The data thus obtained were used to guide sampling and survey sites on satellite imagery for later investigation during the field surveys. These data were also used to inform the assessment of the probability that species of concern could occur in the CHAA (see Section 3.3.3.2 Species of Concern).

Data for studies on macro-invertebrates were collected at two fixed sites on each of five transects of 2 km length (from the shore lake-ward) opposite each of the five well pads. Samples from each transect were collected at about 10 m from the shoreline (inshore) and at the end of the transect, 2 km from shore (offshore).

In the Masika and Kamansinig Rivers, samples were collected mid-stream in the upper, mid and lower reaches. In the Bugoma Lagoon, which is in the lower reaches of the Kamansinig River, the samples were taken about 10 m from the shore and at a point approximately midway across the lagoon.

In Lake Albert, at each sampling site, composite, triplicate sediment samples were collected using a Ponar grab sampler (after APHA 1992). The physical characteristics (for example, soft mud, sandy, stony, etc.) were noted for each composite sample. The samples were processed in the laboratory according to published methods (that is, APHA 1992, Ferraro and Cole 1992, Ochieng 2006, and Ochieng et al. 2008) in order to sort, identify and quantify the macro-invertebrates. Identification was undertaken based on Mandal-Barth (1954), Merrit and Cummins (1984), and de Moor et al. (2003).

Three indices, namely: EPT (Ephemeroptera (mayflies), Plecoptera (stoneflies) and Trichoptera (caddis flies)) taxa richness; total taxa richness; and abundance of macro-invertebrate individuals (after Wenn 2008, Oghenekaro 2011) were determined and used for a description of macro-invertebrate status as indicators of environmental quality. EPT taxa are considered the least tolerant of pollution (for example, organic pollution) and, therefore, aquatic environments with the presence of a high richness of EPT taxa, are regarded to be of good quality. Total taxa richness refers to the total number of all macro-invertebrate taxa (for example, genera) in a sample, with a higher taxa richness having better environmental quality. Based on the works of Wenn (2008), samples with: an EPT score of two to three, and total taxa score of ten to 12, are considered to be indicative of moderate to good environmental conditions; those with EPT scores of one, and total taxa scores between five and 10, were considered to be indicative of fair environmental conditions; while those with EPT scores of zero and varying total taxa scores of one to seven, were considered to be of poor environmental condition.





### 3.2.2.5 Fish

Data for studies on fish were collected at two fixed sites on each of five transects of 2 km length (from the shore lake-ward) opposite each of the five well pads. Samples from each transect were collected at about 10 m from the shoreline (inshore) and at the end of the transect, 2 km from shore (offshore).

In the Masika and Kamansing Rivers, samples were collected mid-stream in the upper, mid and lower reaches. In the Bugoma Lagoon, which is in the lower reaches of the Kamansing River, the samples were taken about 10 m from the shore and at a point approximately midway across the lagoon.

Data were obtained using multifilament gillnets, set perpendicular to the shore, ranging from 1 to 6 inches, increasing by ½ inch increments. Gillnets were set in the evening and hauled at dawn. On retrieval, fish species were identified using Greenwood (1966) and morphometrics (length, weight, and sexual maturity) recorded. Gut content and diet of individuals was determined in the laboratory. Species composition, relative abundance, population structure and relative condition were calculated from the catch statistics.

### 3.2.3 Overall Biodiversity Value

Biodiversity value is a term used by the IFC in PS6 *Biodiversity Conservation and Sustainable Management of Living Natural Resources* (IFC 2012a, 2012b), as well as by a number of other industry bodies; for example, The Energy and Biodiversity Initiative (EBI 2014), and the International Council on Mining and Metals (ICMM) (ICMM 2010). Those publications, together with the CBD's *Voluntary Guidelines on Biodiversity-Inclusive Impact Assessment* (Secretariat of the CBD, 2006), are recognised as standards of good practice for biodiversity impact assessment. Biodiversity values represent components of biodiversity at various levels of biological organisation, such as species or ecosystems that are important for conservation. Those values are reflected in CNOOC's corporate philosophy towards biodiversity conservation (see CNOOC 2014a, b).

To focus in on the key issues relating to biodiversity, as identified by stakeholders, the stakeholder's values placed on components of the biodiversity in the CHAA, and the potential risks posed to those biodiversity values by the Project, the concept of valued components was used. Valued components are physical, biological, economic, social, cultural, and health properties of the environment that are considered important by the proponent, public, government agencies, and/or the scientists involved in the assessment process (that is, the stakeholders) (Treweek 1999). These valued components identified and described during the baseline then become the focus for the impact assessment. In terms of the biodiversity of the CHAA, valued components are defined as the elements of an ecosystem that are identified as having scientific, social, cultural, economic, or aesthetic importance.

The biodiversity baseline concentrated on identifying two levels of biodiversity valued component in the CHAA, which are akin to the requirements of IFC PS6 (IFC 2012a) and the Secretariat of the CBD (CBD 2006); viz.: ecosystems and habitat and species of concern. It also describes the direct and indirect drivers of change to those values. In particular:

a) The ecosystem and habitats level:

This focussed on the broad description and identification of natural and modified terrestrial and aquatic ecosystems and habitat and critical habitat in the CHAA, and the direct and indirect drivers of change to those ecosystems and habitats at the spatial and temporal scale, and the processes and functions that drive those ecosystems and habitats; particularly, the integrity of those systems.

b) The species level:

The focus was on populations of species of concern; in particular, species of high value to people. These species included, amongst others: species of high conservation concern, as gazetted under Ugandan legislation, and the IUCN's Red List (IUCN 2014a); culturally important species; and invasive species.

The assessment of ecosystem services, often seen as a third component of biodiversity, is covered in Chapter 18 Ecosystem Services.



### 3.2.3.1 Ecosystems and Habitats of Concern

The intrinsic values of ecosystem functions and processes, and habitats, supported in the CHAA were assessed on two levels: ecosystem integrity and priority habitat (including critical habitat). The methods used for each of these are discussed below.

The assessment of ecosystem integrity broadly followed the guidance provided by the CBD's *Voluntary Guidelines on Biodiversity-Inclusive Impact Assessment* (Secretariat of the CBD 2006) and Treweek (1999). For the identification and assessment of priority habitats, natural and modified habitats were identified initially based on remote imagery, then verified with field data. For critical habitat, the IFC's approach was followed (refer to IFC PS6 (IFC 2012a, b)).

These guidance documents are complementary, and are recognised as a leading-practice approach for the assessment of impacts to biodiversity through a focus on the protection and conservation of biodiversity values of key conservation concern.

#### 3.2.3.1.1 Ecosystem Integrity

Ecological integrity refers to the abundance and distribution of species and the ecological patterns and processes that maintain biological diversity and ensure ecosystem resilience (Woodley et al. 1993). The major ecosystems and habitat types within the CHAA were initially identified at the desktop level based on the works by: RPS (2006); AWE (2008a, b, c, 2013a, b, 2014a, b); AECOM (2012); EACL (2013, 2014); as well as a land cover assessment undertaken based on SPOT6 data (Appendix E). Thereafter, those ecosystems and habitats were confirmed by field verification undertaken during the terrestrial and aquatic field studies (see Section 3.3.1 and 3.3.2 respectively).

The area of each ecosystem and habitat was determined based on the land cover assessment data built on the SPOT6 and aerial imagery data analysis, and the mapping of communities done during the terrestrial ecology field surveys. A quantitative and qualitative assessment of the integrity of each of the identified ecosystems and habitats was determined from field data (see Appendix C and D). For the purposes of this impact assessment, the integrity of the ecosystems was determined based on the following criteria (after: Kent and Coker 1992, Treweek 1999, Tucker 2005, Secretariat of the CBD 2006):

- **Composition**  
Diversity and complexity - what is there and how abundant (in a particular time frame) it is.
- **Structure (or pattern)**  
How biological units are organised in time and space. Ecosystem 'scale' refers to the space it occupies and the way it changes over time. The structure and interactions that shape the flow of energy and the distribution of biomass.
- **Linkages and corridors**  
To habitat of the same or different ecosystems, which provide an important 'playing field' for ecological processes and enable the goal of their persistence. These linkages are in contrast to a highly-fragmented landscape where patches of natural habitat are effectively isolated.
- **Key processes (including ecosystem function)**  
Which natural (that is, physical and/or biological) and/or human-induced processes are of key importance for the creation and/or maintenance of ecosystems. These are termed drivers of change, and include direct and indirect drivers. Examples of direct drivers include: changes in land use and land cover; fragmentation and isolation; extraction, harvest, or removal of species; external inputs such as emissions, effluents, chemicals; disturbance; introduction of invasive, alien and/or genetically modified species; and restoration. Examples of indirect drivers of change include: demographic; economic; socio-political; cultural; and technological processes or interventions.
- **Representativeness in the landscape.**



The uniqueness of the ecosystems within the CHAA and the wider landscape; this rarity factor is related to the concepts of irreplaceability and vulnerability. The concept of irreplaceability relates to rarity or uniqueness in the landscape, while vulnerability refers to degree of threat (for more detailed definitions, see Section 3.3.3.1.2).

### ■ Resilience and stability

The ability of the ecosystem to absorb change and persist, and maintain the same form.

Based on the assessment of these criteria, the condition of the ecosystems and habitats was estimated and assigned a subjective class, that is, pristine, near-pristine, slightly-degraded, moderately-degraded, heavily-degraded.

Two key drivers for this condition designation were: to aid in the identification of modified and natural habitat (see below), and the IFC's requirement of no net loss of natural habitats within the LSA (IFC 2012a); and the philosophy of net positive impact (NPI) in areas of critical habitat (IFC 2012a, see below). Where NPI is primarily measured in terms of quality hectares, which can be determined as a derivative of the area of an ecosystem multiplied by the condition of different habitat types.

### 3.2.3.2 *Species of Concern*

Although all species occurring in an area of interest are a component of overall biodiversity and ecological value, it is neither practicable, nor necessary, to assess potential effects of a project on every species that might be affected. This is particularly the case for most invertebrates and lower plants (that is, bryophytes and pteridophytes) where the taxonomy and ecology is often poorly established. As such, and in line with global conservation priority setting, terrestrial and aquatic vertebrates and selected invertebrates were used as a surrogate for all animal species, and vascular plants as a surrogate for all plants (Secretariat of the CBD, 2006). The selection was based upon a higher level of knowledge (that is, ecology and conservation status) of these surrogates, and adopts the hypothesis that conditions that support restricted-range vertebrates and selected invertebrates, and/or vascular plants, are also likely to support species of other taxonomic groups, including rare and threatened species. Although this is an approximation of the likely situation, it provides manageable and meaningful conclusions.

For the intents of this biodiversity impact assessment, a species of concern was defined as a plant or animal species that requires special conservation consideration based on certain characteristics, or one which may be particularly sensitive to Project effects. Those characteristics were then used to gauge the sensitivity of the particular species to the development, and how best to manage those sensitivities as part of the development.

The following selection criteria were used to screen and identify terrestrial and aquatic species of concern for the assessment, which are in line with the criteria for critical habitat designations (IFC 2012a):

- a) Threatened and restricted-range/endemic species (Criteria 1 and 2).
- b) Statutory species (national/international legislation, agreements, conventions) (Criteria 1, 2 and 3).
- c) Species of economic and/or cultural importance (Criteria 5 and 13).
- d) Convention on the International Trade in Endangered Species (CITES)-listed species (Criterion 1, 2 and 3).
- e) Evolutionarily distinct species (Criterion 5)
- f) Species that play a critical ecological role, represent guilds of species, or capture effects to other species with similar habitat requirements and sensitivities (Criterion 10).
- g) Invasive or potentially invasive species.

The determination of which level the species of concern was placed at, was used for determining the level of sensitivity of the particular species. A similar approach was used for assessing the sensitivity of the valued components. As an example,





- A species of concern with a moderate sensitivity could be one that is a regional endemic, a species whose distribution is significantly reduced from former extent but currently stable.
- A species of concern with a very high sensitivity may have an IUCN status of critically endangered or endangered, a local endemic, or its range is restricted to the CHAA. Or local temporal concentrations of individuals significant to global population, or much reduced and/or highly fragmented species distribution compared to its former extent, or ecosystem representation whose presence or processes support critically endangered or endangered species' habitat, or buffers it, keystone species, and/or species new to science.

The identification of a list of potential species of concern for the CHAA was determined from the species lists and known distribution records contained in: Emerton and Muramira (1999); AWE (2008a, b, 2013a, b, 2014a, b); Lamprey (2009); NEMA (2010); Plumptre et al. (2003, 2007, 2010, 2011); AECOM (2012, 2013); and the GBIF (2017), and the findings of the field surveys (see Appendix C and D). For those species where actual records of occurrence did not exist, yet were identified as potentially occurring in the area (based on habitat preferences and knowledge of the species), inferred distributions were derived from: Kalema and Beentje (2012) (plants); Mandahl-Barth (1954) (freshwater molluscs); Greenwood (1966) (fish); Carder and Tindimubona (2002), Davenport (2003) (butterflies); Miller and Miller (2003) (Odonata); Channing and Howell (2006) (amphibians); Spawls et al. (2004) (reptiles); Stevenson and Fanshawe (2002) (birds); and Butynski et al. (2013), Happold (2013), Happold and Happold (2013), Kingdon and Hoffman (2013a, b), Kingdon et al. (2013) (mammals).

It is recognised that some species of concern would not actually occur in the CHAA for various reasons, such as unsuitable habitat. Therefore, a screening of the probability of the various species of concern actually occurring in the CHAA was determined through a probability analysis based on:

- Knowledge and experience of the CHAA, and the wider area, as determined based on observations made during the KFDA Scoping Study.
- Findings of previous studies and published scientific literature.
- Species records stored in the GBIF (2017).
- Knowledge of the life histories of the species, habitat preferences, and known ecological requirements as determined through published information and information presented in the species profiles on the IUCN's Red List (IUCN 2017).
- Consultation with experts and professional judgement and experience of the assessors.

Three levels of probability were used: possible, probable and unlikely. These were defined as:

- c) Possible: the species may occur in the CHAA, or move through the CHAA (in the case of migratory and highly mobile species) due to potential habitat and/or resources.
- d) Probable: the species is likely to occur in the CHAA due to suitable habitat and resources being present, and/or known records from the CHAA.
- e) Unlikely: the species will not likely occur in the CHAA due to lack of suitable habitat and resources.

The probability assessment was used as the starting point for the screening of species of concern to occur in the CHAA as per the criteria set out above. Only those species with a possible and probable likelihood of occurrence within the CHAA were considered for inclusion in the valued component assessment. Nevertheless, the other species were not ignored; rather, it was assumed that the species chosen could act as proxies for many of the other species, should they occur, however remote the possibility. As such, a precautionary approach was adopted where there was an uncertainty that a species could potentially occur in the CHAA.



### 3.2.3.3 IFC Priority Habitat

Under the IFC's approach (see IFC 2012b), three classes of habitat are used to assign value to biodiversity: modified habitat; natural habitat; and critical habitat. Modified habitats are found in areas that have been altered by human activity and may contain large portions of non-native plants and animals. Examples include agricultural landscapes and reclaimed areas (IFC 2012b). Modified habitats may or may not retain ecological functions that support significant biodiversity value. Natural habitats are those where the species composition and primary ecological functions of the area have not been fundamentally altered by human activity (IFC 2012b). The definition of "fundamentally altered" is undertaken on a case-by-case basis; however, for the intents of this biodiversity impact assessment, natural habitats were defined as those habitats where the key processes, composition, and structure were largely intact. Critical habitats are a subset of either modified or natural habitats that constitute areas of significant importance for biodiversity conservation. The identification of natural, modified and critical habitat is discussed below.

Different mitigation standards are recognised for development occurring in each of the three habitat classes. Consequently, identifying the types of habitat that might be affected by a development project is a central aspect of understanding baseline conditions.

#### 3.2.3.3.1 Natural and Modified Habitat

As mentioned in Section 3.3.1, the identification of modified and natural habitats was initially based on secondary data sources using SPOT6 imagery as a basis. Although it is recognised that the modified and natural components of particular ecosystems and habitats within land cover classes cannot be accurately determined directly from spectral classes, using SPOT6 data does provide a very good indication of modified and natural habitats. For more information on the approach and methods for the determination of land cover, see Appendix E. Following that initial land cover assessment, each land cover class was assigned to the natural or modified categories, as per the IFC's criteria (see IFC 2012b), as relevant to the CHAA, and based on initial observations made during the scoping visit. Areas within protected and managed areas (for example, Central Forest Reserves) were automatically assumed to be natural habitat, as were the near-shore (that is, within 1 km of the shore) environments of Lake Albert. These classes were then refined using the data collected as part of the terrestrial and aquatic ecology baseline studies (see Appendix C and D).

#### 3.2.3.3.2 Critical Habitat

The identification and assessment of critical habitat followed the approach defined by the IFC (refer to IFC PS6 (IFC, 2012a, b)). The IFC's PS6 uses the concept of critical habitat as an important means to identify biodiversity values of key conservation concern. The purpose of defining critical habitat is to identify areas of a particularly sensitive nature that deserve special attention for avoidance and may require supplementary mitigations, including offsetting.

The IFC's critical habitat concept considers and expands on a variety of pre-existing ideas and definitions of priority sites for biodiversity conservation; for example: Key Biodiversity Areas (IUCN 2010); Endemic Bird Areas and Important Bird Areas (BirdLife International 1998); Alliance for Zero Extinction sites (AZE 2010); World Heritage Sites (UNESCO 2014); and Ramsar Convention on wetlands of international importance (Ramsar 2014). This approach is also supported by a broad array of conservation organisations, and is increasingly accepted and applied by a variety of private companies (as developers) and financial institutions (the Equator Principles Banks as lenders).

As mentioned, critical habitats are a subset of either modified or natural habitats that constitute areas of significant importance for biodiversity conservation. For a development to occur in a critical habitat:

- No other viable alternatives within the region exist for development of the project on modified or natural habitats that are not critical.
- The project does not lead to measurable, irreversible and adverse impacts on those biodiversity values for which the critical habitat was designated, and on the ecological process supporting those biodiversity values.



- The project does not lead to a net reduction in the global and/or national/regional population of any species of concern, for which critical habitat was identified in the CHAA, over a reasonable period of time.
- A robust, appropriately-designed, and long-term biodiversity monitoring and evaluation programme is integrated into the project’s management programme to achieve net gain (or NPI).

If these conditions can be met, it is reasonable that a mitigation strategy can be designed to achieve net gains, over a reasonable period of time, for the biodiversity values for which the critical habitat has been designated; that is, a no net loss in natural habitat and NPI philosophy in critical habitat. Consequently, projects proposed in areas containing critical habitat face challenges not faced by projects in natural or modified habitats that are not classified as critical.

Critical habitat was identified by delineating spatial units of analysis (DMUs), screening biodiversity features (that is, at the species, ecosystem and landscape scales), and evaluating the distribution of critical habitat in the CHAA. For a detailed discussion of the methods and approach used, see Appendix B.

### 3.3 Impact Assessment

The impact assessment process identifies the magnitude of a particular impact from the project and then compares that magnitude with the sensitivity of the receiving environment to derive an overall significance for the impact. This method relies on a detailed description of both the impact and the biodiversity valued component that is the receptor. The magnitude of an impact depends on its characteristics, which may include such factors as its duration, reversibility, area of extent, and nature in terms of whether positive, negative, direct, indirect or cumulative.

#### 3.3.1 Key Questions and Indicators

One of the main purposes of an impact assessment is to provide answers to questions that people have about how a project could affect something that matters to them, such as a valued component. To focus this assessment and ensure that the impact assessment clearly addressed the key issues raised by the stakeholders (see Section 3.1), and the objectives set for this impact assessment (see Section 2.1), questions were formulated that captured the concerns relative to a particular issue. In this report, those concerns are expressed as ‘key questions’, and they form the basis of the investigations of potential effects and impacts of the Project.

Two key questions were established:

- 1) **What effect could the Project have on habitats and ecosystem integrity?**
- 2) **What effect could the Project have on species of concern?**

Under each of these key questions, sub-questions were developed that focused on the specific phases of the Project, in particular, the construction, operation and decommissioning phase.

Indicators for the impact assessment were selected to assess the level of potential impact. Changes to the indicators were analysed to determine the effectiveness of the mitigation hierarchy, and identify Project constraints and opportunities for additional avoidance and mitigation. Indicators and context for impact assessment are outlined for the key questions and associated valued components in Table 1.

**Table 1: Key questions and indicators**

Key Question	Valued Component	Indicator	Indicator Description	Context
What effect could the Project have on habitats and	Ecosystem integrity Priority habitat	■ regional representative -ness	The uniqueness of an ecosystem or habitat in the CHAA and wider landscape	The persistence of species of concern in the CHAA and wider area.





Key Question	Valued Component	Indicator	Indicator Description	Context
ecosystem integrity?		<ul style="list-style-type: none"> <li>changes in soil, water flows and quality, and vegetation</li> </ul>	Drivers of change affecting key processes	<p>Maintain the distribution and abundance of species of concern such that self-sustaining and ecologically effective populations can be maintained.</p> <p>Achieve net gains for species of concern for which critical habitat is affected by the proposed Project, and at least no net loss for all other valued components.</p> <p>Maintenance of ecosystem processes and functions and connectivity.</p>
		<ul style="list-style-type: none"> <li>ecosystem composition</li> </ul>	The diversity and complexity of an ecosystem or habitat – species composition and abundance	
		<ul style="list-style-type: none"> <li>ecosystem configuration</li> </ul>	The structure or pattern of an ecosystem, including linkages and corridors	
What effect could the Project have on species of concern?	All species of concern	<ul style="list-style-type: none"> <li>habitat quantity and quality</li> </ul>	The extent and integrity of preferred foraging and breeding habitat	
		<ul style="list-style-type: none"> <li>habitat connectivity</li> </ul>	Connectivity to adjacent areas of suitable habitat and potential for dispersal	
		<ul style="list-style-type: none"> <li>abundance and distribution</li> </ul>	Expected vs actual population numbers and distribution	
		<ul style="list-style-type: none"> <li>survival and reproduction</li> </ul>	Likelihood of continued survival/reproduction compared to baseline	

Valued components were assessed in the context of maintaining self-sustaining and ecologically effective populations (or sub-populations) (in the case of species of concern), or a functioning and resilient ecosystem, at the scale of the CHAA. A self-sustaining population is one that will be maintained into the future with a low risk of loss (that is, extirpation). Long-term population persistence is the outcome of maintaining self-sustaining, ecologically effective populations, and population persistence is frequently applied as a conservation target by conservation biologists and resource managers (Ruggiero et al. 1994, With and Crist 1995, Fahrig 2001, Nicholson et al. 2006).

Maintaining self-sustaining and ecologically effective populations often will result in the protection of the ecological services humans benefit from when ecosystems are functional. Such benefits include the continued opportunity for consumptive and non-consumptive use of non-timber forest products by people that value those resources as part of their culture and livelihood (Hooper et al. 2005). For more information on the assessment of impacts to ecosystem services, see Chapter 18 Ecosystem Services.

### 3.3.2 Impact Assessment

Key questions were answered using a two-step process. Firstly, an overall written analysis, also known as a reasoned narrative, identified the indicators used for the assessment, and walks the reader through the logic of the assessment and the conclusions reached. Secondly, a formal impact classification was applied.

For a detailed explanation of the approach and method used for the reasoned narrative and the impact classifications, see Appendix F.





### 3.4 Limitations

This biodiversity impact assessment has been undertaken based on historical data and field surveys focusing on the current extent of the proposed Project. Any subsequent design changes and or alterations may require new surveys to be conducted (for example, if infrastructure layout is changed and extended beyond the current CHAA). Additionally, this biodiversity impact assessment should be read with the following limitations in mind:

- In a few instances, access limited the areas that could be sampled during the terrestrial ecology field campaigns. The escarpment area, for example, was extremely steep in many places and it was not practical to access some of the natural habitats.
- Comprehensive field trapping and sampling programmes were limited in many instances due to theft and/or vandalism of traps. Therefore, multiple-trap-night surveys were not always able to be employed, nor were trapping regimes for reptiles and small mammals.
- A thorough assessment of the migratory species of bird in the CHAA has not yet been completed. A survey needs to be completed in late September or October to assess the importance of the area for migrants.
- The selection of species of interest for the impact assessment was based on the level of knowledge (that is, ecology and conservation status) of the species to act as surrogates for all species in the area, and adopts the hypothesis that conditions which support restricted range vertebrates and/or vascular plants are likely to also support rare species of other taxonomic groups.

Despite these limitations to baseline data, the conclusions contained within this report are based upon a robust and transparent procedure, and represent an accurate evaluation and assessment of likely impacts.

Your attention is also drawn to Appendix A of this report for further limitations.

## 4.0 RELEVANT LEGISLATION

This chapter presents a summary of Ugandan national legislation, associated regulations and policies that are pertinent to biodiversity, to assist and guide the ESIA. It describes international conventions and regional frameworks to which Uganda is a signatory; and also includes a summary of the international standards and guidelines that represent good industry practise, to which CNOOC wishes to adhere. Publicly available documents, and reports supplied by CNOOC were used to compile this review.

### 4.1 Uganda's Policy, Legal and Institutional Framework on Biodiversity

#### 4.1.1 The Constitution of the Republic of Uganda (1995)

The over-arching government policy on natural resource conservation in Uganda is provided for in the Constitution of the Republic of Uganda. The relevant constitutional provisions in the National Objectives and Directive Principles of State Policy include the following:

- **Principles of State Policy XXVII (iv):** mandates the State (both central and local government) to create and develop parks, reserves and recreational areas, and to ensure conservation and promote the rational use of natural resources so as to safeguard and protect the bio diversity of Uganda.
- **Article 237 (2) (b):** the Government or local government, as determined by Parliament by law, shall hold in trust for the people and protect, natural lakes, rivers, wetlands, forest reserves, game reserves, national parks and any land, to be reserved for ecological and touristic purposes for the common good of all citizens.
- **Article 245:** the utilisation of natural resources of Uganda shall be undertaken in such a way as to meet the development and environmental needs of present and future generations of Ugandans and, in particular, the State shall take all possible measures to prevent or minimise damage and destruction to land, air and water resources resulting from pollution and other causes.





### 4.1.2 Uganda Wildlife Bill (2017)

The primary objectives of the Uganda Wildlife Bill are to provide for the conservation and sustainable management of wildlife, to strengthen wildlife conservation and management; to continue the Uganda Wildlife Authority; and to streamline roles and responsibilities for institutions involved in wildlife conservation and management.

The Bill re-aligns the Uganda Wildlife Act Cap. 200 with the 2014 Uganda Wildlife Policy, the Oil and Gas policy and laws, the Land use policy and law, the National Environment Act, the Uganda Wildlife Education Centre Act, the Uganda Wildlife Research and Training Institute Act and all other laws of Uganda and developments which came into force after the enactment of the Uganda Wildlife Act in 1996.

For the first time, nationally-protected species were declared in the 2017 Uganda Wildlife Bill. Wildlife species listed in the Third Schedule of Act V are protected species in Uganda, in addition Act V states that wildlife species protected under any international convention or treaty to which Uganda is a part (and to which the regulations set out in section 86 applies), are protected species.

#### 4.1.2.1 Uganda Wildlife Act (1996)

The Uganda Wildlife Act defines two types of conservation areas: “wildlife protected” and “wildlife managed” areas. Although the Act *made provision for* the declaration of protected species, no protected species were declared in the Act.

##### **Wildlife Protected Areas**

- **National Park:** these are protected areas of international and national importance because of their biological diversity, landscape or national heritage, and in which biodiversity conservation, recreation, scenic viewing, scientific research and other economic activity may be permitted.
- **Wildlife Reserve:** these are protected areas of importance for wildlife conservation and management and in which conservation of biological diversity, scenic viewing, recreation, scientific research, and regulated extractive utilisation of natural resources are permitted.

##### **Wildlife Management Areas**

- **Community Wildlife Areas:** these are wildlife management areas where wildlife is protected, whilst taking into account the continued use of the land and the sustainable exploitation of wildlife in the area by people and communities ordinarily residing there. Sustainable exploitation of the natural resources of the area, including by mining and other methods, is permitted - providing that it is in a manner compatible with the continued presence of wildlife in the area.

#### 4.1.2.2 Uganda Wildlife Act Cap 200 of 2000

The Uganda Wildlife Act cap 200 of 2000 was enacted by an Act of Parliament to provide for sustainable management of wildlife (UWA 2014). The Act consolidated wildlife management law in Uganda and established the Uganda Wildlife Authority (UWA) as the responsible authority for wildlife management and conservation, and enforcement of wildlife laws and regulations. The Act covers all wildlife protected areas (PAs) and wildlife outside PAs, and specifically mandates UWA to control and monitor industrial and mining developments in wildlife protected areas.

#### 4.1.2.3 Uganda Wildlife Policy (2014)

The Uganda Wildlife Policy generally promotes long-term conservation of wildlife and biodiversity in a cost-effective manner, which maximises the benefits to the people of Uganda in terms of ecology, economy, aesthetics, science and education. The policy aims at achieving this through promoting conservation and sustainable utilisation of wildlife throughout Uganda. The policy seeks to exclude industrial development, including mineral exploration and extraction, from wildlife protected areas (that is, national parks and wildlife reserves).





The Ugandan Government resolved to review Uganda's Wildlife Policy, to harmonise it with related instruments like the National Environment Policy, the Wetland Policy and the Constitution, in the form of the 2014 Uganda Wildlife Policy. New aspects incorporated in the policy included:

- To provide for incentives that supports the private sector to invest more in wildlife development in Uganda.
- To guarantee safety for tourists by enhancing security in national parks and game reserves, under the expanded anti-terror surveillance in Uganda.
- To increase resource allocation to the tourism sector, specifically for extending and improving infrastructure to, within and around tourism sites.
- To reconcile the needs for wildlife conservation and human beings, particularly in areas that have been affected by insurgency and civil strife.
- To ensure that any infrastructural development within and around wildlife conservation areas does not compromise the support eco-systems for flora and fauna in the respective areas.
- Demands for land in national parks will not be entertained, except in very exceptional circumstances where survival of communities is involved.

The 2014 edition of the Policy (first formulated in 1994) updates the Policy of 1999. The Policy guides the conservation and development of wildlife resources in Uganda. The policy requires that new developments and interventions with potential to affect wildlife resources are subject to environmental impact assessments.

#### **4.1.2.4 Uganda Forestry Policy (2001) and the National Forestry and Tree Planting Act (2003)**

The Forestry Policy is implemented through the National Forestry and Tree Planting Act (2003). The Act provides for:

- The conservation, sustainable management and development of forests.
- The declaration of forest reserves for the purposes of protection and production of forests and forest produce.
- The sustainable use of forest resources and enhancement of productive capacity of the forests.
- The promotion of tree planning.
- Consolidation of the law relating to the forestry sector and trade in forest produce.

Parts of Uganda's permanent forest estate carry dual status as National Parks, Wildlife Reserves and Animal Sanctuaries; such areas are subject to additional regulations under the Uganda Wildlife Act (1996).

Declared forest reserve categories include Central Forest Reserves (CFRs), Local Forest Reserves, Community Forests, private forests, and forests forming part of a wildlife conservation area (declared under the Uganda Wildlife Act, Cap 200).

CFRs fall in two main categories, namely those designated for production and those for protection. Such forest reserves are subsequently managed in a manner consistent with the purpose for which they were declared:

- **Production forests:** includes savanna bushland and grassland areas - reserved for supply of forest products and future development of industrial plantations.
- **Protection forests:** includes all the tropical high forests, savanna woodlands and/or grasslands – reserved forests include those that protect watersheds and water catchments, biodiversity, ecosystems and landscapes that are prone to degradation under uncontrolled human use.

CFRs are held in trust for the people of Uganda and managed by the National Forestry Authority (NFA) and are classified according to the following categories:





- Site of special scientific interest;
- Strict nature reserve;
- Joint management forest reserve;
- Recreation forest for purposes of eco-tourism; and
- Any other area, for a purpose prescribed in the order.

In a forest reserve, it is prohibited to cut, disturb, damage, burn or destroy any forest produce, remove or receive any forest produce, or undertake activities not consistent with the specific management plan except under conditions set out in the Act or in accordance with a licence granted under the Act. The Act also makes provision for classification of trees as reserved/protected and therefore subject to specific controls. In addition, Section 38 of the Act requires that an environmental impact assessment be undertaken for any project or any activity which may, or is likely to have a significant impact on a forest.

### **4.1.2.5 The Land Act (1995)**

Section 43 of the Land Act provides for management and utilisation of land in accordance with the Uganda Wildlife Act, and other laws. Section 44 (i) mandates the government or local governments to protect national parks, wetlands and forest reserves (amongst others) for ecological and tourism purposes, and hold these in trust for the people of Uganda.

### **4.1.2.6 Uganda National Land Policy (2013)**

The Uganda National Land Policy makes provisions in relation to natural resource management and biodiversity. These include Government resolutions to ensure that land use practises conform to land use plans, and that the principles of sound environmental management including biodiversity preservation, soil and water protection, conservation and sustainable land management are applied. The policy commits the Government to take measures including to

- Provision of special protection for 'fragile' ecosystems (that is, unique and sensitive biodiversity features).
- Development of harmonised criteria for gazetting and de-gazetting conservation areas.
- Establishment and implementation of effective mechanisms for management of wildlife outside protected areas.
- Incentivise community participation in conservation on privately-owned land and co-management of conservation on public land.
- Regulate the use of hilltops and other sensitive ecosystems.
- Develop mechanisms to resolve human-wildlife conflict.

### **4.1.3 Uganda National Biodiversity Strategy and Action Plan (2015-2025)**

Published by the Ugandan National Environmental Management Authority (NEMA) in October 2016, the National Biodiversity Strategy and Action Plan (NBSAP) provides a framework to guide the setting of conservation priorities, channelling of investments and building of the necessary capacity for the conservation and sustainable use of biodiversity in the country.

The overarching principles of the NBSAP are:

- a) Sustainable development and environmental sustainability
- b) Mainstreaming of biodiversity conservation, sustainable use of biological resources and equitable sharing of benefits from biological resources into existing policy, legislative, institutional and development frameworks as appropriate;





- c) Stakeholder participation in the development and implementation of biodiversity strategy and action plans;
- d) Awareness creation, education, training and capacity building at local, national and institutional levels to enhance effective participation and implementation of biodiversity measures;
- e) Recognition, promotion and upholding of traditional and indigenous knowledge of biological resources and sustainable resource management and where benefits arise from the use of this knowledge;
- f) Engagement and collaboration with international partners to enhance conservation and sustainable use of Uganda's biological diversity;
- g) Integrated implementation of Multi-Lateral Environmental Agreements;
- h) Equal consideration of the three objectives of the Convention on Biological Diversity – conservation; sustainable use; and benefit sharing arising from the use of biological resources

The Uganda NPSAP is a useful policy guide for addressing Uganda's concerns in biodiversity conservation and the utilisation of its components, as well as for implementation of the requirements of the Convention on Biological Diversity.

#### 4.1.4 The National Policy for the Conservation and Management of Wetland Resources, 1995

Prepared in 1995 by the Ministry of Natural Resources, the policy aims at "promoting the conservation of Uganda's wetlands in order to sustain their ecological and socio-economic functions for the present and future well-being of the people." The policy also complements the Ramsar Convention on Wetlands of International Importance especially as Waterfowl Habitat, and provides for the requirement of EIA for all planned wetland developments.

#### 4.1.5 The National Fisheries Policy, 2004

The Policy provides strategies for sustainable management of fisheries through decentralisation and community involvement.

### 4.2 Conventions and International Agreements

Uganda is a signatory to the following international conventions and agreements:

- Convention on Biological Diversity: Under the convention, each contracting party is expected to develop national strategies, plans or programs for the conservation and sustainable use of Biological diversity.
- Convention on International Trade in Endangered Species (CITES).
- Convention on the Conservation of Migratory Species of Wild Animals, (the Bonn Convention).
  - African-Eurasian Water-bird Agreement (AEWA).
  - International Gorilla Agreement (Uganda is in the process of acceding to this agreement – it has been signed but is not yet ratified).
- Convention on Wetlands of International Importance (the Ramsar Convention).
- UNESCO World Heritage Commission.
- Lusaka Agreement on the Cooperative Enforcement Operations Directed against Illegal trade in Fauna.

### 4.3 International Guidance

#### 4.3.1 International Finance Corporation's Performance Standards

At the project financing level, the management of biodiversity is addressed by PS6: *Biodiversity Conservation and Sustainable Management of Living Natural Resources*. PS6, and the associated GN6 relates to:





- The protection and conservation of biodiversity.
- Maintenance of ecosystem services.
- Sustainable management of living natural resources.

The requirements set out in PS6 have been guided by the Convention on Biological Diversity. PS6's main priority is that the Project should seek to avoid impacts on biodiversity and ecosystem services. When avoidance of impacts is not possible, measures to minimise impacts and restore biodiversity and ecosystem services should be implemented.

However, when a project occurs in critical habitat supporting exceptional biodiversity value, a net gain in biodiversity value is required.

PS6 sets specific biodiversity protection and conservation standards relating to potential project impact. The specific requirements are separated according to the following categories:

- **Modified Habitat:** areas that may contain a large proportion of plant and/or animal species of non-native origin, and/or where human activity has substantially modified an area's primary ecological functions and species composition. PS6 relates to areas of modified habitat that have significant biodiversity value, and requires that impacts on such biodiversity must be *minimised, and mitigation measures implemented* as appropriate.
- **Natural Habitat:** viable assemblages of plant and/or animal species of largely native origin, and/or where human activity has not essentially modified an area's primary ecological functions and species composition. In such areas, the conservation outcome required by PS6 is *no-net-loss of biodiversity value* achieved using the "like-for-like" or better principle of biodiversity offsets, where feasible.
- **Critical Habitat:** areas with high biodiversity value, including (i) habitat of significant importance to Critically Endangered and/or Endangered species; (ii) habitat of significant importance to endemic and/or restricted-range species; (iii) habitat supporting globally significant concentrations of migratory species and/or congregatory species; (iv) highly threatened and/or unique ecosystems; and/or (v) areas associated with key evolutionary processes. When a project occurs in critical habitat supporting exceptional biodiversity value, a *net gain in biodiversity value* is required by PS6. This is achievable through appropriate biodiversity offsets.
- **Legally Protected and Internationally Recognised Areas:** such areas often have high biodiversity value; when this is the case these areas are likely to qualify as critical habitat and, as such, the conservation outcome required by PS6 is also a *net gain in biodiversity value*, as well as obtaining the relevant legal permits, following standard governmental regulatory procedures, and engagement of affected communities and other stakeholders.
- **Invasive Alien Species:** the development project should not intentionally introduce any new alien species (unless carried out within the appropriate regulatory permits) and should not deliberate any alien species with a high risk of invasive behaviour under any circumstance. PS6 requires that any introduction of alien species be the subject of a *risk assessment* for potential invasive behaviour, and that the project should *implement measures to avoid* the potential for accidental or unintended introductions.
- **Management of Ecosystem Services:** where a project is likely to adversely impact ecosystem services, an *ecosystem service review to identify priority ecosystem services* is required. Priority ecosystem services are (i) those services on which project operations are most likely to have an impact and, therefore, which result in adverse impacts to Affected Communities; and/or (ii) those services on which the project is directly dependent for its operations (for example, water). If adverse impacts on Priority ecosystem services are unavoidable, these must be *minimised and mitigation measures* that aim to maintain the value and functionality of priority services implemented. With respect to impacts on priority ecosystem services on which the project depends, *impacts on ecosystem services should be minimised and measures that increase resource efficiency* of their operations implemented. For a full assessment of ecosystem services, see.



### 4.3.2 Regional Frameworks

At the regional level, Uganda is a member of the **African Union** and one of its objectives is to promote sustainable development at the economic, social and cultural level.

In the East African region, Uganda is obliged to implement the articles of the Treaty for the establishment of the **East African Community**, which it ratified together with other member states in 2000. In article 119, Partner States agreed to promote close cooperation in culture and sports.

Uganda is signatory to the **Nile Basin Initiative** (NBI). The NBI was established in 1999 by the Nile basin countries, to oversee the implementation of the Nile River Basin Action Plan. This process is still ongoing; once concluded, the resulting agreement will supersede all the existing Nile water agreements, pending establishment of a permanent legal and institutional framework for the Nile Basin.

Uganda is also a member of the **Lake Victoria Basin Commission** (LVBC). Releases from Lake Victoria have a controlling role on the water balance and level of Lake Albert, and hence on the flows within the Victoria and Albert Niles. The LBVC was established by the East African Community as a mechanism for coordinating the various interventions on the Lake and its Basin. The LBVC also serves as a centre for promotion of investments and information sharing among the various stakeholders.

## 5.0 KEY ISSUES RELATING TO BIODIVERSITY

As identified through stakeholder consultation, review of background biodiversity and environmental reports, published ecological literature, and consideration of the IFC's Environmental, Health, and Safety Guidelines for Onshore Oil and Gas Development (IFC 2007b) and the performance standards (that is, IFC 2012a, b and c), broadly, the key issues pertaining to the CHAA's biodiversity include:

- Habitats and ecosystems
  - Construction and operation of the pipeline and the potential effects that the construction and operation could have on wetlands, streams, forests and reforestation areas (including potential critical habitat), agricultural areas, and soils.
  - Construction and operation of the wells and CPF on the environment of the Buhuka Flats and Lake Albert.
  - The potential effects the construction and operation of the Project could have on Lake Albert, wetlands and environment of the Buhuka Flats. These include: vibration; pollution (oil, erosion and sedimentation, other run-off, effects to groundwater); increased fishing pressure from in-migration; long-term damage to the lake ecosystem.
  - Potential effects to the Bugoma Central Forest Reserve due to upgrade of the existing road, including the possible need for offsets.
  - Potential effects the construction and operation of the Project could have on the corridors connecting the wild areas along Lake Albert from Semliki to Murchison Falls National Park.
- Species of concern
  - Concern for the loss of animal species from the Buhuka Flats. Potential effects to the populations of Hippopotamus (*Hippopotamus amphibius*), Nile Crocodile (*Crocodylus niloticus*), and Grey Crowned Crane (*Balearica regulorum*), amongst others.
  - The identification of migratory and threatened species inhabiting the CHAA.
  - Potential effects to the populations of Chimpanzees (*Pan troglodytes*), Nahan's Francolin (*Ptilopachus nahani*) and African Elephant (*Loxodonta africana*) in Bugoma Central Forest Reserve.

In summary, the main issues related to potential effects to the biodiversity of the CHAA from the construction and operation of the Project relate to the to changes in ecosystem composition (for example, species





composition), configuration (for example, patch size and connectivity) and function of the wider CHAA through the direct loss, disturbance or change in condition of natural and modified habitats, including critical habitat.

## 6.0 BASELINE ENVIRONMENT

This section describes the baseline biodiversity environment of the LSA and CHAA. It draws upon existing, published information, local knowledge and comprehensive, multi-season field surveys. The detailed baseline study reports for the terrestrial and aquatic ecology are presented in Appendix C and D respectively.

### 6.1 Terrestrial Biodiversity

This section focuses on describing the baseline terrestrial biodiversity of the LSA. It summarises the findings reported in the terrestrial ecology baseline report, as presented in Appendix C.

#### 6.1.1 Vegetation Communities and Flora Species

A summary of the baseline of the vegetation communities and flora species of the CHAA are presented based on the findings of the desktop study and the field investigations of the LSA.

##### 6.1.1.1 Vegetation Communities

According to Langdale-Brown et al. (1964), the CHAA is mapped as supporting the following vegetation communities:

- a) Dry *Hyparrhenia* Grass Savanna, with undifferentiated deciduous Thicket (Q3/V1) on the Buhuka Flats.
- b) *Themeda-Chloris* Grass Savanna (Q4) on the Buhuka Flats.
- c) Dry *Combretum-Hyparrhenia* Savanna (N2) on the escarpment.
- d) Moist *Combretum-Terminalia-Albizia-Hyparrhenia rufa* Savanna/Medium Altitude Forest/Savanna Mosaic (K/F2) beyond the escarpment, towards Bugoma Central Forest Reserve along the pipeline route.
- e) Moist *Combretum-Terminalia-Albizia-Hyparrhenia rufa* Savanna (K) beyond the escarpment, towards Bugoma Central Forest Reserve along the pipeline route.
- f) *Cynometra-Celtis* Medium Altitude Moist Semi-deciduous Forest (D2) beyond the escarpment, towards Bugoma Central Forest Reserve along the pipeline route.

It is noted that the delineation of these communities is 50 years old. Since that time, large tracts of vegetation have been altered in the CHAA, in particular, along the pipeline route, and, therefore, strict alignment with Langdale-Brown et al.'s (1964) classification was not possible. These areas have been subjected to high-intensity, subsistence agriculture, which has altered much of the original natural landscape (Forest Department 2002). These drivers of change, together with widespread cattle grazing and charcoal manufacture, have put pressure on the natural vegetation communities in the CHAA as compared to Langdale-Brown et al.'s (1964) original work. This is particularly noticeable in the areas on the escarpment, between the escarpment proper and the Bugoma Central Forest Reserve (see Figure 4). These areas have, for the most part, been converted to subsistence agricultural fields.

Nevertheless, significant natural vegetation still exists long the escarpment (Figure 5). This corridor extends from the areas south of the CHAA, northwards toward the Kabwoya Wildlife Reserve and the Kaiso-Tonya Community Wildlife Area. Indeed, Plumptre et al. (2007) identified these corridors to be part of an important linkage from the Semliki/Toro Wildlife Reserve in the south, the Budongo–Bugoma–Kagombe–Itwara Forest Reserves, right through to the Murchison Falls National Park in the north.

The field surveys (Appendix C) identified seven broad vegetation communities within the LSA (as depicted in Figure 5 and Figure 6):

- a) Wooded Grassland;
- b) Woodland;



- c) Thicket-Grassland Mosaic;
- d) Open Grassland;
- e) Bushed Grassland;
- f) Bushland and Shrubland; and
- g) Wetlands (including permanent wetlands of *Phragmites*, *Typha*, and *Cyperus*, and seasonally flooded grassland (floodplains) of *Sporobolus pyramidalis* and *Cynodon dactylon*).

These communities broadly align with those described by Langdale-Brown et al. (1964); however, their current distributions are different to those originally described, primarily due to the increased pressures from agriculture and human disturbance over the last 50 years. The characteristics and condition of each of these communities is summarised below.





# BIODIVERSITY IMPACT ASSESSMENT




Figure 4: Land cover and land use in the CHAA





# BIODIVERSITY IMPACT ASSESSMENT




Figure 5: Vegetation communities of the CHAA (focus on Buhuka Flats)





# BIODIVERSITY IMPACT ASSESSMENT




Figure 6: Vegetation communities of the CHAA





#### 6.1.1.1.1 Wooded Grassland



Figure 7: Typical wooded grassland in the CHAA

Aligns with the Dry *Combretum-Hyparrhenia* Savanna (N2) of Langdale-Brown et al. (1964); the ground layer in these communities is dominated by grasses, while woody plants comprise 10% to 50% of the upper storey structure (Langdale-Brown et al. 1964). In the LSA, the ground layer is dominated by *Hyparrhenia rufa* and *Sporobolus pyramidalis* (Figure 7). The woody plant fraction was dominated by *Rhus natalensis*, *Combretum collinum*, *Acacia hockii* and *Annona senegalensis*.

These communities comprise about 3.35% of the CHAA, and included the sub-communities of: wooded grassland (WGI); open wooded grassland (OWGI); and dense wooded grassland (DWGI) (Figure 4a). These communities are largely restricted to the escarpment and beyond. The original extent of these communities on the plateau above the escarpment appears to have been significantly reduced in the last 50 years due to subsistence agricultural practices. On the escarpment, clearing for agricultural fields, and grazing by cattle are the primary drivers of change, together with frequent fires.



### 6.1.1.1.2 Woodland



Figure 8: Typical woodland along a watercourse in the CHAA

Aligns with the *Cynometra-Celtis* Medium Altitude Moist Semi-deciduous Forest (D2) of Langdale-Brown et al. (1964); these communities are characterised by an upper storey canopy layer that does not interlock and remains open, which allows for the growth of herbaceous vegetation; however, the herbaceous layer lacks a multi-layered structure (Langdale-Brown et al. 1964).

Within the CHAA, these communities were mostly dominated by *Acacia* species, while some were mixed with *Crateva* sp. and *Acalypha* sp. (Figure 7). The most abundant species in these communities were typically *Tamarindus indica*, *Rhus natalensis* and *Acacia sieberiana*, while the understory was typically dominated by *Grewia trichocarpa* and *Azima tetraacantha*. *Hypoestes forskalii*, *Panicum deustum* and *Sporobolus pyramidalis* were common herbaceous layer species.

These communities comprise about 0.70% of the CHAA, and included the sub-communities of: open woodland (OWo); and riverine woodland (RiWo) (Figure 8), and they tend to be restricted to the riverine areas along the mainly seasonal watercourses. Harvesting of large trees for the manufacture of charcoal is a noticeable driver in these communities, as well as trampling and grazing by cattle.



### 6.1.1.1.3 Thicket-Grassland Mosaic



Figure 9: Typical Thicket-grassland mosaic community in the LSA

Aligns with the Dry *Hyparrhenia* Grass Savanna, with undifferentiated deciduous Thicket (Q3/V1) of Langdale-Brown et al. (1964); these communities are characterised by thickets of 2 to 4 m height, with a dominance of much-branched, thorny, woody species that form dense clumps or continuous thickets sometimes approaching 100% cover (Langdale-Brown et al. 1964).

Within the CHAA, particularly on the Buhuka Flats, these thickets are interspersed with grassland, forming a mosaic (Figure 9). The most abundant species were *Acacia brevispica*, *Acalypha fruticosa*, *Azima tetraacantha*, *Euphorbia candelabrum*, *Cadaba farinosa* and *Dichrostachys cinerea*. Common species in the herbaceous and ground layer include *Sansevieria* spp., *Sporobolus pyramidalis*, *Cynodon dactylon*, *Aloe* sp. and *Cissus oliveri*.

These communities comprise about 1.37% of the CHAA, and included the sub-communities of: grassland with thicket (GIWT), which is largely restricted to the Buhuka Flats (Figure 5). Within this area, noticeable effects of heavy cattle grazing were observed, which appears to be the primary driver for maintaining the mosaic nature of this community, which also limits bush encroachment.



### 6.1.1.1.4 Open Grassland



Figure 10: Typical grassland community of the LSA

Aligns with the *Themeda-Chloris* Grass Savanna (Q4) of Langdale-Brown et al. (1964); these communities are dominated by a grass layer, with woody species typically constituting less than 5% (Langdale-Brown et al. 1964).

Within the CHAA, the dominant grass species in these communities included *Sporobolus pyramidalis*, *Pennisetum purpureum*, *Imperata cylindrica*, *Cynodon dactylon* and *Panicum maximum* (Figure 10). Occasional woody species included *Acacia polyacantha* subsp. *campylacantha* and *Vernonia amygdalina*. Often, these grasslands supported some thicket communities.

These communities comprise about 0.71% of the CHAA, and included the sub-communities of: open grassland (OGI) (Figure 5) and were largely restricted to the Buhuka Flats. Livestock grazing is by far the most dominant driver affecting the structure of the community, keeping it at very low stature.

### 6.1.1.1.5 Modified Habitats

Most areas of modified habitat occur along the proposed Feeder pipeline route. These areas are modified from their original, natural habitat - having been converted to cultivation. Patches of natural vegetation occur within these modified areas, although most have been largely altered from their original state. These remnants show affinities to the various vegetation communities identified in the wider area, including woodland, bushland, wooded grassland and wetland.



### 6.1.1.2 Flora Species

Few studies have focused specifically on the flora of the CHAA. Those studies that do exist tend to focus on protected areas and forest reserves. For example: Plumptre et al. (2009), in a study of Kabwoya Wildlife Reserve, which is about 30 km north of Buhuka Flats and the same ecoregion as the CHAA, identified 167 flora species in that reserve; NEMA (2010) identifies the Bugoma Central Forest Reserve as a particularly species rich area; while Kalema (2005) compiled a list of species for the Semliki Wildlife Reserve, which is about 40 km south of the CHAA. These studies do provide useful references for species assemblages of the wider area, and formed a good basis upon which to develop a list of potential species for the CHAA. These lists, together with those provided by the IUCN (2017), formed the basis of the identification of threatened species occurring in the CHAA.

#### Species Richness, Diversity and Abundance

In the area of the Buhuka Flats, the most abundant species were *Cynodon dactylon*, *Sporobolus pyramidalis*, *Acalypha fruticosa*, *Phragmites kirkii*, *Capparis erythrocarpos*, *Senna* sp., *Asparagus africanus*, *Cissus oliveri*, *Typha capensis*, *Cyperus articulatus* and *Dichrostachys cinerea*.

On the escarpment, where the soil conditions are more marginal and fragile, the most abundant species were *Acalypha fruticosa*, *Rhus natalensis*, *Hypoestes forskalii*, *Terminalia brownii*, *Acacia brevispica*, *Cissus oliveri*, *Sporobolus pyramidalis* and *Enteropogon macrostachyus*.

Beyond the escarpment, on the plateau, soil conditions (including drainage) were better, with only localised areas of water-logging. Within these areas, however, there is a strong element of human influence in the form of agricultural activities. Here, *Pennisetum purpureum*, *Acacia polyacantha*, *Imperata cylindrica*, *Vernonia amygdalina*, *Panicum maximum*, *Combretum collinum*, *Acanthus polystachyus* were the most abundant species.

*Sporobolus pyramidalis* had the highest relative abundance across all the LSA and all communities sampled. This was followed by *Acalypha fruticosa*, *Cynodon dactylon*, *Panicum maximum*, *Vernonia amygdalina* and *Rhus natalensis*. This indicates that grasses form a substantial proportion of all the vegetation communities within the LSA.

Ninety-four species were uncommon in the LSA. These include *Cordia africana*, *C. millenii*, *Cynometra alexandrii*, *Pterygota mildbraedii*, and *Markhamia lutea*. These are all good timber tree species, which are under intense pressure from logging. *Cordia millenii* is listed under Uganda's National Forestry Authority as a Reserved Species (Kalema and Bleentje 2012), and is, therefore, flagged for protection owing to excessive felling for its high-grade timber. None of the other species are listed under Ugandan legislation, or the IUCN's Red List (IUCN 2017). These species are widespread in the region and Africa (Kalema and Beentje 2012).

Overall, 96 families and 635 species were recorded in the LSA (Appendix C). Although the highest species richness was recorded in bushland (369 species), woodland (318 species) and wetland communities (301 species), with thicket-grassland and bushed grassland communities recording 91 and 202 species respectively, the number of species recorded was strongly correlated with sampling effort. In particular, the latter two communities had less sampling effort applied than the aforementioned communities (see Appendix C). A better measure was the mean number of species per survey plot, which identified bushed grassland (13.47 spp./plot), wooded grassland (8.89 spp./plot) and woodland (6.91 spp./plot) to have the highest species richness, while the lowest was wetland (5.02 spp./plot) and open grassland (5.79 spp./plot).

The general observation from these findings is that the more wooded vegetation communities recorded higher species richness per sampling unit. Accordingly, the least wooded communities of wetland and open grassland had the lowest species richness per sampling unit.

Across the LSA, woody species contributed 38.6% of species richness as compared to 61.4% for the non-woody species (Appendix C). On the Buhuka Flats, herbs and shrubs dominated; while on the escarpment, shrubs dominated; beyond the escarpment, herbs and trees dominated.





### Species of Concern

Four species of conservation interest were recorded in the LSA. These were the: *Milicia excelsa* (Mvule Tree) (listed as Lower Risk/Near Threatened by the IUCN, and a restricted species on the Ugandan list of Reserved Tree Species, as promulgated under Uganda's *National Forestry and Tree Planting Act 2003*); *Tamarindus indica* (Tamarind Tree) (Not Evaluated by the IUCN, yet a restricted species on the Ugandan list of Reserved Tree Species, as promulgated under Uganda's *National Forestry and Tree Planting Act 2003*); *Cordia millenii* (Drum Tree) (listed as Lower Risk/Least Concern by the IUCN); and the CITES Appendix II-listed *Euphorbia candelabra* (Candelabra Tree) and *Aloe* sp. (Aloe) (both Not Evaluated by the IUCN).

Five invasive species were recorded in the LSA, although they were uncommon. *Mimosa pigra* (Giant Sensitive Tree), *Lantana camara* (Lantana), and *Eichhornia crassipes* (Water Hyacinth) were the commonest species recorded, predominantly on the Buhuka Flats and the shore of Lake Albert (Appendix C). These species are recognised as some of most noxious weeds in the world (Lowe et al. 2000).

Other invasive species recorded included: *Pistia stratiotes* (Water Lettuce) (in a wetland community on the Buhuka Flats, where it was locally abundant); *Parkinsonia aculeata* (Parkinsonia) (recorded on the Buhuka Flats in open grassland); and *Ricinus communis* (Castor Oil Plant) (recorded from Wetland and Woodland communities).

For further discussion and assessment of these species, see Section 7.2.

### 6.1.2 Invertebrates

As mentioned, the sampling of terrestrial invertebrates was limited to butterflies, dragonflies and damselflies for the reasons discussed. Summarised below are the findings of the baseline surveys presented in Appendix C.

#### 6.1.2.1 Butterflies

A summary of the baseline of the butterfly species of the LSA is presented, based on the findings of the desktop study and the field investigations. The detailed baseline studies are presented in Appendix C.

### Species Richness, Diversity and Abundance

One-hundred-and-fifty-five species of butterfly were recorded in the LSA. Of these: 38 are forest-dependent species, including one forest highland species; 27 forest edge/woodland species; 25 migrant species; 20 open habitat species; 42 widespread species; and two wetland-dependent species (Appendix C). Based on their ecological preferences, 27.3% of the butterfly species recorded were those that are typically widespread; 13% were typical of open habitats; 16.2% were migrants; 17.5 % were forest edge/woodland species; 24.7% were characteristic of forest habitats; and 1.3% were wetland dependent.

The habitat mosaic of the escarpment area was the most species rich, along with the Kamansinig River, and the Hohwa River along the pipeline route, as well as the Kibale-Butoole area.

### Species of Concern

No Albertine Rift endemic species were recorded in the LSA. Only four of the species recorded have been evaluated by the IUCN, and are all listed as Least Concern; these were:

- Jeffry's Bush-brown (*Bicyclus jefferyi*)

This widespread species favours forest clearings and edges of relatively wet forests, and although it is common, its population trend is unknown (Larsen 2011a).

- Small Grass Yellow (*Eurema brigitta*)

This species is one of the most common butterflies of Africa and the Oriental region, thus having a vast EOO (Larsen 2011b). It is believed to have a stable population trend and favours a wide variety of savanna and grassland habitats.

- Dark Blue Pansy (*Junonia oenone*)



This species occurs across the entire African continent, and, although its population trend is currently unknown, there are no present threats to its global population (Larsen 2011c).

### ■ Dark Grass Blue (*Zizina antanossa*)

This species is one of the most widely distributed butterflies in Africa. It inhabits grassy, open areas in savannah, and disturbed areas of forest. Its population trend is unknown, but it is not known to be affected by any major threats at present (Larsen 2011d).

For further discussion and assessment of other species that may potentially occur in the CHAA, see Section 7.2.

### Habitats

Although no IUCN-listed, or Uganda-listed threatened species were recorded in the LSA, it does support a rich diversity of species. Habitats of importance for butterflies within the LSA, based on species richness and diversity, were the watercourses draining off the escarpment and along the pipeline route, the vegetation communities of the escarpment, and forest. It is noted that the forest patches may represent relicts of the original habitat in the wider area. However, it is noted too that the majority of species recorded are habitat generalists or ecotone species, and are not dependent upon intact habitat for their survival. Nevertheless, forest dependent species did constitute ~25% of the species recorded, and ~1.5% were wetland dependent. Therefore, certainly for butterflies, the array of habitats supported in the LSA is important to maintain species diversity.

For further discussion and assessment of habitats, please refer to Section 7.1.

### 6.1.2.2 Dragonflies and Damselflies

A summary of the baseline of the dragonfly and damselfly species of the LSA is presented based on the findings of the desktop study and the field investigations. The detailed baseline studies are presented in Appendix C.

### Species Richness, Diversity and Abundance

Forty-six species of dragonflies and damselflies were recorded from the LSA. The seasonally flooded wetlands along the Kamansinig River and the permanent wetlands of Masika River were the most species rich.

### Species of Concern

All the species recorded have been assessed by the IUCN, and all are categorized as being of Least Concern, with stable or unknown population trends. One species, the Common Riverjack (*Mesocnemis singularis*), although listed as Least Concern, is recommended by the IUCN for further monitoring due to possible declining population trends (Clausnitzer et al. 2010).

For further discussion and assessment of other species that may potentially occur in the CHAA, see Section 6.3.2.

### Habitats

Although no IUCN-listed, or Uganda-listed threatened species were recorded in the LSA, it does support a rich diversity of species. Habitats of importance for dragonflies and damselflies within the LSA, based on species richness and diversity, were the seasonally flooded and permanent wetlands on the Buhuka Flats and along the pipeline route.

For further discussion and assessment of habitats, please refer to Section 7.1.

### 6.1.3 Reptiles and Amphibians

A summary of the baseline of the reptile and amphibian species of the LSA is presented based on the findings of the desktop study and the field investigations. The detailed baseline studies are presented in Appendix C.



### 6.1.3.1 Amphibians

#### Species Richness, Diversity and Abundance

Twenty-three amphibian species were recorded from the LSA; this is between 76% and 100% of the expected species in the CHAA (Appendix C). These represent seven families and ten genera. The most species rich sites for amphibians were along the pipeline route: Kabakete, near the proposed Kabaale refinery site, had 12 species; and Zorobe had eight species. On the Buhuka Flats, the wetland on the lower reaches of the Masika River had eight species, with the wetlands of the Kamansinig River having up to seven species.

The most common species was a Ridged Frog (*Ptychadena* sp.<sup>13</sup>), which was recorded at 42% of the sampling sites. This was followed by the Cinnamon-bellied Reed Frog (*Hyperolius cinnamomeoventris*) at 37% of the sites, the Kivu Reed Frog (*H. kivuensis*), and the Crowned Bullfrog (*Hoplobatrachus occipitalis*) at 32% of sites, while the Banded Banana Frog (*Afraxalus fulvovittatus*), Common Toad (*Amietophrynus regularis*) and Anchieta's Ridged Frog (*Ptychadena anchietae*) were found at 26% of the sites.

#### Species of Concern

All the species recorded are listed as Least Concern by the IUCN, with the majority believed to have stable population trends, or unknown trends, except one. The Lake Victoria Toad (*Amietophrynus vittatus*) is listed as Data Deficient by the IUCN, with an unknown population trend (IUCN SSC Amphibian Specialist Group 2014a). This species was recorded on the Buhuka Flats in the seasonally flooded wetlands associated with the Kamansinig River (Appendix C, AWE 2008a, 2008b, 2013a). DeSaeger's River Frog (*Amietia desaegeri*), although listed as Least Concern by the IUCN, is a range restricted species (IUCN SSC Amphibian Specialist Group 2014b).

For further discussion and assessment of this species, see Section 7.2.

#### Habitats

Habitats of importance for amphibians within the LSA, based on species richness and diversity, were the seasonally flooded and permanent wetlands on the Buhuka Flats and along the pipeline route, and watercourses draining off the escarpment.

For further discussion and assessment of habitats, please refer to Section 7.1.

### 6.1.3.2 Reptiles

#### Species Richness, Diversity and Abundance

Twenty-one reptilian species, belonging to eight families and 11 genera, were recorded in the LSA (Appendix C), which could account for ~70% of the species in the CHAA. Generally, however, the reptilian diversity for each sampling site was poor, which may be an artefact of the sampling effort (see Section 3.5). The most diverse site was the area where the CPF will be located with five species, followed by Kasoga/Buhumuro-Nsanga on the pipeline route (four species), Masika River and wetlands associated with the Kamansinig River, each with three species. The rest of the sites had one, two or no reptiles recorded at them.

The most common species were the Speckle-Lipped Skink (*Trachylepis maculilabris*), recorded at 56% of the sites, followed by the Ground Agama (*Agama agama*) at 44% of the sites, and the Tree Agama (*Acanthocercus atricolis*) and the Striped Skink (*Trachylepis striata*) at 33% of the sites.

#### Species of Concern

The majority of species recorded were of Least Concern or Not Evaluated by the IUCN, and tended to be common species in the area. Four species, the Nile Crocodile (*Crocodylus niloticus*), Nile Monitor (*Varanus niloticus*), Smooth Chameleon (*Chamaeleo laevigatus*), and Graceful Chameleon (*C. gracilis*) are listed under CITES Appendix II (UNEP-WCMC 2018).

<sup>3</sup> This species is noted to not be new to science; however, it is difficult to separate from other species. Therefore, a specimen was sent to the museum for determination, the results of which were not available at the time of writing this report (M. Behangana, pers. comm.)





The Nile Soft-shelled Turtle (*Trionyx triunguis*) is known from Lake Albert, however it was not recorded from the LSA during baseline data gathering surveys. Although the species has been assessed as Vulnerable by the IUCN (2017), and populations in central and north-eastern Africa are understood to be stable (van Dijk et. al., 2017), it is facing pressures from human exploitation within Lake Albert, where adults and eggs are hunted for food and medicinal purposes; the carapace can fetch a high price in the markets of Kampala (Appendix C).

For further discussion and assessment of these and other species that may potentially occur in the CHAA, see Section 7.2.

### Habitats

Habitats of importance for reptiles within the LSA, based on species richness and diversity, were the seasonally flooded and permanent wetlands on the Buhuka Flats and along the pipeline route, and wooded grasslands. The wetlands associated with Lake Albert's shoreline (in particular, the lagoon, and the lower reaches of the Masika River) are important breeding and nursery areas for the Nile Soft-shelled Turtle and the Nile Crocodile. The ravines associated with the watercourses draining off the escarpment are important habitats for a variety of reptiles. Importantly, the heterogeneity of habitats in the wider LSA is important for maintaining reptile diversity.

For further discussion and assessment of habitats, please refer to Section 7.1.

### 6.1.4 Birds

A summary of the baseline of the bird species of the LSA is presented based on the findings of the desktop study and the field investigations. The detailed baseline studies are presented in Appendix C.

#### Species Richness, Diversity and Abundance

Two-hundred-and-eighty-three species were recorded in the LSA; mostly composed of species typical of the area. However, no forest specialists were recorded, and only a few forest generalists, reflecting the almost total loss of the original forest cover of the escarpment and the land above it. The seasonally flooded and permanent wetlands of the Buhuka Flats supported a wide variety of waterbirds, while grassland species were well-represented in the open areas. Interestingly, the diversity of aerial feeder species (like martins, swifts and their kin) was low, but the number of individuals was high.

The richness of species along the pipeline route was less than the flats; however, the diversity was still quite high, with 29 species recorded that were unique to that area.

Fifty-five species of waders and waterbirds were recorded in the LSA, primarily from the shore of Lake Albert and the wetlands on the Buhuka Flats. Notable species include: the first record of the Terek Sandpiper (*Xenus cinereus*) from Lake Albert; the second record of the Lesser Sandplover (*Charadrius mongolus*) for Uganda.

The species richness and abundance of individuals increased as expected during the peak migratory period of September/October. Some Palearctic migratory stragglers were still present in February-March in some numbers. Although very large numbers of migratory species were not recorded, it is conceivable that Lake Albert as a whole could support significant numbers. For example, a 1 km count along the shores of the lake, south from the jetty, produced over 400 birds of 27 species. Given this, and the fact that the lake is ~180 km long, it is conceivable that the shores of Lake Albert could support as many as 100,000 birds, just on the Ugandan side. The number of these species and individuals had significantly dropped during the May-June survey indicating that the majority of individuals had migrated away from the lake. Therefore, it can be expected that during the peak migratory period, the numbers of birds in the area will be substantial.

Interestingly, 16 raptor species were recorded. The Albertine Rift is a known migratory route for raptors, yet all the species recorded were residents.

### Species of Concern

The majority of species recorded in the LSA were of Least Concern status, and tended to be common species in the area. Two species listed as Endangered by the IUCN were recorded; these were Grey Crowned Crane (*Balearica regulorum*) and White-backed Vulture (*Gyps africanus*).



Grey Crowned Crane was recorded on the Buhuka Flats, where up to 14 individuals were regularly seen. (Appendix C). Indications were that breeding pairs were beginning to form at the end of May. Twenty White-backed Vultures were seen overflying the LSA in February.

Twelve regionally listed species were recorded in the LSA. These included: the vulnerable Martial Eagle (*Polemaetus bellicosus*), African Skimmer (*Rynchops flavirostris*), Grosbeak Weaver (*Amblyospiza albifrons*), Saddle-billed Stork (*Ephippiorhynchus senegalensis*), Great White Egret (*Ardea alba*); and the near threatened Purple Heron (*Ardea purpurea*), Grey Heron (*Ardea cinerea*), Goliath Heron (*Ardea goliath*), Brown Snake-eagle (*Brown Snake-eagle*), African Marsh Harrier (*Circus ranivorus*), Black-bellied Firefinch (*Lagonosticta rara*), and Vieillot's Black Weaver (*Ploceus nigerrimus*). Interestingly, individuals of most of these species were encountered throughout the entire LSA; that is, the Buhuka Flats, the escarpment and the pipeline route.

Other species of interest included the east African endemics: Spotted-flanked Barbet (*Tricholaema lacrymosa*), White-headed Saw-wing (*Psolidoprocne albiceps*), Black-lored Babbler (*Turdoides sharpie*), Red-chested Sunbird (*Cinnyris erythrocerus*), Baglafaecht's Weaver (*Ploceus baglafaectii*), Red-headed Quelea (*Quelea erythrops*), and Grey-headed Oliveback (*Nesocharis capistrata*).

Palaearctic migratory species were more abundant during the dry season and the second wet season (corresponding to the peak migratory period of September/October), when 39 species were recorded. These included: Black-winged Stilt (*Himantopus himantopus*); Common Ringed Plover (*Charadrius hiaticula*); Lesser Sandplover (*Charadrius mongolus*); Little Stint (*Calidris minuta*); Ruff (*Philomachus pugnax*); Common Snipe (*Gallinago gallinago*), Common Greenshank (*Tringa nebularia*), Green Sandpiper (*T. ochropus*); Wood Sandpiper (*T. glareola*); Terek Sandpiper (*Xenus cinereus*); Common Sandpiper (*Actitis hypoleucos*); White-winged Tern (*Chlidonias leucopterus*); Eurasian Reed Warbler (*Acrocephalus scirpaceus*); Great Reed Warbler (*A. arundinaceus*); Willow Warbler (*Phylloscopus trochilus*); Garden Warbler (*Sylvia borin*); Black Stork (*Ciconia nigra*); Black Kite (*Milvus migrans*); Eurasian Marsh Harrier (*Circus aeruginosus*); European Honey Buzzard (*Pernis apivorus*); Lesser Spotted Eagle (*Aquila pomarina*); Tawny Eagle (*Aq. rapax*); Booted Eagle (*Hieraaetus pennatus*); Great Spotted Cuckoo (*Clamator glandarius*); Blue-cheeked Bee-eater (*Merops persicus*); Eurasian Bee-eater (*M. apiaster*); African Hoopoe (*Upupa epops*); Sand Martin (*Riparia riparia*); Eurasian Swallow (*Hirundo rustica*); Yellow Wagtail (*Motacilla flava*); Red-throated Pipit (*Anthus cervinus*); Whitchat (*Saxicola rubetra*); Northern Wheatear (*Oenanthe oenanthe*); Isabelline Wheatear (*O. isabellina*); Pied Wheatear (*O. pleschanka*); Woodchat Shrike (*Lanius senator*); Spotted Flycatcher (*Muscicapa striata*); Semi-collared Flycatcher (*Ficedula semitorquata*); and Nightingale (*Luscinia megarhynchos*). Overall, numbers of both Palaearctic and Afrotropical species were quite high; however, the majority of waterbirds are resident in Uganda, making only local movements in response to rainfall.

For further discussion and assessment of these and other species that may potentially occur in the CHAA, see Section 7.2.

### Habitats

The birds recorded in the LSA represent guilds that are closely tied to the various habitats of the area, as well as many generalist species. For example, woody vegetation associated with wooded grassland and woodland communities was important for a variety of tree-dependent species; while the more open areas are important for species such as Temminck's Courser (*Cursorius temminckii*), Grey Crowned Crane (*Balearica regulorum*) and a variety of plovers and lapwings, whilst the seasonally flooded and permanent swamps, support a wide variety of bird species.

Nevertheless, habitats of importance for birds within the LSA, based on species richness and diversity, included: the ecotonal habitat at the foot of the escarpment and the escarpment face; the permanent wetlands associated with the Masika and Kamansing Rivers; and the shore of Lake Albert. The proposed pipeline route passes through predominantly agricultural country. Although largely disturbed and modified, these habitats are, nevertheless, species rich.

For further discussion and assessment of habitats, please refer to 7.1.



### 6.1.5 Mammals

A summary of the baseline of the mammal species of the LSA is presented based on the findings of the desktop study and the field investigations. The detailed baseline studies are presented in Appendix C.

#### Species Richness, Diversity and Abundance

Generally, the LSA supports a depauperate community and populations of medium to large-sized mammals, presumably due to the strong influence of human disturbance and associated pressures (*viz.*, AWE 2008a, b, 2013a, 2014a, EAC 2013, 2014). The mammal fauna assemblage is dominated by small to medium-sized species such as rodents, shrews, bats, mongoose, rabbits, duiker, bushbuck and primates.

Thirteen species of non-volent small mammals (that is, rodents and shrews) were recorded on the Buhuka Flats and escarpment area, which comprised 11 species of rodent and two shrews. Those areas surveyed along the pipeline route only recorded two species of rodent. These low figures for species richness and diversity may be an artefact of the sampling effort (see Section 3.4). The Pygmy Mouse (*Mus minutoides*) was the most abundant and common, followed by the Multimammate Mouse (*Mastomys natalensis*).

Two species of fruit bat (Sub-order: Megachiroptera), the Ethiopian Epauletted Fruit Bat (*Epomophorus labiatus*) and Peter's Dwarf Epauletted Fruit Bat (*Micropteropus pusillus*) were recorded from Buhuka Flats. Sixteen confirmed species of insectivorous bats (Sub-order: Mirochiroptera) were recorded from the Buhuka Flats, with possibly another two unconfirmed species. These represent the majority of species expected to occur in the CHAA.

The most common and abundant species were the Yellow House Bat (*Scotophilus dingani*), the Dark-winged Lesser House Bat (*Scotoecus hirundo*) and the Banana Pipistrelle Bat (*Neoromicia nanus*) appeared to occur in survey areas at higher levels of activity.

Thirteen medium-sized mammals were recorded in the LSA. These species tended to be habitat generalists, and included four primates: Vervet Monkey (*Cercopithecus aethiops*), Red-tailed Monkey (*C. ascanius*), Colobus Monkey (*Colobus guereza*) and Olive Baboon (*Papio anubis*); three viverrids: African Civet (*Civetta civetictis*), African Palm Civet (*Nandinia binotata*) and Small-spotted Genet (*Geneta tigrina*); a herpestid: the Marsh Mongoose (*Atilax paludinosus*); three rodents: Lesser Cane-rat (*Thryonomys gregorianus*), Bunyoro Rabbit (*Poelagus marjorita*) and Stripped Ground Squirrel (*Xerus erythropus*); and two bovids: Bush Duiker (*Sylvicapra grimmia*), and Bushbuck (*Tragelaphus scriptus*). These species appear to be sparsely distributed throughout the LSA. Of interest though is the relatively high diversity of carnivores and omnivores. The populations of all the species recorded within the LSA are unknown.

A small population of Hippopotamus occurs on the Buhuka Flats. No accurate counts of the number of individuals occur, however, it is doubtful that more than five animals exist in this isolated population (pers. obs.).

#### Species of Concern

As discussed, the majority of species recorded were cosmopolitan, generalist species with a Least Concern status and stable populations, as assessed by the IUCN. The notable exception being the small population of Hippopotamus on the Buhuka Flats; this species is listed as Vulnerable by the IUCN, with a decreasing population (Lewinson and Oliver 2008). Within Uganda, this species is recognised as having a restricted distribution, although it is locally abundant; as such, it is fully protected under the law (Lewinson and Oliver 2008).

Five species of bats recorded in the LSA are potentially cave or cavity roosting species. These include the Little Free-tailed Bat (*Chaerephon pumila*), Angolan Free-tailed Bat (*Mops condylura*), a Bent-wing Bat (*Miniopterus* sp.), the Dark-winged Lesser House Bat and Yellow House Bat. Although no caves were identified within the LSA that could house colonies of these species, there is a possibility that such features could exist, particularly on the escarpment. Of most interest is the Bent-Wing Bat. This species is listed in CITES Appendix II (UNEP-WCMC 2018), and is known to roost in caves housing hundreds or even thousands of individuals (Monadjem et al. 2010).



For further discussion and assessment of these and other species that may potentially occur in the CHAA, see Section 6.3.2.

### Habitats

As mentioned, the majority of the species recorded in the LSA are habitat generalists and quite wide ranging. There are obvious exceptions. The Bugoma Lagoon area and swamps of the lower reaches of the Masika River are important daytime refuges for the local Hippopotamuses, along with many other small and medium-sized mammals. Similarly, the seasonally flooded wetlands of the upper reaches of the Masika River and its tributaries, and the ones associated with the Kamansinig River, as well as those along the pipeline route, are important habitats for the small mammal assemblages in the LSA.

The escarpment is an important habitat for all the mammal species recorded in the LSA. It not only forms a continuous corridor along the length of Lake Albert, but it also offers important refugial sites for many of the small mammal species; notably the cavity and cave roosting bat species. The escarpment is dissected by numerous watercourses draining off the plateau; these form incised ravines in the escarpment that have the potential to support significant roosting sites for bats.

The thicket communities on the Buhuka Flats, and remnant vegetation patches along the pipeline route form important refuges and resource areas for small mammals.

For further discussion and assessment of habitats, please refer to Section 7.2.

## 6.2 Aquatic Biodiversity

This section focuses on describing the baseline aquatic biodiversity of the LSA and CHAA. It summarises the findings reported in the aquatic ecology baseline reports, as presented in Appendix D.

### 6.2.1 Water Quality

The water quality characteristics, as relevant to aquatic ecology, of the near-shore environment of Lake Albert, the Masika River, the Kamansinig River and the Well Pad 2 Stream are summarised below.

#### General Parameters – Lake Albert

Generally, the water quality parameters assessed for Lake Albert fell within the accepted limits of Uganda's national drinking water standards (the Ugandan standards), as published by the National Water and Sewerage Cooperation (NEMA 1996). Generally, the water quality at all sites sampled in Lake Albert was suitable for fisheries productivity and maintenance of other aquatic biodiversity (Appendix D).

Dissolved oxygen ranged between 7.03 and 7.95 mgL<sup>-1</sup>, levels conducive to an environment supporting a high fish biomass (Romaine 1985), and tended to be higher in the wet season.

The pH had a narrow range between 9.45 and 9.66, which, although above the Ugandan standards, is considered normal for Lake Albert. Interestingly, this pH is higher than the values for the rivers draining off the escarpment, which ranged between 8.8 and 9.2. These findings are similar to the findings from 50 years ago, when the pH ranged from 8.9 to 9.1 (Talling 1963). Typically, a pH above 9.5 can lead to ammonia toxicity to fisheries (Beveridge, 1996); however, this high level apparently has little effect on the biodiversity of the lake.

Conductivity had a narrow range of 632 to 634 µS.cm<sup>-1</sup>; these values were less than those recorded by Talling (1963), which ranged from 720 to 780 µS.cm<sup>-1</sup>.

Interestingly, this pH is higher than the values for the rivers draining off the escarpment, which ranged between 8.8 and 9.2.

#### General Parameters – Rivers and Wetlands

Generally, the water quality parameters assessed for the rivers and wetlands fell within the accepted limits of Uganda's national drinking water standards (the Ugandan standards), as published by the National Water and Sewerage Cooperation (NEMA 1996). Dissolved oxygen varied at all sites sampled, ranging from 1.3 mg.L<sup>-1</sup>, near the mouth of the Masika River, to 9.2 mg.L<sup>-1</sup> in the Well Pad 2 River mid-stream. The pH also varied, ranging from 7.1 at the mouth of the Masika River, to 9.6 at the mid-stream of Well Pad 2 stream. Similarly,





the conductivity ranged  $278 \mu\text{S}\cdot\text{cm}^{-1}$  at the mouth of the Masika River, to  $966 \mu\text{S}\cdot\text{cm}^{-1}$  in the middle of the Bugoma Lagoon.

### Phosphorous Compound – Lake Albert

The total phosphorus concentration ranged from  $21.1$  to  $43.9 \mu\text{g}\cdot\text{L}^{-1}$  between the dry and wet season; these concentrations supply soluble reactive phosphorus (SRP) in the range of  $0.0$  to  $3.4 \mu\text{g}\cdot\text{L}^{-1}$ , a range which would typically lead to the lake waters off the Buhuka Flats being classed as nutrient poor and not very productive (that is, oligotrophic) (OECD 1982). However, other nutrient sources within the water column, such as soluble reactive silica (SRSi) (see below), which drives diatom biomass, counter the oligotrophic conditions. Furthermore, green and blue-green algae readily absorb the SRP, thereby increasing the biomass in the otherwise nutrient poor (that is, oligotrophic) environment.

Within such a system, as represented by Lake Albert, aquatic biodiversity can be unique, with endemic and range-restricted species encountered that are adapted to the nutrient poor conditions. Such a system is sensitive to phosphorous loading, such as that occurring from agricultural run-off and pollution, which, in turn, can adversely affect fish production.

### Phosphorous Compound – Rivers and Wetlands

The overall total phosphorus concentration ranged from  $79.4$  to  $350.3 \mu\text{g}\cdot\text{L}^{-1}$ , supplying SRP in the range of  $0.2$  in the middle of the lagoon, to  $34.5 \mu\text{g}\cdot\text{L}^{-1}$  at the mouth of the Masika River. The influence of soap products, from local people doing their washing in the rivers, may have resulted in the high concentrations observed.

### Silica Compound – Lake Albert

The SRSi levels were relatively high in all samples from Lake Albert, and ranged from  $410$  to  $1096 \mu\text{g}\cdot\text{L}^{-1}$  in the wet and dry season. As mentioned above, SRSi is readily absorbed by diatoms; in particular, the diatom genera *Cyclotella* and *Nitzschia* are abundant in the lake, and form an important food source for fish.

### Silica Compound – Rivers and Wetlands

At the time of the first wet season sampling, sufficient rainfall had not yet fallen; hence, there was no water connectivity between the lake and the rivers and wetlands draining off the escarpment and the Buhuka Flats. As such, the concentration of SRSi was similar in these systems compared to the lake. The SRSi ranged from  $456.7$  to  $929.7 \mu\text{g}\cdot\text{L}^{-1}$  in all the nine sites sampled. It is expected that, following sufficient rainfall, these systems will be flushed and the concentrations of SRSi may drop.

### Nitrogen Compound – Lake Albert

Total Nitrogen concentration decreased in the wet season, ranging from  $149$  to  $872 \mu\text{g}\cdot\text{L}^{-1}$ , compared to the dry season, which ranged from  $407.8$  to  $729.2 \mu\text{g}\cdot\text{L}^{-1}$ . These concentrations support high levels of phytoplankton, which is the major carbon producer, and food source for fish in Lake Albert. As expected, the total nitrogen concentrations in the vicinity of the confluence of the Masika River and Lake Albert were the highest, at  $872 \mu\text{g}\cdot\text{L}^{-1}$ .

Ammonia concentrations within the water column were similar in the wet and dry seasons, ranging from  $3.1$  to  $28.7 \mu\text{g}\cdot\text{L}^{-1}$ . The higher concentrations of  $22.3$  and  $28.7 \mu\text{g}\cdot\text{L}^{-1}$  were recorded from the sampling locations associated with Pad 2 and the confluence of the Masika River and Lake Albert. The reason as to why the concentration at the Pad 2 sampling locations is unclear; however, the high concentrations recorded at the Masika River's confluence were not unexpected. Although ammonia becomes detrimental to fish health at concentrations greater than  $20 \mu\text{g}\cdot\text{L}^{-1}$ , typically, within a healthy aquatic ecosystem, it is converted immediately into nitrate in the presence of oxygen (via the intermediate product nitrite) (Delince 1992). Within the samples from Lake Albert, nitrate concentrations ranged from  $15.3$  to  $95.3 \mu\text{g}\cdot\text{L}^{-1}$  indicating efficient conversion of toxic ammonia, driven by good dissolved oxygen concentrations, which were greater than  $7.0 \text{mg}\cdot\text{L}^{-1}$  at all sites (see above). Consequently, ammonia concentrations were observed to drop below  $5 \mu\text{g}\cdot\text{L}^{-1}$  further into the lake. As such, the aquatic ecosystem offshore from the Buhuka Flats is healthy, as indicated by the efficient conversion of ammonia and good concentrations of dissolved oxygen.





### Nitrogen Compound – Rivers and Wetlands

The total nitrogen concentration ranged from 104.2 to 863.  $\mu\text{g.L}^{-1}$  at all sites sampled, which presented an ammonia concentration ranging from 4.4 to 33.9  $\mu\text{g.L}^{-1}$ . Nitrate concentrations ranged from 18.6 to 179.7  $\mu\text{g.L}^{-1}$ . The higher concentrations were observed at the mouth of the Masika River and could be the result of an oxygen deprived environment.

### Chlorophyll a – Lake Albert

Chlorophyll a, an indirect determinant of algal biomass (Heckey 1993), ranged from 1.0 to 3.1  $\mu\text{g.L}^{-1}$  in the wet season, and 1.7 to 8.7  $\mu\text{g.L}^{-1}$  in the dry season. Similar to the observed concentrations of phosphorous, the concentrations of chlorophyll a qualifies the lake waters offshore from the Buhuka Flats as nutrient poor (that is, oligotrophic – being in the range of 0 to 8.5  $\mu\text{g.L}^{-1}$  according to OECD 1982). Within Lake Albert, this range indicates a stable food supply for the fish without visible algal blooms.

### Chlorophyll a – Rivers and Wetlands

Chlorophyll a, ranged from 0.0 to 18.3  $\mu\text{g.L}^{-1}$ , which could be expected to decrease during the wet season as the algal growth is suppressed by the growth of emergent wetland plants and the shade they produce.

### Faecal Coliform – Lake Albert

The National Water and Sewerage Cooperation (NEMA 1996) identifies that zero colony forming units (CFU) per 100 mL is the recommended concentration of faecal coliform for drinking water, it is also the recommended level for the maintenance of healthy aquatic biodiversity.

Faecal coliform concentrations within Lake Albert, at the locations sampled, ranged from 2 to 8 CFU per 100 ml in the dry season, jumping to 2 to 100 CFU per 100 ml in the wet season. As expected, concentrations were higher at the inshore sampling sites, and decreased further out into the lake.

The marked difference between the wet and dry season concentrations can be attributed to the increased run-off experienced during the wet season into the lake.

### Faecal Coliform – Rivers and Wetlands

All sites within the rivers and wetlands assessed on the Buhuka Flats had faecal coliform ranging from four to 504 CFU per 100 ml. These values are well above the National Water and Sewerage Cooperation standard.

## 6.2.2 Phytoplankton

The phytoplankton communities of the near-shore environment of Lake Albert, the Masika River, the Kamansinig River and the Well Pad 2 stream are summarised below.

### Species Richness, Diversity and Abundance – Lake Albert

Thirty-five genera were identified in the phytoplankton communities of Lake Albert, offshore from the Buhuka Flats, over the wet season compared to the 26 genera identified during the dry season. These comprised: blue-green algae – 20 species comprised of eight genera; simple algae (that is, Cryptomonads, Dinoflagellates, Euglenoids) – two species each (interestingly, the Euglenoids were not recorded in the dry season); diatoms – eight species comprised of six genera; filamentous green algae – 18 species comprised of 12 genera. Generally, the species diversity increased across all groups from the dry season to the wet season.

Among the blue-green algae, six species (*Anabaenopsis tanganikae*, *Aphanocapsa incerta*, *Ap. nubium*, *Chroococcus limneticus*, *Planktolyngbya circumcreta* and *P. limnetica*) were present at all sites sampled. Other taxa, such as *Cylindrospermopsis*, *Merismopedia* and *Coelosphaerium*, were rare. One species, *Microcystis aeruginosa*, recently identified as the only toxin-producing blue-green algae in Ugandan freshwater habitats (Okello et al. 2010), was conspicuously absent at all sites sampled. This is a notable observation given that this species prefers polluted environments, and its absence suggests that the waters of Lake Albert are a relatively pristine environment (Okello et al. 2010).

Of the five genera of Diatoms identified, three taxa: *Cyclotella* species; *Nitzschia acicularis* and *N. fonticola*, were represented at all sites sampled, while *Navicula* species were found at fewer sites. The genus



*Aulacoseira* was conspicuously absent in the dry season, yet was encountered during the wet season. This difference between the seasons could be attributed to this taxon being selectively fed upon by zooplankton and fish. Interestingly, this taxon has almost disappeared from Lake Victoria, which has apparently led to the virtual total decline of some native fish species which feed almost exclusively upon it (Ogutu-Ohwayo et al, 2002).

The majority of the 18 filamentous green algae species identified were widespread across all the sampling sites. Some taxa increased during the wet season compared to the dry season, although seven genera (*viz.*, *Closterium*, *Dictyosphaerium*, *Didymocystis*, *Gonatozygon*, *Kirchneriella*, *Oocystis* and *Pediastrum* were rare during the dry and wet seasons.

In general, there was an increase in the species composition per site from the dry season to the wet season; ranging from 15 to 23 species in the dry, to 18 to 26 species in the wet.

The composition of the phytoplankton community within Lake Albert differed between the dry and wet seasons. During the dry season, the phytoplankton communities were dominated (>50%) by diatoms (ranging from 15.46 to 85.35 mm<sup>3</sup>.L<sup>-1</sup>), with blue-green algae comprising between 13.92 and 70.34 mm<sup>3</sup>.L<sup>-1</sup> of the biomass volume. In the wet season, blue-green algae dominated (ranging from 0.84 to 3.53 mm<sup>3</sup>.L<sup>-1</sup>), while the diatom communities decreased markedly to between 0.91 and 3.09 mm<sup>3</sup>.L<sup>-1</sup>. Both diatom and blue-green algae biomass tended to be higher in the offshore areas (~1 km) compared to the inshore areas.

It is interesting to note that the diatom biomass recorded in Lake Albert during the dry season was lower than that recorded in Lake Edward and Lake George, which ranged from 45 to 230 mm<sup>3</sup>.L<sup>-1</sup> and 90 to 420 mm<sup>3</sup>.L<sup>-1</sup>, respectively (Okello and Kurmayer, 2011).

Single-celled algae (that is, Cryptomonads) populations increased from the dry season (~0.94 mm<sup>3</sup>.L<sup>-1</sup>) to the wet season (from 0.09 to 0.14 mm<sup>3</sup>.L<sup>-1</sup>), although they only constituted a minor proportion of the phytoplankton communities. Multi-celled algae (in particular, Dinoflagellates) were not recorded during the dry season, yet, during the wet season, their populations increased to around 0.12 mm<sup>3</sup>.L<sup>-1</sup>. Green algae biomass ranged between 1.38 and 37.19 mm<sup>3</sup>.L<sup>-1</sup> in the dry season, and dropped to between 0.05 and 8.46 mm<sup>3</sup>.L<sup>-1</sup> in the wet season.

The observed difference between the seasons could be attributed to the increase in grazing pressure from the larger zooplankton population, which increased during the wet season (see below).

### Species Richness, Diversity and Abundance – Rivers and Wetlands

Thirty-one genera, belonging to five families were identified in the phytoplankton communities of the rivers and wetlands of the LSA. These included: blue-green algae – ten species in nine genera; simple algae – one species in a single genus (which was only recorded in the lagoon); diatoms – eight species in eight genera; filamentous green algae – 13 species in 11 genera.

Among the blue-green algae, five species (*Anabaena circinalis*, *Aphanocapsa nubilium*, *Chroococcus limneticus*, *Merismopedia tenuissima* and *Planktolyngbya limnetica*) were present at all sites sampled. Like the blue-green algal community of Lake Albert, *Microcystis aeruginosa*, was conspicuously absent at all sites sampled. This is a notable observation given that this species prefers polluted environments, and its absence suggests that the waters of the rivers and wetlands of the LSA are a relatively pristine environment (Okello et al. 2010).

Ten of the identified species of diatom were recorded in the Masika River and the lagoon. Only four species were recorded in the Kamansinig River, these were: *Navicula gastrum*; *Nitzschia acicularis*; *Ni. onticola* and *Surirella* sp.). The rare *Aulacoseira* was present in the lagoon.

The 13 species of filamentous green algae were mainly present in the Bugoma Lagoon sites, being absent from the Kamansinig River, and only represented by one genus, *Actinastrum*, in the Masika River.

The Bugoma Lagoon wetland had the highest total phytoplankton biomass of ~190 mm<sup>3</sup>. L<sup>-1</sup>. Of this, more than 50% was contributed by the blue-green algae, while diatoms and filamentous green algae shared similar proportions. Simple algae appeared in minor quantities.



### Species of Concern

Of all the taxa of phytoplankton identified during the dry and wet season, only the genus *Aulacoseira* is a group of special interest. As mentioned above, this has almost disappeared from Lake Victoria, which has apparently led to the virtual total decline of some native fish species which feed almost exclusively upon it (Ogutu-Ohwayo et al. 2002). Its existence in Lake Albert suggests that this lake is still relatively undisturbed and unpolluted.

For further discussion and assessment of these and other species that may potentially occur in the CHAA, see Section 6.3.2.

### Habitats

The near-shore environment of Lake Albert, the watercourses and wetlands of the Buhuka Flats support a diverse assemblage of phytoplankton. These areas are naturally higher in nutrients that support higher biomasses of phytoplankton, which, in turn, drive the food web of the south-eastern shores of Lake Albert, and, potentially, contribute to the food web of the wider lake.

For further discussion and assessment of habitats, please refer to Section 6.3.1.

### 6.2.3 Zooplankton

The zooplankton communities of the near-shore environment of Lake Albert, the Masika River, the Kamansinig River and the Well Pad 2 stream are summarised below.

#### Species Richness, Diversity and Abundance – Lake Albert

The zooplankton communities within Lake Albert and the Bugoma Lagoon wetland of the Buhuka Flats are dominated by three taxa: Copepod crustaceans (Order: Copepoda) (comprising three species); water fleas (Order: Cladocera) (comprising seven species), and rotifers, or wheel animals (Phylum: Rotifera) (comprising nine species). Depending on the location: Copepods comprised, on average, 56% (range: 0% to 100%) of the zooplankton biomass in the dry season samples, and 69% (range 8% to 100%) in the wet season samples; water fleas comprised, on average, 49% (range: 10% to 90%) of the biomass in the dry season samples, and 38% (range 0% to 92%) in the wet season samples; rotifers comprised, on average, 26% (range: 0% to 80%) of the biomass in the dry season samples, and 33% (range 0% to 100%) in the wet season samples.

In Lake Albert, zooplankton abundance was dominated by the Copepods *Thermocyclops neglectus* and *Mesocyclops* sp., the water fleas *Diaphanosoma excisum* and *Moina micrura*, and the rotifer *Keratella tropica*. The relative abundance was higher in the offshore environments compared to the inshore environments. In contrast, the zooplankton abundance of the Bugoma Lagoon was dominated by the rotifers *Brachionus angularis*, *B. calyciflorus*, and *B. falcatus*, with the Copepod *T. neglectus* and water fleas *M. micrura* and *D. excisum* also strongly represented.

The Bugoma Lagoon supported the highest number of rotifer species, yet also exhibited the highest decline in species richness from the dry season to the wet season. There was a marked difference in species diversity, compositions and abundance between the dry season and the wet season at all sites sampled. In the lagoon, total abundance of zooplankton decreased from an average of ~1,000,000 individuals per square metre in the dry season, to an average of ~180,000 individuals per square metre in the wet season. The dominant taxa comprising these abundances also shifted between the dry and wet seasons; as mentioned above, Copepods were more abundant in the wet season than the dry, while rotifers dominated in the dry season. The opposite trend was observed in Lake Albert. Here the zooplankton abundance increased from an average of ~350,000 individuals per square metre in the dry season, to an average of ~1,400,000 individuals per square metre in the wet season. Certainly, in Lake Albert, there was a tremendous increase in abundance of all taxa in the wet season compared to dry season.

In comparison with other lakes in the Victoria basin (for example, see Vincent et al. 2012, Mwebaza-Ndawula et al. 2003), the diversity of zooplankton in this particular part of Lake Albert is generally low. Nonetheless, key species reported to characterize zooplankton assemblages of Ugandan lakes, and which are important food species for fish (Mwebaza-Ndawula et al. 2004), occurred in relatively high abundance.





### Species of Concern

No species of concern were identified in the zooplankton communities in the LSA.

For further discussion and assessment of other species that may potentially occur in the CHAA, see Section 6.3.2.

### Habitats

The zooplankton communities of near-shore and off-shore environments of Lake Albert reflect a healthy water habitat dominated by Cyclopoid Copepods. A similar healthy environment was reported in Lake Albert over a decade ago by Lehman et al. (1998). The exception is the Bugoma Lagoon, where high numbers of rotifers were observed. Typically, a high abundance of rotifers, notably species of *Brachionus*, as observed in the lagoon, reflect elevated levels of pollution, or eutrophication, of waterbodies (Radwan and Popiolek 1989, Tasevska et al. 2010). This is possibly the current situation in the Bugoma Lagoon where *B. angularis* and *B. calyciflorus* were abundant. Another notable observation was the absence of the predatory Cyclopoid copepod, *Mesocyclops* sp., in the lagoon, yet it was recorded in >75% of the other sites sampled. The changes in the zooplankton assemblages in the Bugoma Lagoon, and the dominance of rotifers, could also be due to the lack of connectivity with the lake and the consequent stagnation of the water coupled with the increase in nutrient loads in-flowing from the Kamansinig River and the hinterland, as well as the presence of Hippopotamus adding to the nutrient loading.

Nevertheless, the dominance of Copepods in the aquatic habitats of Lake Albert is important. These taxa are keystone species in sustaining fish communities in most water bodies (Mwebaza-Ndawula et al. 2001, Mwebaza-Ndawula et al. 2003, Mwebaza-Ndawula et al. 2004). As such, the near-shore habitats, wetlands of the lower Masika River and Kamansinig River, and the large bays of the Buhuka Flats are important habitats for zooplankton.

For further discussion and assessment of habitats, please refer to Section 6.3.1.

### 6.2.4 Macro-invertebrates

The macro-invertebrate communities of the near-shore environment of Lake Albert, the Masika River, the Kamansinig River and the Well Pad 2 stream are summarised below.

#### Species Richness, Diversity and Abundance – Lake Albert

Species richness of mayflies (Order: Ephemeroptera), stoneflies (Order: Plecoptera) and caddis flies (Order: Trichoptera) (EPT) in Lake Albert was low, yet similar in the dry season and both wet season samples, ranging from zero to 15.

The species diversity was equally low during the dry and both wet seasons, being dominated by a few taxa; the mayfly *Povilla adusta*, the molluscs *Melanoides tuberculata*, *Gabbia humerosa*, and *Bellamya unicolor*, and the freshwater clam *Corbicula africana*. In terms of abundance of individuals, mayflies comprised the largest number (0 to 1707 larvae per square metre), yet only included three species (dominated by *Povilla adusta*, with *Caenis* sp. and *Baetis* sp. also occurring); while no stoneflies were recorded during any of the seasonal surveys. Among the caddis flies, only members of the family Psychomidae were recorded in the dry season, only *Dipseudopsis* spp. (Family: Dipseudopsidae) was recorded in the first wet season, and both were recorded during the second wet season survey.

The larvae of phantom midges (Family: Chaoboridae) were common and abundant (up to 294 larvae per square metre) in the inshore and offshore sediments of Lake Albert across all sampling periods. Larvae of non-biting midges (Family: Chironomidae) and biting midges (Family: Ceratopogonidae) were also relatively abundant.

Snails (Phylum: Gastropoda), freshwater mussels (bivalves in the Order: Unionoidea), freshwater clams (*Corbicula africana*) and aquatic worms (Phylum: Annelida, Class: Oligochaeta) were the most common and abundant macro-invertebrates recovered from bottom sediments in the dry and wet seasons.

During the dry season, five snail taxa (*Melanoides tuberculata*, *Cleopatra* sp., *Gabbia* (*Gabiella*) *humerosa*, *G. walleri* and *Bellamya unicolor*) were the most abundant and widely distributed. In contrast, during the first



wet season two species (*viz.*, *M. tuberculata* and *B. unicolor*) were the most abundant and widely distributed taxa. Their densities ranged from zero to 1064 individuals per square metre in the dry season, and zero to 896 in the wet season. In the second wet season, *Gabbia humerosa* was most abundant with densities of 1401 - 1078 individuals per square metre recorded at two locations (Pad 2 and Pad 3 inshore); this species was not recorded at any other location on that occasion.

Two species of freshwater mussel (*viz.*, *Byssanodonta parasitica* and *Corbicula africana*) were common and abundant. Their densities in the dry season were markedly less than the first wet season survey (that is, zero to 294 individuals per square metre in the dry, compared to zero to 672 individuals per square metre in the first wet). In the second wet season survey, significantly higher number of individuals were recorded at the Pad 1 offshore site only (zero to 518 individuals of *B. parasitica* and zero to 140 individuals of *C. africana*) with the other sites supporting relatively low numbers.

The densities of aquatic worms (Oligochaetes) ranged from zero to 98 individuals per square metre in the dry season, up to 168 per square metre in the first wet season, and up to 42 per square metre in the second wet season; the highest densities being recorded in inshore habitats adjacent to the lagoon.

Total taxa richness scores in Lake Albert were similar in both of the wet and dry season samples, with an average of seven species (range: three to 15 species) recorded in the inshore habitats, and eight species (range: two to 15 species) recorded in the offshore habitats. Importantly, the densities of the various groups varied, sometimes significantly, between the dry and wet seasons, with higher densities observed for all groups during the first wet season.

Based on EPT and total taxa richness scores for the dry and wet season surveys, the sampling sites in Lake Albert and the wetlands of the Buhuka Flats showed varying environmental conditions. Only the inshore sampling sites in the vicinity of Pad 1 and Pad 4-2 reflected a moderate to good environmental condition. Those sampling locations in the vicinity of Pad 2 (offshore), Pad 3, Pad 4-2 (offshore), Pad 5 (offshore), and in the Bugoma Lagoon reflected fair environmental conditions. Interestingly, the remainder of the sites, namely Pad 1 (offshore), Pad 2 (inshore) and Pad 5 (inshore) reflected poor environmental conditions.

Typically, densities of benthic macro-invertebrates, especially the insect group (that is, mayflies, stoneflies, caddis flies and midges), tend to increase from the dry to wet season (Wetzel 2001). The numbers of the mayfly *Povilla adusta* and phantom midges (*Chaoboridae*) recorded across the three sampling periods reflected this trend, however, the same trend was not observed in any of the other macro-invertebrate samples from the LSA. The reasons for this are not clear based on the current data and sampling regime.

It should be noted that a dedicated search conducted in November 2014 as part of the second wet season survey did not find *Bellamya rubicunda* or *Gabiella candida* at either the Lake Albert inshore stations or the wetland streams of the Buhuka Flats (see Appendix D).

### Species Richness, Diversity and Abundance – Rivers and Wetlands

The most common macro-invertebrates in the rivers and wetlands were: mayflies (*Baetis* spp., Order: Ephemeroptera) and caddis flies (*Cheumatopsyche* spp., Order: Trichoptera), with densities ranging from 0 to 6,723 individuals/m<sup>2</sup> and 2801 individuals/m<sup>2</sup> respectively; dragonflies (*Brachymesia* spp., Order: Libellulidae; and) with mean density of 0 to 126 individuals/m<sup>2</sup> recorded in the River Masika at the base of the escarpment; water bugs (*Macrocoris* spp., Order: Hemiptera), with mean density of 0 to 98 individuals/m<sup>2</sup>; non-biting midges (Order: Diptera, Chironomidae), with 0 to 98 individuals/m<sup>2</sup>; and aquatic worms, with 0 to 385 individuals/m<sup>2</sup>. Other notable groups included black flies (Order: Diptera, Simuliidae), with densities ranging from 0 to 210 individuals/m<sup>2</sup>, recorded from the mouth of Well Pad 2 stream. Importantly, the densities and species composition of the various groups varied, sometimes significantly, between the dry and wet seasons, with higher densities observed for all groups during the wet seasons; particularly the dragonflies which were present in much greater densities during the second wet season survey than the previous sampling events.

Freshwater snails (*Biomphalaria* spp.) were recorded in the Kamansinig River, Masika River and Well Pad 2 stream.

Based on the EPT scores, the Kamansinig River at the foot of the escarpment reflected a moderate to good environmental condition, the remainder of the river and wetland habitats were fair to poor. Total taxa richness



scores for the Kamansinig River and Well Pad 2 stream reflected a moderate to good environmental condition compared to the other sites sampled.

It is important to note that the watercourses draining off the escarpment are seasonal. Therefore, low counts of macro-invertebrates would be expected in some of these sites during the dry season. This was reflected in the observed EPT diversity recorded during the second wet season; *Baetis* spp. (Ephemeroptera) and *Cheumatopsyche* spp. (Trichoptera) were abundant during this time, yet absent from the dry season surveys. Similarly, the Bugoma Lagoon appears to be isolated from the lake during the dry season. This could account for the low dissolved oxygen in the bottom sediments resulting from inadequate mixing and decomposition of plant material. These conditions in the Bugoma Lagoon support the fact that intolerant species of mayflies, stoneflies and caddis flies were absent, yet more tolerant species like non-biting midges and aquatic worms were common.

### Species of Concern

All the species recorded in the LSA were Least Concern, or not yet evaluated by the IUCN. The Critically Endangered freshwater Mud Snail (*Gabiella candida*) has, to date, only been recorded from Butiaba, about 90 km north of the LSA (Kyambadde 2010a, GBIF 2017).

It should be noted that a dedicated search conducted in November 2014 as part of the second wet season survey did not find *Bellamyia rubicunda* or *Gabiella candida* at either the Lake Albert inshore stations or the wetland streams of the Buhuka Flats (see Appendix D). Nevertheless, taking a precautionary approach, there remains a potential that these species could be found within suitable habitat in the LSA (see Section 6.3.2).

No other known macro-invertebrate species of concern were identified as occurring, or have a potential to occur, in the LSA.

### Habitats

The near-shore and inshore habitats of Lake Albert in the vicinity of the Buhuka Flats offer a diverse array of substrates (that is, clay/snail shells, sand/plant materials, soft mud, and rock/shells) that support a rich diversity of the benthic macro-invertebrates. Similarly, the wetlands associated with the Masika River, Kamansinig River, Well Pad 2 stream, and the watercourses along the pipeline route, all provide important habitat for macro-invertebrates. The Bugoma Lagoon offers a unique habitat for fish and other organisms including macro-invertebrates.

For further discussion and assessment of habitats, please refer to Section 6.3.1.

## 6.2.5 Fish

The freshwater fish communities of Lake Albert, the Masika River, the Kamansinig River and the Well Pad 2 stream are summarised below.

### Species Richness, Diversity and Abundance

Twenty-four fish species comprising 19 genera and nine families were recorded from Lake Albert in the LSA. This represents ~45% of the 53 fish species reported to occur in Lake Albert (Greenwood 1966).

Four species contributed the most biomass (% weight of catch as a surrogate for abundance – although the limitations of this are recognised) in the first wet season survey from all sampling locations in the LSA; these included: the Nile Perch (*Lates niloticus*) (42%), *Ragoge* (*Brycinus nurse*) (17%), *Ngassa* (*Hydrocynus forskahlii*) (12%) and *Angara* (*A. baremoze*) (3%). These are also some of the most commercially important species in Lake Albert (Taabu-Manyahu et al. 2012). In comparison, five species contributed the most biomass by catch in the dry season samples: Nile Perch (40%), *Ngassa* (18%), *Angara* (9%), Shield-head Catfish (*Synodontis schall*) (9%) and Black Nile Catfish (*Bagrus bajad*) (8%).

During the second wet season, *Ragoge* (*B. nurse*) dominated the catch (73%) with similar numbers being caught at Pad 1, Pad2, Pad 3 and Pad 5 sampling sites. The remainder of the catch was largely made up of Nile Perch (*L. niloticus*) (8%), *Ngassa* (*H. forskahlii*) (4%), Haplochromines (3%), *Angara* (*A. baremoze*) (1%) and *Bagrus bayad* (1%).



Other important species, in terms of abundance, included: Silver Butter Catfish (*Schilbe intermedius*) (6%), which was only recorded at two sampling locations (that is, Pad 2 and Pad 4-2) during the first wet season; a Cichlid (*Thoracochromis (Haplochromis) wingati*) (~5%) at Pad 5 during the first wet season; and another Cichlid (*T. loati*) (3%), which was recorded in sizable numbers in the vicinity of Pad 1.

Five species (*Imberi (Alestes macrolepidotus)*, *Muziri (Neobola bredoi)*, *Mpoi (Barilius (Distichodus) niloticus, Citharinus citharus, C. latus)*, *Kisinja (Barbus (Labeobarbus) bynni)*, Lake Albert Cichlid (*Thoracochromis (Haplochromis) avium*)), were not recorded during the dry season, yet were recorded in the first wet season. Conversely, the Shield-head Catfish, a Barb (*Barbus (Labeobarbus) perience*) and Mango Tilapia (*Sarotherodon galilaeus*) were recorded in the dry season, yet not in the wet season. Similarly, Senegal Bichir (*Polypterus senegalis*), *Imberi*, Barbel (*Clarius gariepinus*), *Mpoi*, *Muziri*, and Lake Albert Cichlid that were recorded in the first wet season, were not recorded in the during the second wet season. One catfish species, Sudan Squeaker (*S. frontosus*), and the African Carp (*Labeo horrie*), were recorded for the first time during the second wet season surveys. These seasonal differences seen in the community composition may be due to migratory responses to inflow of run-off into the lake.

Species composition within the Bugoma Lagoon was noticeably different compared to Lake Albert. Five species dominated the biomass of each catch; these included: Nile Tilapia (*Oreochromis niloticus*) (36%); Singidia Tilapia (*O. leucostictus*) (18%); *Imberi* (18%); Senegal Bichir (*Polypterus senegalis*) (16%); and *Angara* (8%).

In general, the fish community in the near-shore zone of the LSA is composed of a fairly uniform, multi-species mix of various ages in good condition. The diversity, age classes and condition of the species assessed is a reflection of adequate food and a healthy environment. Species distribution within the lake environment reflects that habitat requirements of each species.

The near-shore artisanal fishery is dominated by gillnets, and is mostly focused on Nile Perch, *Ragoge*, *Ngassa*, and *Angara*. *Muziri* features strongly in the seine net fishery. For more information and discussion on the fisheries, see Chapter 18 Ecosystem Services Review.

### Species of Concern

One locally threatened species, the African Electric Catfish (*Malapterurus electricus*) may occur in the LSA. Although this species has not been recorded from LSA, it is known from the wider Lake Albert (Azeroual et al., 2010).

Eight commercially important species are known to occur within the LSA. These include the *Imberi*, *Angara*, Catfish (*Clarias lazera*), *Mpoi*, *Ngassa (Hydrocynus vittatus)*, Nile Tilapia, Mango Tilapia, and Zill's Tilapia (*T. zillii*). The three species of *Mpoi*, the *Angara*, *Ngassa*, Butter Catfish, and Shield-head Catfish are of particular importance because these species have become very rare in Lake Albert (Wandera and Balirwa 2010). Lake Albert supports at least ten endemic, range-restricted fish species, notably the commercially important *Angara*, *Ngassa* and *Imberi* (Wandera 2000, Campbell et al. 2005, Wandera and Balirwa 2010), all of which have been recorded in the LSA.

Lake Albert has previously experienced increases in populations of *Ragoge* and *Muziri* (Wandera and Balirwa 2010). In 2012, these two species constituted 51% and 34% of commercial fish catches, and indications then were that their populations were increasing in the lake (Taabu-Manyahu et al. 2012). An increase in the populations of these species, which feed exclusively on zooplankton, could have implications for the zooplankton community structure due to increased pressure on these micro-organisms. However, recent baseline work conducted in November 2017 as part of the social baseline update indicated that fisheries are now under significant pressure in Lake Albert since the construction of the escarpment road. This new easy access to the Lake Albert shore has facilitated a large increase in fishing demand, with trucks arriving from Kampala on a near-daily basis to collect the catch. The size of fish being caught by local communities on the Flats appears to have decreased significantly, with only white bait being observed drying during 2017, compared to much larger Nile Perch observed during fieldwork conducted prior to the escarpment road being in place (see Vol 4c Social Assessment, Chapter 18 Ecosystem Services Review).



For further discussion and assessment of other species of concern that may potentially occur in the CHAA, see Section 6.3.2.

### Habitats

The near-shore habitats of Lake Albert and the wetlands of the lower Masika River and Kamansing River, and the large bays of the Buhuka Flats are important habitats for zooplankton. In particular, the dominance of Copepod zooplankton is important because these taxa are keystone species in sustaining fish communities in the lake (Mwebaza-Ndawula et al. 2001, Mwebaza-Ndawula et al. 2003, Mwebaza-Ndawula et al. 2004). This is supported by the high biomass, high catch rates, and strong multispecies fishery in the near-shore waters of Lake Albert in the LSA.

For further discussion and assessment of habitats of concern, please refer to Section 6.3.1.

## 7.0 OVERALL BIODIVERSITY VALUE

Presented below are the findings of the baseline biodiversity value of the CHAA, as described based on a description of the two identified components of overall biodiversity defined in Section 3.3.3, that is, ecosystems and habitats, and species of concern. Within each of these, valued components are described.

### 7.1 Ecosystems and Habitats of Concern

Uganda falls at the confluence of a number of regional centres of endemism (White 1983): the Guinea-Congo Forest; Lake Victoria Basin; Afro-Tropical Highlands; Somali-Masai; and Sudan and Guinea Savannah. This has resulted in the region having a unique suite of biodiversity. Indeed, the Albertine Graben is an area of high endemism and threatened species (Critical, Endangered and Vulnerable); with over 50% of birds, 39% of mammals, 19% of amphibians and 14% of reptiles and plants of mainland Africa occurring in this region (Plumptre et al. 2003). Furthermore, the Albertine Graben and Lake Albert, within which the CHAA is located, is recognised as: part of the Eastern Afrotropical Biodiversity Hotspot (CI 2014); an Endemic Bird Area (Stattersfield et al. 1998); a Key Biodiversity Area (IUCN 2010); and within three globally important ecoregions, notably, the Albertine Rift Montane Forests, the East Sudanian Savanna, and the Rift Valley Lakes Freshwater Ecoregion (Olson and Dinerstein 1998). As such, the Graben is recognised as an area of global importance for conservation. For further discussion and analysis of the biodiversity importance of the wider Albertine Graben and Lake Albert, see, for example, Emerton and Muramira (1999), Wandera (2000), Plumptre et al. (2003, 2005, 2007, 2010, 2011), NEMA (2010), Taabu-Munyaho et al. (2012), AECOM (2012, 2013), CI (2014). The near-shore environment of Lake Albert, and the wetlands associated with the rivers draining off the escarpment, support important populations of commercially important fish species (including important breeding areas), freshwater turtles, crocodiles, invertebrates, algae and aquatic plants, staging grounds for migratory birds, as well as biogeochemical processes that drive the ecosystem.

The Project is located within the Albertine Graben, and adjacent to Lake Albert. As such, it can be expected that the CHAA includes ecosystems and habitats that are important components of the biodiversity and biogeographical significance of the Albertine Graben and Lake Albert.

Nevertheless, the CHAA is located in an area that is heavily influenced by human activities, and currently does not support a high species richness and diversity. Historically, the Buhuka Flats was part of the Buhuka Community Wildlife Area (CWA) (UWA, pers. comm.); however, it was not maintained as a CWA, and, subsequently, the large animal populations declined (UWA, pers. comm.), and, apart from a few Hippopotamus, all the large mammals have disappeared from the area. As such, it is no longer recognised as a CWA and was degazetted in 2002 (RPS 2006). Nevertheless, as identified in Section 6.1.1.1, significant natural vegetation still on the Buhuka Flats and along the escarpment (Figure 4a), 84% of which is considered natural habitat under the IFC's definitions (Figure 15). These vegetation communities and habitats support populations of smaller mammals, mammals, reptiles, amphibians, and invertebrates, including species of concern (see Section 6.3.2).

The Buhuka Flats and the adjacent escarpment area are currently primarily used for livestock grazing, subsistence agricultural purposes, firewood collection, charcoal manufacture, and harvesting of non-timber forest products, while the near-shore environment of Lake Albert supports a strong artisanal fishing industry. These activities exert pressures on the ecosystems, and, undoubtedly have affected the ecological integrity of





the ecosystems of the LSA (for example, see NEMA 2010, Wandera and Balirwa 2010). In addition, the ease of access to the Buhuka Flats has been altered with the construction of the escarpment road; this increased ease of access may facilitate increased anthropogenic demand for services such as livestock grazing and firewood harvest, exacerbating pressure on nearby ecosystems.

Similarly, the ecosystems along the pipeline route have experienced severe pressure from subsistence agricultural practices that have resulted in the transformation of a large proportion of the natural habitats in recent history (see Figure 15) (for a review, see AECOM 2012).

Regardless of the intense agricultural, fishing and subsistence pressures within the CHAA, the entire area supports ecosystems and habitats of greater or lesser value. For the intents of this impact assessment, these have been grouped into the following broad ecosystems: the near-shore environment of Lake Albert; the vegetation communities and corridors along the rift valley escarpment; wetlands of the Buhuka Flats and pipeline route; and the ecosystem services they offer (see Chapter 18 Ecosystem Services) for a more detailed discussion of these ecosystems as they relate to the supply of ecosystem services); and the Bugoma Central Forest Reserve.

Information on the ecology, ecosystem functioning and processes of some of these systems are available, either in published literature (for example: Lake Albert (see, Green 1971, Wandera 2000, Campbell et al. 2005, Wandera and Balirwa 2010, Taabu-Munyaho et al. 2012); the importance of the vegetation along the escarpment as a corridor (see Plumptre et al. 2003, 2007, Ayebare et al. 2013); the Bugoma Central Forest Reserve (see Plumptre et al. 2010, 2011)), or in previous studies undertaken in the area (for example, AWE 2008a, b, 2013a, 2014a, EAC 2013, 2014). As mentioned in Section 3.3.3.1.1, these available studies were supplemented by seasonal surveys of the LSA to gauge an understanding of the current integrity of the ecosystems in the CHAA (see Section 6.1, 6.2, and Appendix C and D).

As relevant to this impact assessment, a summary of those studies, and other relevant scientific literature, is provided below to provide a description of the ecological integrity of each of these ecosystems (based upon: composition and structure; linkages and corridors; key processes, including drivers of change; representativeness; resilience and stability). Therefore, as relevant to the Project, and the potential effects the Project may have on those biodiversity values, presented below is a description of the baseline of ecosystems and habitats of concern that comprise the receptors/valued components for impact assessment.

### 7.1.1 Near-Shore Habitats of Lake Albert

For the intents of this impact assessment, the near-shore environment of Lake Albert is defined as the area within 500 m of the shore of the lake, with water depths typically between 0 m to 12 m; beyond this distance the depth of the lake increases to 18 m and more (NEMA 2010). This area supports a diversity of habitats, water depths, temperature regimes and concomitant flora and fauna communities. These near-shore habitats are also recognised as sensitive ecological areas by NEMA (2010).

Lake Albert supports the most diverse commercial fisheries in Uganda, with at least 55 species (Wandera and Balirwa 2010). This diverse fish fauna supports an array of multi-species fisheries along the lake's length, and, in terms of fisheries production, Lake Albert is second only to Lake Victoria (Taabu-Munyaho et al. 2012). Although this is currently the case, Lake Albert is facing multiple environmental changes, including declining fish species diversity, over fishing and reduced catches, hyper-eutrophication in places and hypoxia (Campbell et al. 2005). Furthermore, the ecology of the lake, and the processes driving the ecosystem, remains poorly understood (for example, Lehman et al. 1998, Wandera 2000, Wandera and Balirwa 2010).

#### Composition and Structure

The south-central part of the lake, in the vicinity of the LSA, is dominated by the steep sided escarpment, and predominantly deep waters close to the shore (Taabu-Munyaho et al. 2012). In the immediate vicinity of the LSA, that is, the Buhuka Flats, the ecosystem of the lake is composed of a diversity of habitats. Indeed, all the six main habitat types in Lake Albert identified for fishes by Wandera and Balirwa (2010) (*viz.*, shallow river-associated waters, open sandy shores, lagoons, large bays, rocky escarpments, and, open-water habitats), occur within the near-shore areas of the LSA (Figure 5). Of these habitats, those workers identified river mouths, lagoons, near-shore waters of large bays, and rocky areas as priority habitats for fish breeding.



All these habitats are well represented in the CHAA (Figure 5), and constitute approximately 5.7 km (river mouths 0.3 km; lagoons 1.4 km; large bays 2 km; rocky shores 1.9 km).

Wandera and Balirwa (2010) reported that, together, throughout the lake, these habitats support 40 species of fish. River mouths contributed the highest number of species, while lagoons supported the highest number of individuals and biomass (Wandera and Balirwa 2010). Indeed, these workers specifically recognise the Bugoma Lagoon of the Buhuka Flats as one of these important habitats in the lake. The importance of the near-shore habitats within the CHAA is reflected in the diversity and abundance of fish species recorded during the aquatic ecology surveys (see Section 6.2.5, and Appendix C). Approximately 36% of the species known to inhabit Lake Albert were recorded from the LSA, and these were represented by a range of age and size classes, all in good condition. This suggests that the diversity in composition and structure of the habitats in the LSA is important for the life cycle of a range of species, including many commercial species.

Not only are the shoreline waters, and their concomitant structural diversity, important habitats for juvenile fishes (Wandera and Balirwa 2010), they are also important areas for invertebrate and algae communities that form an important food source for juvenile and adult fish. The submerged and emergent aquatic plants that are common in the shallow waters off the Buhuka Flats and the wetlands associated with the Masika River and Kamansinig River provide a diversity of structural refuge and breeding sites not only for fish, but also for many invertebrate prey species (such as, Copepod crustaceans, water fleas, rotifers, midges, mayflies, stoneflies, caddis flies). Importantly, the dominance of Copepod crustaceans in the aquatic habitats of Lake Albert is important. These taxa are keystone species in sustaining fish communities in most water bodies (Mwebaza-Ndawula et al. 2001, Mwebaza-Ndawula et al. 2003, Mwebaza-Ndawula et al. 2004). As such, the near-shore habitats, wetlands of the lower Masika River and Kamansinig River, and the large bays of the Buhuka Flats are important habitats for these components of the zooplankton community.

### Linkages and Corridors

The near-shore aquatic habitats within Lake Albert are all part of the wider lake ecosystem. In terms of size, the lake, as a contained ecosystem, is not large, being approximately 5300 km<sup>2</sup> (~150 km long by ~35 km wide) (Wandera and Balirwa 2010). And, although it is primarily fed by the Victoria Nile (which drains Lake Kyoga and Lake Victoria) and the Semliki River system, which account for 83% of inflow, the biodiversity of the lake is unique. Lake Albert supports at least ten endemic fish species, notably the commercially important *Angara*, *Ngassa* and *Imberi* (Wandera 2000, Campbell et al. 2005, Wandera and Balirwa 2010), as well as numerous mollusks, crabs and seed shrimp (Crustacea: Ostracoda) (Plumptre et al. 2003).

The movement of fish and other animals within Lake Albert is currently not understood, nor is the movement of fish and other animals into and out of the lake (for example, see Campbell et al. 2005, Wandera and Balirwa 2010). Nevertheless, certain species, notably *Angara*, *Ngassa*, African Butter Catfish (*Schilbe niloticus*), and African Catfish (*Clarias lazera*) do make lateral movements within the lake to shallow waters in bays and up river systems to breed (Akinyi et al. 2010a, Azeroual et al. 2010c, e, Lalèyè et al. 2010), and these breeding movements tend to be seasonally based (Kusnierz et al. 2014). How far these species travel within the lake to reach their preferred breeding areas is not known. However, given that there are only a few major river systems entering the lake that would be suitable for breeding, and the afore-mentioned species are caught throughout the lake (see Taabu-Munyaho et al. 2012), it is conceivable that individuals would travel large distances within the lake to breed. Likewise, juveniles would disperse widely within the lake.

Based on the above, and the fact the CHAA supports important breeding habitat for fish (that is, river mouths, lagoons, near-shore waters of large bays, and rocky areas (after Wandera and Balirwa 2010)), it is possible that the near-shore habitats within the vicinity of the Buhuka Flats are an important end point, or starting point, for fish.

### Key Processes and Drivers of Change

As mentioned, Lake Albert is primarily fed by the Victoria Nile and the Semliki Rivers. Certainly, in the case of the Victoria Nile, and many of the other rivers and streams feeding the lake, these flow through areas of high agricultural activity. Consequently, it can be expected that these rivers carry with them high levels of nutrients and other contaminants. Certainly, Wandera (2000) and Wandera and Balirwa (2010) identified that



agricultural run-off was having real effects on increasing the nutrient levels of the lake, a process known as eutrophication.

Intense fishing pressure from commercial and artisanal fisheries has led to the decrease in populations of many of the commercially important species of the lake. Available data suggest that the overall shape of Lake Albert's food web is triangular, dominated by predatory species like the Nile Perch and *Ngassa* at the top, and the various tilapia species, *Ragoge* and *Muziri* forming the broader base (Campbell et al. 2005). As noted by Campbell et al. (2005), the Nile Perch has undergone dietary shifts suggesting that recent overfishing of it and its prey species may be changing the nature of the entire food web within the lake, and ultimately the ecosystem integrity. Another observation by Wandera and Balirwa (2010) was the shift towards increased catches of *Ragoge* and *Muziri* in recent times. These workers suspected that this shift was due to the decreased populations of larger species, and the consequent reduced catches of those species. With the increased pressure on *Ragoge* and *Muziri* populations (which are food species for the larger species, and the main predators of zooplankton, especially Cyclopoid copepods (Green 1971)), it is not unreasonable to assume that the fish community within the wider lake may change in the long-term.

The fishing activities based out of the 11 fishing villages in the CHAA and vicinity (that is, five on the Buhuka Flats, with six located to the north and south of the Buhuka Flats), undoubtedly put pressure on the local fish populations. Indeed, apart from the published accounts of the decrease in commercial fish stocks (Wandera 2000, Wandera and Balirwa 2010, Taabu-Munyahu et al. 2012), anecdotal accounts from the local fisher-folk also identify noticeable decreases in catches of fish per unit effort.

Likewise, the presence of approximately 22,000 people on the Buhuka Flats and other nearby villages who do not have access to running water and sanitation, will put large pressures on the nutrient loading of the inflowing lake waters of the LSA.

Despite the current human population of the Buhuka Flats and LSA, the near-shore aquatic environment was generally healthy (as reflected by the water quality, and the phytoplankton and zooplankton communities, in particular the Copepod crustacean communities – see Section 6.2). The notable exception was the Bugoma Lagoon. Here the water quality parameters and zooplankton communities reflected an environment with a high nutrient load and stagnant water. However, this was expected given that this wetland was isolated from the lake, possibly due to the lack of significant surface water inflow (at the time of sampling) and the resident Hippopotamus population. It is expected that the water quality, and the associated zooplankton diversity, in the Bugoma Lagoon will improve as inflow increase during the rainy season. As such, it is plausible that the health status of this system will fluctuate between the seasons, driven by increased water flows in the rainy season, and the subsequent drying in the dry season. These seasonal effects will be observed for all the wetlands in the CHAA (see Section 6.3.1.2.3).

Current climate change models predict that Uganda is likely to experience more extreme periods of intense rainfall, an erratic onset and cessation of the rainy seasons and more frequent episodes of drought (GCCA 2012). An overall increase of approximately 180 mm per annum is predicted, which will result in a mean annual rainfall for the CHAA of 880 mm to 1580 mm for the period 2020 to 2039. Current records of the lake's water level indicate that it varies by approximately 4 m every year, as influenced by rainfall. How this increased rainfall could influence that water level of the lake is unknown.

Within the CHAA, the main drivers of change influencing the near-shore aquatic habitats derive from the human population exerting pressure on fish stocks, including catching breeding individuals, and polluted run-off from the 11 villages in the area. There has been a substantial increase in the local population over the past 10 years, driven by a multitude of factors such as regional instability, attractive livelihood opportunities to engage in fishing on Lake Albert, and more recently, interest in capitalising from opportunities related to oil and gas developments.

### Representativeness

The majority of the aquatic habitats within the CHAA are represented widely around the lake. Within the southern section of the Ugandan side of the lake, from the northern end of Kaiso-Tonya Community Wildlife Area in the north, to the Uganda-Democratic Republic of Congo (DRC) border in the south (approximately 210 km of shoreline), shallow river-associated waters comprise ~14 km (7%) of shoreline, and include the mouths





of, amongst others, the Wasi River, the Muzizi River, the Nkussi River, the Masika River, and a number of others. Open sandy shores account for ~68 km (32%), lagoons (including the wetland systems of the Wasi and Muzizi Rivers) account for ~54 km (26%), large bays account for ~13 km, and rocky escarpments account for ~25 km (12%).

Based on these figures, the CHAA supports 3% (0.4 km) of the shallow river-associated waters, 13% (8.5 km) of the open sandy shore habitat, 2% (1.2 km) of the lagoons, 14% (1.8 km) of large bays, and 7% (1.6 km) of rocky escarpment (Figure 14). Of particular significance is the lagoon, which, according to Wandera and Balirwa (2010), is only one of six such lagoons in the lake.

### Resilience and Stability

Lake Albert is approximately ~5500 km<sup>2</sup>. Given its size, the buffering effect of the large water body, it is expected to be reasonably resilient and stable.

### Overall Condition

Based on the findings of the baseline studies, the condition of the near-shore habitats of Lake Albert in the CHAA are near pristine. Despite the pressures of local the local population and their associated fishing activities, these habitats are still in a natural state and support health populations of phytoplankton, zooplankton, macro-invertebrates and fish. Some localised areas, particularly around the fishing villages are slightly degraded due to polluted run-off and gross pollution.

#### 7.1.2 Escarpment Vegetation Corridors

The escarpment supports natural vegetation bounded on the east by highly modified subsistence agricultural landscapes, and the Buhuka Flats and Lake Albert on the west. The vegetation communities form part of a continuous strip of vegetation to the south and to the north.

### Composition

Within the CHAA, the vegetation communities on the escarpment are composed of a mix of the described vegetation communities of the area. Four communities dominate; comprising 66%, that is, open wooded grassland (28%), followed by dense wooded grassland (23%), dense bushland (14%), and riverine bushland (10%). The remaining communities make up the rest. These form a continuous corridor of vegetation of approximately 2443 ha within the CHAA, which is bounded on the east by agriculturally modified landscapes, and the Buhuka Flats on the west (Figure 11).




Figure 11: Escarpment vegetation communities

**Structure**

The vegetation communities form a diverse mosaic of habitats across the escarpment, with 15 different communities represented (Table 2).

**Table 2: Extent of escarpment corridor vegetation communities within the CHAA**

Escarpment corridor vegetation communities	Area (Ha)
Dense Bushland	337.7
Dense Riverine Woodland	28.3
Dense Wooded Grassland	589.0
Grassland with Thicket	5.2





<b>Escarpment corridor vegetation communities</b>	<b>Area (Ha)</b>
Open Bushland and Shrubland	71.5
Open Grassland	31.5
Open Wooded Bushland	214.0
Open Wooded Grassland	552.2
Open Woodland	85.2
Open Woodland/Dense Bushland	353.2
Riverine Bushland	89.7
Riverine Woodland	69.8
Riverine Woodland and Bushland	2.9
Seasonal Wetland	5.1

The vegetation communities appear to be driven by landscape and geology. For example, open wooded grassland and dense wooded grassland tend to be restricted to the crest of the escarpment; dense bushland tends to be restricted to the steep slopes; with the watercourses support riverine communities.

### Linkages and Corridors

In the regional context, these vegetation communities contribute to maintaining the continuity of the many high-priority conservation sites within the wider Albertine Graben. These corridors are also recognised as important for the maintenance of the evolutionary processes unique to the Albertine Graben (Ayebare et al. 2013).

On the local scale, within the CHAA, the escarpment is an important habitat for many species recorded in the LSA. It not only forms a continuous corridor along the length of Lake Albert, but it also offers important refugial sites for many of the small mammal species; notably the cavity and cave roosting bat species.

### Key Processes and Drivers of Change

Within the CHAA, the main drivers of change influencing the vegetation communities along the escarpment are from livestock grazing, fuel wood harvesting, charcoal manufacture and the conversion of natural vegetation for subsistence agriculture. As mentioned, this is particularly noticeable between the current vegetation on the escarpment and the Bugoma Central Forest Reserve (see Figure 4).

The keeping of livestock forms a substantial component of the local socio-economic structures in the CHAA, particularly on the Buhuka Flats. Livestock numbers are large and there is strong evidence for overgrazing on the Buhuka Flats extending up onto accessible regions of the escarpment corridor.

Large trees on the escarpment are becoming rarer as these individuals are selectively harvested for the manufacture of charcoal, which is typically on-sold. Smaller woody species are regularly harvested for fuel wood used directly in the fishing villages.

The harvest of fibre and other house construction materials is common on the escarpment. For example, thatching grass is regularly harvested on the escarpment and transported to the local fishing villages (Figure 12).



*Figure 12: 'Chutes' used for transport of thatching grass harvested from the escarpment*

The occurrence of frequent fires was also evident on the escarpment. Too-frequent fire is known to detrimentally affect the functioning and processes of savanna ecosystems (Smith et al. 2013).

The condition of the vegetation communities on the escarpment suggests that all these processes are contributing to changes in their composition and structure. In particular, encroachment of bushland into the grassland and woodland communities appears to be quite frequent. Bush encroachment is a typical consequence of changed land use practices like intense livestock grazing and removal of large, ecosystem engineer species like African Elephant (Wigley et al. 2009). Nevertheless, local communities often prefer these changes because they result in increased woody resources for building and firewood and increased browse availability for goats (Wigley et al. 2009). Consequently, in the long-term, a positive feedback loop could result in these communities becoming completely transformed.

### **Representativeness**

The vegetation communities of the escarpment form part of a contiguous vegetation corridor, extending for approximately 70 km, from the Toro-Semliki Wildlife Reserve in the south, to the Kabwoya Wildlife Reserve in the north. This corridor is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2007). This corridor is recognised as being of regional importance for savanna species, and potential climate refugia in the future (Ayebare et al. 2013).

### **Resilience and Stability**

The natural vegetation along the escarpment is under increasing pressure from surrounding land uses (that is, subsistence agriculture) and increased population pressure. The condition of the vegetation communities on the escarpment suggests that all the current drivers of change are contributing to changes in their composition and structure. In particular, encroachment of bushland into the grassland and woodland



communities appears to be quite frequent. Bush encroachment is a typical consequence of changed land use practices like intense livestock grazing and removal of large, ecosystem engineer and keystone species like African Elephant (Wigley et al. 2009). Nevertheless, as previously mentioned, local communities often prefer these changes because they result in increased woody resources for building and firewood and increased browse availability for goats. Consequently, in the long-term, a positive feedback loop could result in the frequency of open grassland and wooded communities becoming transformed to dense bushland and shrubland.

### Overall Condition

Based on the findings of the baseline studies, and more up-to-date knowledge of the current pressures on the escarpment vegetation communities since the opening of the escarpment road, the overall condition of vegetation communities on the escarpment in the CHAA are slightly to moderately degraded. The combined pressures of livestock grazing, natural resource harvesting and frequent fire indicate that the composition of these communities are changing.

### 7.1.3 Wetlands and Riparian Habitats

The CHAA supports permanent wetlands and seasonally flooded grasslands, and riverine forest associated with riparian watercourses. These habitats form potentially important habitat for species of concern; in particular, Grey Crowned Crane, Madagascar Pond Heron and migratory birds.

#### Composition

The seasonal or permanent nature of the wetland communities of CHAA tended to determine the species composition. Seasonal wetlands were comprised largely of *Cyperus articulatus* and sparse *Typha* sp., while the permanent wetlands were largely composed of *Cyperus papyrus*, *Phragmites mauritianum* and *Typha capensis* (see Section 6.1.1.1).

In the pipeline region of the CHAA, the mainly seasonal watercourses support a riverine woodland vegetation community that is dominated by *Acacia* species, with *Crateva* sp. *Acalypha* sp. *Tamarindus indica*, and *Rhus natalensis* also occurring in the canopy layer, and an understorey dominated by *Grewia trichocarpa* and *Azima tetraacantha*.

#### Structure

The seasonally flooded grassland communities had a very simple structure that, as mentioned, is largely determined by the water inundation frequency. The vegetation typically dies back during the dry season, and undergoes vigorous growth during the wet season. Consequently, the vegetation is dominated by short species that are quick growing. The underlying soils also form a determinant of these wetlands, with all the seasonally flooded areas being underlain by black cracking clays (Vertisols).

The permanent wetlands are dominated by tall growing species, such as *Phragmites* sp. and *Typha* sp. These species tend to be perennial and do not die back during the dry season. As such, the permanent wetlands maintain a dense, tall and emergent vegetation cover throughout the year.

The riverine forest associated with the riparian watercourses are characterised by an upper storey canopy layer that does not interlock and remains open, with a relatively structurally-homogenous herbaceous layer beneath.

#### Linkages and Corridors

The wetlands of the CHAA are typically part of larger watercourses that drain the area. For example, the Masika River forms permanent wetlands along its lower reaches on the Buhuka Flats, while the Kamansing River forms seasonally flooded grasslands on some of its lower reaches.

Although these wetlands are restricted to the particular watercourses of the CHAA, they do form stepping-stone habitats for wetland species moving up and down the Albertine Graben. In particular, migratory bird species. The Albertine Rift is an important stopping point for migratory birds during their annual migration, between October and March, from Europe and Asia (Byaruhanga et al. 2001).





The larger permanent wetlands along the pipeline route form part of a more extensive network of wetlands on the plateau, many of which connect to those in the Bugoma Central Forest Reserve.

### Key Processes and Drivers of Change

Currently, the wetlands of the CHAA appear to be functioning and stable. Water flows into, and through, the permanent wetlands of the CHAA appear to follow the natural wet-dry season cycles. This is reflected in the seasonal differences in the water quality data, and seasonality of particularly the insects and other animals of the wetlands (see Section 6.2.1 and 6.2.4, respectively). Seasonal water flows through these wetlands are important for flushing nutrients through the system and adding nutrients to the system. This is especially evident in the Bugoma Lagoon. The changes in the zooplankton in the lagoon, and the dominance of rotifers, could be due to the lack of flow and connectivity with the lake during the dry season.

An exception is the seasonally flooded grasslands associated with the Kamansing River. The road leading from the foot of the escarpment and the borrow pit has noticeably influenced the flow regimes and drainage patterns of this wetland. The wetland on the western side of the road appears to have been altered and no longer fully functional (see Figure 19). This is a good example of how surface and sub-surface flows are important to maintaining a wetland's functionality and processes.

The permanent wetlands of the CHAA are important sources of fibre for house construction and container manufacture. This is especially evident in those wetlands close to human settlements, such as those along the pipeline route and the lower Masika River. Given their use as fibre sources, the frequency of human-induced fire in these wetland communities appears to be reduced.

Harvesting of large trees for the manufacture of charcoal is a noticeable driver in the riverine forest communities associated with the riparian watercourses within the CHAA, as well as trampling and grazing by cattle.

### Representativeness

The CHAA supports approximately 1157.9 ha of wetlands, of which 85.3 ha are classified as seasonal. In addition, 840 ha of riparian vegetation communities are associated with drainage lines and riparian areas. 151 ha or 12% of all wetlands within the CHAA occur on the Buhuka Flats. The representativeness of these wetlands within the wider area is unknown; however, within the CHAA, wetlands constitute approximately 1.5% of habitats. This figure may under-represent the true extent of wetlands in the CHAA, particularly in the Bugoma Central Forest Reserve. Wetlands within that forest area were not confirmed through field investigations.

### Resilience and Stability

Many factors contribute to the resilience and stability of wetlands, and those factors are dependent on the location and type of wetland (Carvalho et al. 2013). Typically, the overriding factor determining a wetland's resilience is the maintenance of the hydrological regime, and the amount of water entering and leaving the wetland. Other factors include nutrient loading, species diversity, trampling and grazing by livestock, and fire frequency (Carvalho et al. 2013).

The seasonal or permanent nature of the wetland communities of the CHAA tended to determine the species composition. Seasonal wetlands were comprised largely of *Cyperus articulatus* and sparse *Typha* sp., while the permanent wetlands were largely composed of *Cyperus papyrus*, *Phragmites mauritianum* and *Typha capensis* (see Section 6.1.1.1).

As described, the altered flow of water to the seasonally flooded grasslands on the western side of the road leading from the escarpment has noticeably affected the functioning of the portion of the wetland downstream of the road crossing (Figure 19). It is unknown whether reinstatement of those flows would resurrect those wetlands. Consequently, this scenario suggests that these seasonally flooded grasslands are not very resilient, and highly susceptible to changes in water flow patterns. Additionally, these wetlands appear to be favoured by livestock for grazing. This is possibly because the conditions within the wetlands support lush vegetation growth during the wet season, which persists well into the dry season. Consequently, this grazing pressure and trampling may be adversely influencing the species composition of these habitats.



The permanent wetlands are associated with the larger watercourses in the CHAA. Therefore, these wetlands could be reasonably resilient provided the flow volumes of those watercourses are maintained. Livestock grazing in these wetlands has intensified since the opening of the escarpment road (Figure 13); hence, trampling and overgrazing are expected to affect the resilience and stability of these systems. These permanent wetlands are dominated by *Phragmites* sp. and *Typha* sp.; these plants are known to be very resilient to pollution and increased nutrient levels. Indeed, members of these taxa are very efficient at removing pollutants from water and are used to treat polluted water in constructed wetlands (Vymazal 2011).



Figure 13: Cattle watering at Kamansinig crossing, Buhuka Flats (Nov 2017)

The macro-invertebrate communities within the permanent wetlands are susceptible to changes in water quality, and the composition of these communities change with the seasons (see Section 6.2.3 and 6.2.4). However, these seasonal changes in these communities appear to be part of the natural cycle of these wetlands.

### Overall Condition

Based on the findings of the baseline studies, and more up-to-date knowledge of the current pressures on wetlands in the Buhuka Flats, the overall condition of the wetlands in the CHAA are slightly to moderately degraded due to pressures of livestock grazing and natural resource harvesting.

#### 7.1.4 Bugoma Central Forest Reserve

The Bugoma Central Forest Reserve is widely recognised for its biodiversity importance. For example, it is one of a handful of forests that constitute a network and corridor of critical biodiversity sites in Uganda, and supports populations of, amongst other species, Eastern Chimpanzee (*Pan troglodytes schweinfurthii*), African Elephant (*Loxodonta africana*), Nahan's Francolin (*Ptilopachus nahani*), as well as a variety of endemic birds and butterflies (NEMA 2010). It is also: the source of numerous rivers in the region, including the Nguse and Rutowa Rivers (NEMA 2010); and an Important Bird Area (BirdLife International 2014a). Of the 65 forested protected areas surveyed for biodiversity in Uganda, Bugoma Central Forest Reserve ranked 11 in overall biodiversity value, and 15 in terms of rarity value (BirdLife International 2014a). For a more detailed assessment and other studies on this forest's biodiversity and biogeographical importance, see, for example:



Plumptre et al. (2010, 2011), and Ayebare et al. (2013). Hence, a very limited description is provided here, with only important features, as they relate to the potential effects of the Project, discussed.

### Composition and Structure

This forest is a medium altitude, moist, semi-deciduous forest with a high biodiversity. About half of the forested portion of the Bugoma Central Forest is dominated by Iron Wood (*Crynometra alexandri*); a further 38% is mixed forest (BirdLife International 2014a). Two-hundred-and-fifty-seven species of trees and shrubs have been recorded in the forest, seven of which are Albertine Rift endemics, 12 are globally threatened and 14 are listed in IUCN red list (Plumptre et al. 2003, 2010, 2011).

### Linkages and Corridors

The Bugoma Central Forest Reserve is recognised as being an important part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwaru Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2007). However, it is disconnected from other forest reserves, such as Budongo Central Forest Reserve, and is bordered by subsistence agricultural communities and settlements that are placing increased pressure on it (NEMA 2010, Plumptre et al. 2010, 2011).

### Key Processes and Drivers of Change

While the Bugoma Central Forest Reserve is a reserve for timber resources, and harvesting of timber is reportedly undertaken on a sustainable basis (NEMA 2010); it is not a designated *wildlife* protected area. Given the increased human population around the forest, there have been increased incidents of illegal logging, fire wood collection and charcoal manufacture, and bush meat harvesting (NEMA 2010, Plumptre et al. 2010). Between 2011 and 2012, increased immigration into the areas surrounding the forest resulted in increased deforestation, with an estimated 5,000 ha of the forest subject to encroachment by about 1,000 families and pit-sawyers (AECOM 2013). Recent reports of illegal logging of hardwood timber, and illegal land clearance in Bugoma CFR with the intention of transformation for sugarcane plantations (ACBF, 2016) suggest that pressure on the reserve has increased substantially.

Many of the tree species within the forest require elephants or large primates to disperse their seeds. With the declining populations of these species, particularly elephants, many tree species are declining (Plumptre et al. 2010).

### Representativeness

This forest is one the last large tracts of remaining medium altitude, moist, semi-deciduous forest in western Uganda (Plumptre et al. 2010, 2011). The CHAA encompasses the entire Central Forest Reserve of approximately 401 km<sup>2</sup>.

### Resilience and Stability

Given the Bugoma Central Forest Reserve's size (approximately 401 km<sup>2</sup>), it should be relatively resilient and stable from disturbance. However, the forest, and the populations of species it supports, is under intense pressure from the surrounding human population. What the long-term effects of the increase in the human population around the forest, including the illegal settlements within the forest (see Mugerwa 2013), could have on the forest are unknown, but are expected to be damaging to the resilience and stability of the forest. This forest is isolated from other forests in the region, and is being eroded on the outer edges (see Figure 4).

### Overall Condition

Based on the findings of the studies and research conducted by others, the overall condition of the Bugoma Central Forest Reserve is slightly degraded to moderately degraded due to pressures of illegal logging, natural resource harvesting, and the current human pressures surrounding the forest.



## 7.2 Species of Concern

Appendix G provides the details of the critical habitat screening and appraisal of species of concern that could occur in the CHAA.

The information and data reviewed, together with the baseline field surveys, identified that the CHAA (Figure 2) supports a diverse and rich species assemblage, with numerous species of concern potentially occurring in the area. Based on those reports and surveys, the CHAA has the potential to support a possible 96 species of concern (excluding invasive species, which are discussed in Section 7.2.5), as per the definitions in Section 3.2.3. These include: six plant species; two macro-invertebrate species; 14 fish species; two butterfly species; one dragonfly and damselfly species; three amphibian species; nine reptile species; 44 bird species; and 14 mammal species (Appendix G). Forty-six of those species were recorded during the field surveys (Appendix C and D).

### 7.2.1 Threatened, Range-Restricted/Endemic and Statutory Species

Overall, the CHAA has a potential to support populations of seven globally recognised Critically Endangered and Endangered species (one Critically Endangered macro-invertebrate (the Mud Snail (*Gabbiella candida*), four Endangered birds (Madagascar Pond-Heron, Grey Crowned-Crane (*Balearica regulorum*), White-backed Vulture (*Gyps africanus*), Hooded Vulture (*Necrosyrtes monachus*)), one Endangered mammal (Eastern Chimpanzee (*Pan troglodytes schweinfurthii*)). The CHAA supports the range-restricted Nahan's Francolin (*Ptilopachus nahani*) which was down-listed from Endangered to Vulnerable in 2017 (IUCN 2017); and potentially supports five populations of other globally recognised Vulnerable birds and four populations of mammals; as well as one Near Threatened tree, one macro-invertebrate, 12 birds and four mammals (Appendix G).

Of these, 23 species (one Critically Endangered, three Endangered, five Vulnerable, 14 Near Threatened), could occur in the LSA, and hence potentially be affected by direct Project impacts. A further nine species (one Endangered, six Vulnerable, two Near Threatened) could occur in the CHAA, and hence potentially be affected by indirect, induced and cumulative impacts from the Project.

The CHAA also supports approximately 19 Palearctic migratory bird species that are listed under Appendix II of the Convention on Migratory Species (CMS 2014). For more details on those, and other locally important species, see Appendix G.

Of the above-mentioned species, those that have the potential to trigger critical habitat were carried forward for a more detailed and formal appraisal to determine if they could trigger critical habitat, as per the methods presented in Appendix B. That appraisal is presented in Appendix G, and the results are summarised in Section 6.3.2.2.

### 7.2.2 Species of Economic and/or Cultural Importance

Lake Albert supports a strong commercial fishing industry. As discussed in Section 6.3.1.2.1, this fishery is second in importance only to that of Lake Victoria. Within the CHAA, many of the species that form the mainstay of this fishery were recorded, or are probably present (as summarised in Appendix G, Table G1). These include: *Imberi*, *Angara*, *Mpoi*, African Catfish, *Ngassa*, Nile Tilapia, African Butter Catfish, Mango Tilapia, and Zill's Tilapia.

The off-shore habitats within Lake Albert form an important fishing ground for the 11 fishing villages in the CHAA and vicinity (that is, five on the Buhuka Flats, with six located to the north and south of the Buhuka Flats). As discussed in Section 6.3.1.1.1, the near-shore habitats supported in the CHAA are most likely important breeding habitats for many of the commercially important species supporting those 11 fishing villages.

Consequently, the near-shore environments of the CHAA also trigger critical habitat for Criterion 13 because these are seen to be important breeding areas for commercially important fish species, as well as important fishing grounds for those species. This near-shore habitat accounts for about 794 ha of the CHAA (Figure 5).



### 7.2.3 Species listed under CITES

CITES (the Convention on International Trade in Endangered Species of Wild Fauna and Flora) is an international agreement between governments (Uganda has been a signatory since 1991 – see Section 4.0), which has the aim to ensure that international trade in species does not threaten their survival (CITES Secretariat 2017).

The CHAA supports known populations of CITES-listed Appendix II plant and animal species. These include Aloes (*Aloe* spp.), Candelabra Tree (*Euphorbia candelabrum*), Graceful Chamaeleon, Smooth Chamaeleon, Serrated Hinge-back Tortoise, Leopard Tortoise, Nile Monitor, Nile Crocodile, Grey Parrot, and Hippopotamus (see Appendix G, Table G1). Other CITES-listed Appendix II species that may occur in the CHAA, yet have not been recorded, include the Secretarybird, White-bellied Pangolin, African Golden Cat, and Giant Ground Pangolin (Appendix G, Table G1; UNEP-WCMC 2018).

Importantly, within the CHAA, and Bugoma Central Forest Reserve in particular, CITES-listed Appendix I species have a high likelihood of occurring, or have been recorded (see Plumptre et al. 2010, 2011). These species include African Elephant, Eastern Chimpanzee and Leopard (Appendix G, Table G1).

CITES-listed species are grouped in the Appendices according to how threatened they are by international trade. Appendix I lists species that are the most endangered and are threatened with extinction; CITES prohibits international trade in specimens of these species, except when the purpose of the import is not commercial (CITES Secretariat 2017). Species listed in Appendix II are not necessarily threatened with extinction now, however, unless trade in these species is controlled, they could be seriously threatened (CITES Secretariat 2014). International trade in individuals of Appendix II species may be authorized by appropriate authorities when specific conditions are met, above all, that trade will not be detrimental to the survival of the species in the wild (CITES Secretariat 2017).

The populations of the CITES-listed species potentially and actually occurring in the CHAA are not precisely known. Indications are that the certain species are relatively common, for example Aloe and Candelabra Tree (Figure 4b), while others, such as African Elephant may be very uncommon (see Plumptre et al. 2010).

### 7.2.4 Evolutionarily Distinct Species

This section identifies those species of concern that could trigger critical habitat for Criteria 5 under the IFC's definitions (IFC 2012a). As discussed in Section 3.3.3.1.2, critical habitat for key evolutionary processes does not have quantitative thresholds (for example, see IFC, 2012a). Therefore, for the purposes of this impact assessment, expert opinion was used to identify critical habitat with respect to evolutionarily distinct species as indicators of the landscape-level features that can influence key evolutionary processes.

The Albertine Rift is known as a centre of endemism driven by unique evolutionary processes; indeed it is the most species rich region in Africa for vertebrates (Plumptre et al. 2003, 2007). It is also a recognised Endemic Bird Area (Stattersfield et al. 1998), testament to a collection of unique species derived from a set of unique evolutionary processes. Lake Albert too is a recognised centre of endemism within the Albertine Rift, and is of biogeographical significance for a number of taxa and species, in particular snails (Plam et al. 2008) and fish (Wandera and Balirwa 2010).

At the species level, Criterion 5 applies for “distinct species”, which include those termed Evolutionarily Distinct and Globally Endangered (EDGE) species (GN 95 IFC 2012b, Jetz et al. 2014, ZSL 2014). Jetz et al. (2014) define evolutionary distinctness as a measure of “a species's contribution to the total evolutionary history of its clade and is expected to capture uniquely divergent genomes and functions”. Based on this definition, these workers identified bird species and particular regions where birds occur, that are of enormous value for protecting evolutionary diversity. Bird species with the greatest evolutionary distinctness are often located outside of areas traditionally identified as conservation priorities (Jetz et al. 2014). Species representing the most evolutionary

Based on the works of Jetz et al. (2014), and a geographically-based search of the EDGE species database that covered the spatial extent of the CHAA (see ZSL 2017), two such EDGE species were identified as potentially occurring in the CHAA. These were the Shoebill and Secretarybird, which are discussed and assessed in Appendix G.





### 7.2.5 Invasive and Potentially Invasive Species

The CHAA has been affected to a greater or lesser extent by human activities for a very long time. These activities have altered the landscape to a greater or lesser extent, the most noticeable being the conversion of the natural vegetation on the plateau above the escarpment to agricultural crop land (AECOM 2013). The influence of people on the Buhuka Flats is also very noticeable, the majority of the flats affected by livestock grazing, small-scale agriculture, fuel wood harvest, and building material collection.

As identified in Section 6.1.1, five invasive species were recorded in the LSA, they tended to be localised, and uncommon, although within certain areas, local populations were high (Figure 14). Three of these, the Giant Sensitive Tree (*Mimosa pigra*), Lantana (*Lantana camara*) and Water Hyacinth (*Eichhornia crassipes*) were the commonest species recorded, predominantly on the Buhuka Flats and the shore of Lake Albert (Appendix D). These species are recognised as some of most noxious weeds in the world (Lowe et al. 2000).

Other species recorded include: Water Lettuce (*Pistia stratiotes*) (in a wetland community on the Buhuka Flats, where it was locally abundant); and Castor Oil Plant (*Ricinus communis*) (recorded from wetland and woodland communities).

Other potentially invasive species to note were Neem (*Azadirachta indica*), Jatropha (*Jatropha curcas*) and Parkinsonia (*Parkinsonia* sp.). It is noted that many of these species have been planted by the local communities and offer important cultural and other ecosystem services.



# BIODIVERSITY IMPACT ASSESSMENT




Figure 14: Invasive species recorded on the Buhuka Flats





## 8.0 NATURAL AND MODIFIED HABITAT WITHIN THE CHAA

An assessment and identification of the natural and modified habitats was restricted to the CHAA (Figure 2). The results of the assessment are summarised in Table 2.

**Table 3: Natural and modified habitat in the CHAA**

Habitat	Area (ha)	Area (km <sup>2</sup> )	Proportion of CHAA
Modified	12,944	129.4	16%
Natural	68,303	683.0	84%

The majority of the Buhuka Flats and the escarpment is natural habitat, while the pipeline route is dominated by modified habitat (Figure 15). In the case of the Buhuka Flats, although this area is inhabited (including the villages of Kiina, Nsunzu, Nsonga Kyabasambu, Kyakapere) and under pressure from subsistence agriculture and livestock grazing, a large proportion of the flats is dominated by natural vegetation communities. These include, amongst others: thicket-grassland mosaic, open grassland, bushed grassland, and Wetlands (see Section 6.1.1.1, and Figure 4a and b). Of interest is the connectivity of the natural habitat that occurs on the Buhuka Flats and the escarpment. These habitats form part of the wider Murchison Falls National park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2003) (see Section 6.3.1.2.1).

The plateau above the escarpment is markedly different. Here the area has been largely transformed into cropland and plantations, interspersed with settlements and patches of natural habitat, comprised largely of wetlands (see Figure 6).




Figure 15: Natural and Modified habitat in the CHAA





## 9.0 CRITICAL HABITAT WITHIN THE CHAA

Appendix G presents the screening and assessment of all species of concern that could trigger Criterion 1, 2, 3 and 5 critical habitat within the CHAA, as per the methods presented in Appendix B. Ecosystems that could trigger Criterion 4 critical habitat are discussed in Section 9.1.2.

As discussed, quantitative thresholds for critical habitat are available for Criteria 1 to 3. No such thresholds exist for the other criteria. Therefore, a qualitative assessment of the remaining criteria for valued components of the CHAA was undertaken.

### 9.1 Assessment of Potential Triggers of Critical Habitat

The short-list of possible triggers of critical habitat that are likely to occur in the CHAA, derived from the critical habitat screening exercise (see Appendix G) are discussed in detail in the sections that follow.

#### 9.1.1 Species Potentially Triggering Criterion 1, 2 and 3 Critical Habitat

As discussed in Section 3.2.3.3 and Appendix B, quantitative thresholds for distinguishing the tiers of critical habitat have been developed by the IFC for Criteria 1 to 3. Part of these thresholds require the need to have a reasonable understanding or knowledge of the global populations of species that could trigger those criteria, as well as knowledge of their global distributions (EOO) and occupancy (AOO).

Very few of the species assessed against the screening criteria (see Table G1) had detailed population-level information available or detailed distribution records. Nevertheless, some did have the required information. Where spatial data for EOO and AOO were available (as derived from the IUCN (2017) and other references), these were used as the species' distribution owing to a need to quantify the amount of overlap of these distributions with the CHAA, and, in some instances, were used as a proxy for population level data to assess the values against the Tier 1 and Tier 2 triggers (Appendix B).

##### 9.1.1.1 Mud Snail (*Gabbiella candida*)

- Potential Critical Habitat triggered – Criterion 1 and 2

This Mud Snail is a globally listed Critically Endangered species (Table G2). It's known AOO is less than 10 km<sup>2</sup>, around the port of Butiaba on Lake Albert, and is presumed to be endemic to this area (Kyambadde 2010a). Further surveys to understand this species's EOO have not been undertaken, yet it is possible that it could occur in a much wider area, within suitable habitat.

Two sister species (Bithyniidae: *G. humerosa* and *G. walleri*) were recorded in the LSA (see Section 3.3.2.4 of the main report, Appendix D). These two species have been recorded in the same locality and habitat that *G. candida* was recorded (GBIF 2017). Therefore, it is conceivable that it could occur within the CHAA within suitable habitat.

Major threats to this species include declining habitat quality due to erosion and silting from agriculture and water pollution (Kyambadde 2010a).

Although this species has not been recorded in the CHAA, and its known AOO (that is Butiaba, about 90 km north of the Buhuka Flats) is not within the CHAA, this species may occur in the near-shore habitats off the Buhuka Flats. Suitable, near-shore aquatic habitat within the CHAA (see Section 6.3.1.1.1) for this species includes the Bugoma Lagoon (33.2 ha), large bays (73.7 ha), open sandy shores (554.8 ha) and shallow river-associated water (37.2 ha), totalling approximately 699 ha (Figure 5).

**Table G2: Population details for the Mud Snail**

Global listing*	Critically Endangered
National listing	Not listed
Restricted range*	Yes
Migratory or congregatory	No
Discrete Management Unit	CHAA (814 km <sup>2</sup> )





Global population*	Unknown
Global EOO*	10 km <sup>2</sup>
Global AOO*	10 km <sup>2</sup>
Regional/national population*	Unknown
Regional/national EOO*	10 km <sup>2</sup>
Regional/national AOO**	10 km <sup>2</sup>
Number of global discrete management units	1 (Butiaba Port)

\* Kyambadde (2010a)

Although the Mud Snail has not been recorded in the CHAA, there is a potential that it could occur, due to the presence of potentially suitable habitat. Assuming it potentially could occur in the wider Lake Albert, within suitable habitat, and knowing that it is endemic to Lake Albert, its EOO and AOO could be less than 5000 km<sup>2</sup>, based on the total area of Lake Albert (5,335 km<sup>2</sup>) and the availability of suitable habitat (that is, less than 18 m water depth) (after Kyambadde 2010b). Within the CHAA, suitable habitat for this species only occurs in the near-shore aquatic habitats, which total about 8.5 km<sup>2</sup>. Using the EOO (that is, 5000 km<sup>2</sup>) as a proxy for the global population for this species (see reasoning above), then the 8.5 km<sup>2</sup> of potential habitat within the CHAA equates to 0.2% of this species population potentially occurring in the CHAA.

Hence, applying the quantitative and qualitative triggers for Criterion 1 and 2 critical habitat (Appendix B, Table B1) to these derived population details for this species identifies that it does not trigger Criterion 2. However, there is a potential that it could occur in the CHAA, hence, because it is a Critically Endangered species, and taking a precautionary approach, this species has been assessed as triggering Criterion 1 Tier 2 critical habitat within the CHAA.

**9.1.1.2 Snail (*Bellamya rubicunda*)**

Potential critical habitat – Criterion 2

This Snail is globally listed as Near Threatened (Table G3). It is an endemic, range restricted species, known only from Lake Albert (Kyambadde 2010b). Given that it is endemic to Lake Albert, its EOO and AOO is less than 5000 km<sup>2</sup>, based on the total area of Lake Albert (5,335 km<sup>2</sup>) and the fact the species only occurs down to a maximum of 18 m depth (Kyambadde 2010b). Accordingly, it is also close to meeting the IUCN’s Endangered category (IUCN 2014); however, current known locations and threats appear to be localised and dispersed (Kyambadde 2010b).

A sister species (Viviparidae: *B. unicolor*) was recorded in the CHAA (see Section 3.3.2.4 of the main report, and Appendix D). This species has been recorded in the same locality and habitat that *B. rubicunda* was recorded (GBIF 2014). Therefore, it is conceivable that it could occur within the CHAA within suitable habitat.

Major threats to this species include declining habitat quality due to erosion and silting from agriculture and water pollution (Kyambadde 2010b).

Although this species has not been recorded in the CHAA, its EOO and AOO (that is Lake Albert) overlaps with the CHAA. Therefore, it is conceivable that this species may occur in the near-shore habitats off the Buhuka Flats. Suitable, near-shore aquatic habitat (see Section 5.3.1.1.1 of the main report) within the CHAA for this species includes: the Bugoma Lagoon (33.2 ha); large bays (73.7 ha); open sandy shores (554.8 ha); and shallow river-associated water (37.2 ha); totalling approximately 699 ha, or ~7 km<sup>2</sup> (Figure 12 of the main report).

**Table G3: Population details for the Snail**

Global listing*	Near Threatened
National listing	Not listed
Restricted range*	Yes
Migratory or congregatory	No
Discrete Management Unit	CHAA (814 km <sup>2</sup> )





Global population*	Unknown
Global EOO*	5000 km <sup>2</sup>
Global AOO*	Unknown
Regional/national population*	Unknown
Regional/national EOO*	5000 km <sup>2</sup> (Lake Albert) ~7 km <sup>2</sup> of 814 km <sup>2</sup> (CHAA)
Regional/national AOO**	Unknown
Number of global discrete management units	Unknown

\* Kyambadde (2010b)

Although the Snail has not been recorded in the CHAA, there is a potential that it could occur, for the above reasons. Within the CHAA, suitable habitat for this species only occurs in the near-share aquatic habitats, which total about 7 km<sup>2</sup>. Using the EOO (that is, 5000 km<sup>2</sup>) as a proxy for the global population for this species (see reasoning above), then the 7 km<sup>2</sup> of potential habitat within the CHAA equates to 0.1% of this species population potentially occurring in the CHAA.

Hence, applying the quantitative and qualitative triggers for Criterion 2 critical habitat (Appendix B, Table B1) to these derived population details for this species identifies that it does not trigger Criterion 2. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

### 9.1.1.3 Madagascar Pond Heron (*Ardeola idae*)

- Potential critical habitat – Criterion 1 and 3

The Madagascar Pond Heron is a globally listed Endangered species (Table G4), because of a very small breeding population limited to Madagascar that is undergoing continuing decline (BirdLife International 2017). It has a very large non-breeding range, which includes central and east Africa, including the Albertine Rift (BirdLife International 2017). Anecdotal evidence suggests that this species exhibits site fidelity and residence in suitable habitat throughout its non-breeding migratory habitat (Ndang’ang’a and Sande 2008).

Non-breeding habitat for this species includes shallow waterbodies fringed with vegetation and trees, the banks of small streams, including those inside forest (BirdLife International 2017). Its diet is broad and typical of a heron this size, that is, fish, insects, small invertebrates and vertebrates (Ndang’ang’a and Sande 2008).

Major threats to this species have primarily been noted in their breeding range, and include habitat loss from clearing, draining and converting wetland habitats to rice fields (BirdLife International, 2017).

Although this species has not been recorded in the CHAA, its non-breeding EOO overlaps the CHAA (IUCN 2017). Nevertheless, based on the presence of suitable, non-breeding habitat within the CHAA (see Table G1, Figure G1), this species may occur in the wetlands and riparian habitats of Buhuka Flats (primarily the Masika River), Bugoma Central Forest Reserve, and along the pipeline corridor.

**Table G4: Population details for Madagascar Pond Heron**

Global listing*	Endangered
National listing**	Vulnerable
Restricted range	No
Migratory or congregatory**	Yes - CMS Appendix II, Categories 1b and 1c of the AEWA
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population**	2000 – 6000 (1,300-4,000 mature individuals)
Global EOO# (non-breeding)	3,322,293 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population**	Unknown
Regional/national EOO**	Unknown





Regional/national AOO**	Unknown
Number of global discrete management units	Unknown

\* IUCN (2017)

\*\* Ndang'ang'a and Sande (2008)

\*\*\* BirdLife International (2017)

# IUCN (2014)

Although the Madagascar Pond Heron has not been recorded in the CHAA, with the nearest records around Lake Edward (~180 km south-west of the CHAA, Ndang'ang'a and Sande 2008), there is a potential that it could occur in the CHAA. Suitable non-breeding habitat for this species occurs across the CHAA, and it may not necessarily be limited to those habitats mentioned above. As such, the whole CHAA was considered to be a DMU that could support a non-breeding sub-population of this species. Furthermore, the global population of this species is estimated to be between 2000 and 6000 individuals (Ndang'ang'a and Sande 2008). The precise distribution of these individuals across the non-breeding EOO is unknown; however, Kenya and Tanzania are believed to be the core areas for its non-breeding EOO (Martínez-Vilalta et al. 2014). Therefore, a disproportionate distribution of this species across its non-breeding EOO can reasonably be assumed, with concentrations in Kenya and Tanzania (after Ndang'ang'a and Sande 2008, Martínez-Vilalta et al. 2014).

Given these uncertainties, and the lack of data, regardless of the apparent disproportionate distribution of this species across its non-breeding EOO, a conservative approach to the application of the critical habitat triggers was used. It was assumed that individuals were evenly distributed across the global (non-breeding) EOO (that is, 3,322,293 km<sup>2</sup>), which was then used as a proxy for the global non-breeding population of this species. Hence, the 814 km<sup>2</sup> of the CHAA equates to <0.01% of this species' non-breeding population.

Hence, applying the quantitative and qualitative triggers for Criterion 1 and 3 critical habitat (Appendix B, Table B1) to the derived population details for this species, identifies that it does not trigger Criterion 1 or 3. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

9.1.1.4 Grey Crowned Crane (*Balearica regulorum*)

■ Potential critical habitat – Criterion 1

The Grey Crowned-crane is a globally listed Endangered species (Table G5) (IUCN, 2017). It has a very large range that includes central, east and southern Africa, including the Albertine Rift (BirdLife International 2017).

The largest remaining populations of this species are concentrated in the northern part of their range, where approximately 30,000 to 55,000 individuals remain; with between 4000 and 5000 birds in the southern part (BirdLife International, 2017).

In East Africa, the populations tend to have peak breeding during the dry season (BirdLife International 2017). Typically, this species nests in solitary, territorial pairs at the edges of wetlands and in marshes with water at least 1 m deep and tall, emergent vegetation (Morrison and Bothma 1998, BirdLife International 2017).

The Grey Crowned-crane's diet is broad, consisting of seed heads, growing tips of grasses, pulses, nuts and grain, and small invertebrates and vertebrates (BirdLife International, 2017).

Major threats to this species are mainly from the loss of suitable breeding wetlands, pesticide use, frequent fires, changes to hydrological regimes, egg collecting and live trade/domestication of chicks (BirdLife International 2017). In Uganda, for example, indications are that that the illegal captive trade is particularly significant (Morrison 2008, 2009 in BirdLife International 2017). Additionally, electrocution and collision with overhead power lines has been identified as a serious threat in Uganda (K. Morrison in litt. 2011, J. Harris in litt. 2012).

This species' EOO substantially overlaps with the CHAA, and it has been recorded in the CHAA, on the Buhuka Flats and along the proposed pipeline route, with 14 individuals seen during the wet season surveys (see Section 6.3.2, Appendix C). Suitable breeding habitat for this species (that is, permanent wetlands (Morrison and Bothma 1998, Archibald et al. 2013)), occurs on the Buhuka Flats (~84 ha) and ~83 ha occurs in the





remainder of the CHAA (Appendix G, Figure G2). Suitable foraging habitat also occurs in the CHAA in the form of seasonally flooded grassland (73.1 ha in the Buhuka Flats and 0.6 ha along the pipeline route corridor), and open grassland (568 ha (Buhuka Flats)) (Appendix G, Figure G2). It is noted that In East Africa this species is most commonly found in human-modified habitats (Archibald et al. 2013). However, a model of habitat suitability for Grey Crowned Crane in Uganda (Stabach et al., 2009) indicates that the CHAA lies within an area of relatively low suitability for this species, more or less in between two key areas of habitat suitability – the southwestern portion of Uganda, and the area just north of Lake Albert along the Albert Nile river. Interestingly, these areas were correlated with having both the highest and lowest values of temperature seasonality in the country, as well as having suitable wetland habitat (Stabach et al., 2009). The importance of temperature variation in defining Grey Crowned Crane habitat suitability is further underlined by the reported threat of the loss of large tree roosting sites used by cranes for sheltering from the midday sun, a factor that may be crucial to crane conservation in warmer areas and during dry periods (Olupot, 2014).

Table G5: Population details for Grey Crowned Crane

Global listing*	Endangered, CITES Appendix II
National listing	Protected (Uganda Wildlife Bill 2017)
Restricted range	No
Migratory or congregatory*	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population**	50,000 – 64,000
Global EOO#	3,561,114 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population*	13,000 – 20,000
Regional/national EOO*	Unknown
Regional/national AOO*	Unknown
Number of global discrete management units	Unknown

\* IUCN (2017)

\*\* BirdLife International (2017)

# IUCN (2014)

Grey Crowned Crane has been recorded in the CHAA, with up to 14 individuals identified on the Buhuka Flats during the wet season surveys (Appendix C). It is possible that other individuals could occur within the wider CHAA, with a conservative estimate of between 15 and 20 individuals potentially occurring in this area at any one time, based on the relatively low habitat suitability of the CHAA modelled by Stabach et al. (2009), the transformation of the majority of the CHAA beyond the Buhuka Flats for agriculture, and the expected likely lack of suitable tree roosting sites in proximity to wetlands within the CHAA beyond the Buhuka Flats. As such, the whole CHAA was considered to be a DMU for this species that supports a sub-population of this species in the region. It could support 0.1% of the regional population (15 to 20 individuals of the Ugandan population of 13,000 to 20,000 individuals).

Hence, applying the quantitative and qualitative triggers for Criterion 1 critical habitat (Appendix B, Table B1) to this population identifies that it does not trigger Criterion 1. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

9.1.1.5 White-backed Vulture (Gyps africanus)

■ Potential critical habitat – Criterion 1

The White-backed Vulture is a globally listed Critically Endangered species (Table G6) (IUCN, 2017). It has a very large range from Senegal, Gambia and Mali in the west, throughout the Sahel region to Ethiopia and Somalia in the east, through East Africa into Mozambique, Zimbabwe, Botswana, Namibia and South Africa in the south (BirdLife International 2017). It is commonest and most widespread vulture in Africa, however, it is suffering rapid declines across much of this range (BirdLife International 2017, Kemp et al. 2014).





The White-backed Vulture is a gregarious species congregating at carcasses, in thermals and at roost sites. Indications are that this species migrates down the Rift Valley in Uganda in July (Kemp et al. 2014). Indeed, observations of this species made during the wet season survey of the CHAA (in May 2014, Appendix B), indicate that a group was moving down the valley.

As with most of the world’s vultures, major threats to this species are mainly from the loss of wild ungulate populations, which has led to a loss in the availability of sufficient carrion, hunting for trade, persecution and poisoning, and collisions with power lines (BirdLife International 2017). In East Africa, poisoning (especially from the highly toxic pesticide carbofuran) appears to be the main problem; although this occurs mainly outside protected areas (Kemp et al. 2014). Recent research has indicated that the use of the veterinary anti-inflammatory drug, diclofenac, has resulted in dramatic declines in vulture numbers, particularly across Asia (for example, see Green et al. 2006, Harris 2013). If this situation has affected the vultures of Africa, in particular East Africa, it is unclear at this time.

This species’ EOO substantially overlaps with the CHAA, and 20 individuals have been recorded in the CHAA (see Section 5.1.4 of the main report, Appendix C). This species prefers open wooded savannah, where it requires tall trees for nesting (BirdLife International 2017). Nests are typically located in the crown of large trees or, less often, in an open fork, frequently along a watercourse; they can be frequently clumped in loose colonies of two to 13 nests (Kemp et al. 2014). Suitable breeding and foraging habitat for this species occurs across the CHAA (Figure G3), and includes open woodland (161.7 ha), woodland (190.4 ha), riverine woodland (69.8 ha), dense wooded grassland (554.7 ha), open wooded grassland (468.9 ha), wooded grassland (184.3 ha), open woodland/dense bushland (147.9 ha), and open wooded bushland (214.0 ha).

Table G6: Population details for White-backed Vulture

Global listing*	Endangered
National listing*	Not listed
Restricted range	No
Migratory or congregatory*	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	270,000
Global EOO#	12,348,146 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population***	~40,000
Regional/national EOO*	Unknown
Regional/national AOO*	Unknown
Number of global discrete management units	Unknown

\* BirdLife International (2017)

\*\*\* Kemp et al. (2014)

# IUCN (2014)

The White-backed Vulture has been recorded in the CHAA, with up to 20 individuals identified soaring along the escarpment (Appendix C). It is possible that other individuals could occur within the wider CHAA. However, individuals of this species are very wide ranging, and are known to move up and down the Albertine Rift (Kemp et al. 2014). Therefore, it is difficult to put a precise number on the population utilising the CHAA. Regardless, the whole CHAA was considered to be a DMU for this species that supports a sub-population of this species in the region. It could support <0.05% of the regional population (20 individuals of the regional population of ~40,000 individuals).

Hence, applying the quantitative and qualitative triggers for Criterion 1 critical habitat (Appendix B, Table B1) to this population identifies that it does not trigger Criterion 1. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.





**9.1.1.6 Hooded Vulture (*Necrosyrtes monachus*)**

■ Potential critical habitat – Criterion 1

The Hooded Vulture is a globally listed Endangered species (Table G7) (IUCN, 2017). It has a very large range, tending to be confined to sub-Saharan Africa, from Senegal and southern Mauritania, through south Niger and Chad to South Sudan, Ethiopia and Somalia, to Namibia and Botswana, Zimbabwe, Mozambique and north-east South Africa (Kemp and Christie 2013).

As with most of the world’s vultures, major threats to this species are mainly from hunting for trade, persecution and poisoning (Kemp and Christie 2013, BirdLife International 2017). In Kampala, where scavenging birds were very conspicuous between the 1970s and mid-2000s, they have since noticeably decreased in numbers (Kemp and Christie 2013). Declines could also be attributed to land-use change through development, and improvements of abattoir hygiene and refuse disposal in some areas. Hooded Vultures also appear to suffer some mortality from avian influenza (H5N1), probably acquired from feeding on dead poultry (Kemp and Christie 2013). In East Africa, poisoning (especially from the highly toxic pesticide carbofuran) appears to be the main problem; although this occurs mainly outside protected areas (Kemp and Christie 2013). Recent research has indicated that the use of the veterinary anti-inflammatory drug, diclofenac, has resulted in dramatic declines in vulture numbers, particularly across Asia (for example, see Green et al. 2006, Harris 2013). If this situation has affected the vultures of Africa, in particular East Africa is unclear at this time.

This species’s EOO substantially overlaps with the CHAA; although it has not yet been recorded in the CHAA. This species prefers open woodland and savanna, also forest edge, and is generally absent from desert and dense forest, although it has been known to utilise secondary forest, clearings, settlements and urban areas (Kemp and Christie 2013). This species builds a small stick nest from April to July in the upper fork of large trees, usually deep within foliage and not on the crown (unlike other vulture species) (Kemp and Christie 2013).

Suitable breeding and foraging habitat for this species occurs across the CHAA (Figure G4), and includes open woodland (161.7 ha), woodland (190.4 ha), riverine woodland (69.8 ha), dense wooded grassland (554.7 ha), open wooded grassland (468.9 ha), wooded grassland (184.3 ha), open woodland/dense bushland (147.9 ha), and open wooded bushland (214.0 ha).

**Table G7: Population details for Hooded Vulture**

Global listing*	Endangered
National listing	Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory*	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	197,000
Global EOO#	12,369,089 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population*	Unknown
Regional/national EOO*	Unknown
Regional/national AOO*	Unknown
Number of global discrete management units	Unknown

\* BirdLife International (2017)

# IUCN (2017)

The Hooded Vulture has not yet been recorded in the CHAA (Appendix C), although it is likely that it could occur. Individuals are probably sedentary in most areas, but may range over 200 km when not breeding (Kemp and Christie 2013). Therefore, it is difficult to put a precise number on the population utilising the CHAA, or even the region. Regardless, the whole CHAA was considered to be a DMU for this species that could support a sub-population of this species in the region. Therefore, the CHAA and EOO were used as a proxy for the





population. Based on that assessment, the CHAA could support <0.01% of the global population of this species.

Hence, applying the quantitative and qualitative triggers for Criterion 1 critical habitat (Appendix B, Table B1) to this population identifies that it does not trigger Criterion 1. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

**9.1.1.7 Nahan’s Francolin (*Ptilopachus nahani*)**

■ Potential critical habitat – Criterion 1, 2 and 5

The Nahan’s Francolin is a globally listed Vulnerable species (Table G8) (IUCN, 2017). It has a very restricted distribution, being found only in north-east DRC (within the area bordered by the Aruwimi River, the Nepoko River and the Semliki River) and western and south-central Uganda (in particular the forests of Budongo, Bugoma and Mabira) (McGowan and de Juana 1994).

This species is confined to dense, mature, moist, sometimes swampy medium-altitude forest below 1,500 m (McGowan and de Juana 1994, BirdLife International 2017); and is reasonably common in Budongo Central Forest Reserve (Plumptre et al. 2010, 2011). Large trees with appropriate buttress formation are important for breeding sites for this species (Sande et al. 2009a). Forest disturbance appears to reduce the home range of this species (Sande et al. 2009b).

The population trend of this species appears to be decreasing, with the primary threats thought to be habitat loss through logging and clearance of forest for charcoal burning and agriculture, particularly within Bugoma Central Forest Reserve (BirdLife International 2017). Fragmentation alone probably does not appear to adversely affect the species, but it does appear to be affected by habitat changes associated with human-induced fragmentation, such as the extensive removal of large trees (BirdLife International 2012ad, 2014i).

It appears to have a very restricted EOO, although populations in the wider DRC are unknown, hence its distribution may be larger than thought (BirdLife International 2017). This species’s EOO overlaps with the CHAA; and has been recorded in the CHAA within Bugoma Central Forest Reserve (Plumptre et al. 2010, 2011). Suitable habitat for this species occurs only in Bugoma Central Forest Reserve (40,243 ha), within the CHAA (Figure G5).

**Table G8: Population details for Nahan’s Francolin**

Global listing*	Vulnerable
National listing	Not listed
Restricted range	Yes
Migratory or congregatory*	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	50,000 - 99,999
Global EOO#	100,339 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population*	44,038 (95% CI: 32,827-59,079)
Regional/national EOO	1046 km <sup>2</sup> (Budongo (435 km <sup>2</sup> ), Bugoma Forest (401 km <sup>2</sup> ) and Mabira Forest (210 km <sup>2</sup> ))
Regional/national AOO	1046 km <sup>2</sup> (Budongo (435 km <sup>2</sup> ), Bugoma Forest (401 km <sup>2</sup> ) and Mabira Forest (210 km <sup>2</sup> ))
Number of global discrete management units	3 (Budongo (435 km <sup>2</sup> ), Bugoma Forest (401 km <sup>2</sup> ) and Mabira Forest (210 km <sup>2</sup> ))

\* IUCN (2017)

\*\* BirdLife International (2017)

# IUCN (2014)







The Nahan’s Francolin has been recorded in the CHAA (Plumptre et al. 2010, 2011). Although the precise number of individuals occurring within the CHAA is unknown, the CHAA does support approximately 38% of this species’s known regional AOO (Table G8). Therefore, it is conceivable that the CHAA could support ~16,700 individuals (38% of 44,038).

Hence, applying the quantitative and qualitative triggers for Criterion 1 critical habitat (Appendix B, Table B1) to this population identifies that it triggers only Criterion 1 Tier 1. Therefore, this species has been assessed as triggering critical habitat within the CHAA.

**9.1.1.8 Eastern Chimpanzee (*Pan troglodytes schweinfurthii*)**

■ Potential critical habitat – Criterion 1

The Eastern Chimpanzee is a globally listed Endangered species (IUCN, 2017; Wilson et al. 2008) (Wilson et al. 2008). Generally, Chimpanzees have a wide, but discontinuous distribution, across Africa, yet tend to be confined to the equatorial belt (Oates et al. 2008). The Eastern Chimpanzee ranges from the Ubangi River/Congo River in Central African Republic and the DRC, to western Uganda, Rwanda and western Tanzania (Wilson et al. 2008).

They prefer mature moist and dry forests, either evergreen or semi-deciduous, and forest galleries extending into savanna woodlands (Wilson et al. 2008). They are omnivorous, and their diet is highly variable according to individual populations and seasons. Chimpanzees form social communities with home ranges larger in woodland forest mosaics than in mixed forest, averaging 12.5 km<sup>2</sup> (range 5 to 400 km<sup>2</sup>) (Wilson et al. 2008).

Due to high levels of exploitation, loss of habitat and habitat quality due to expanding human activities, this species is estimated to have experienced a significant population reduction in the past 20 to 30 years (Wilson et al. 2008). Major threats include habitat destruction and degradation (slash and burn agriculture, deforestation, logging), poaching (bush meat, pet trade, traditional medicine, crop protection), and disease (Oates et al. 2008).

This species’s EOO overlaps with the CHAA; and it has been recorded in the CHAA, in particular, Bugoma Central Forest Reserve (Plumptre et al. 2010, 2011). Suitable habitat for this species occurs only in Bugoma Central Forest Reserve (401 km<sup>2</sup>), within the CHAA (Figure G6).

**Table G9: Population details for Eastern Chimpanzee**

Global listing*	Endangered, CITES Appendix I
National listing*	VU, Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	172,700 - 299,700
Global EOO#	5,759,594 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population**	4000 - 5700
Regional/national EOO	1046 km <sup>2</sup> (Budongo (435 km <sup>2</sup> ), Bugoma Forest (401 km <sup>2</sup> ) and Mabira Forest (210 km <sup>2</sup> ))
Regional/national EOO	1046 km <sup>2</sup> (Budongo (435 km <sup>2</sup> ), Bugoma Forest (401 km <sup>2</sup> ) and Mabira Forest (210 km <sup>2</sup> ))
Number of global discrete management units	Unknown

\* Oates et al. (2008)

\*\* Thompson and Wrangham (2013)

# IUCN (2014)





The Eastern Chimpanzee has been recorded in the CHAA (Plumptre et al. 2010, 2011). Although the precise number of individuals occurring within the CHAA is unknown, the CHAA does support approximately 38% of this species known regional EOO (Table G9). Therefore, it is conceivable that the CHAA could support between 1520 and 2160 individuals (38% of 4000 to 5700).

Eastern Chimpanzees are great apes, therefore, under the quantitative and qualitative triggers for Criterion 1 critical habitat (Appendix B, Table B1) they trigger Criterion 1 Tier 1. Therefore, this species has been assessed as triggering critical habitat within the CHAA.

**9.1.1.9 Migratory and Congregatory Species**

**Migratory Birds**

- Potential critical habitat – Criterion 3

The CHAA supports a diversity of migratory and congregatory species as listed under the Convention on Migratory Species (CMS), to which Uganda is a signatory (see Section 4.0 of the main report). All of these species are listed under Appendix II of the CMS. Species listed under Appendix II are identified as having an unfavourable conservation statuses, that is, they may be threatened or near threatened, and, therefore, need international agreements for their conservation and management within the countries within which they are known to range (CMS 2014).

At least 12 Palearctic migratory birds could occur in the CHAA (see Section 5.1.4 of the main report, Appendix C, Table G1). These include, amongst others, Common Ringed Plover, Kittlitz’s Plover, White-winged Tern, Great Snipe, Black-winged Pratincole, Collared Pratincole, Black-tailed Godwit, Eurasian Curlew, African Skimmer, Common Sandpiper, African Wattled Lapwing, and Spur-winged Lapwing. Although individual numbers of these species were low during the surveys conducted (see Section 5.1.4 of the main report), it could be expected that the CHAA may support substantial numbers of these species during peak migration times. In particular, around September and October; a survey for migratory species is planned for that time to gauge the importance of the CHAA for migratory birds. Nevertheless, the CHAA supports suitable habitat for all of the aforementioned species. In particular, the Buhuka Flats supports ~12 km of shoreline (Figure G7), which is favourable to wading and shore birds like the Common Ringed Plover, Kittlitz’s Plover, White-winged Tern, Great Snipe, Black-tailed Godwit, Eurasian Curlew, African Skimmer, and Common Sandpiper. The flats also support ~850 ha of grassland and ~77 ha of seasonally flooded grassland (Figure 4a and b of the main report), that is favoured habitat of Black-winged Pratincole, Collared Pratincole, African Wattled Lapwing, and Spur-winged Lapwing.

**Table G10: Population details for Common Ringed Plover**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	360,000 - 1,300,000
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)





**Table G11: Population details for Kittlitz’s Plover**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G12: Population details for White-winged Tern**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	2,500,000 - 4,500,000
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G13: Population details for Great Snipe**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	465,000 - 1,040,000
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)





**Table G14: Population details for Black-winged Pratincole**

Global listing*	Near Threatened
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	220,000 - 290,000
Global EOO#	9,354,763 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	4,498,364 km <sup>2</sup>
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G15: Population details for Collared Pratincole**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G16: Population details for Black-tailed Godwit**

Global listing*	Near Threatened
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	427,000 – 805,000
Global EOO#	45,772,340 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	11,050,681 km <sup>2</sup>
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)





**Table G17: Population details for Eurasian Curlew**

Global listing*	Near Threatened
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	77,000 - 1,065,000
Global EOO#	31,500,728 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	9,201,510 km <sup>2</sup>
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G18: Population details for African Skimmer**

Global listing*	Near Threatened
National listing*	Protected (Uganda Wildlife Bill 2017)
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	15,000 - 25,000
Global EOO#	10,384,709 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	8,000 - 12,000
Regional/national EOO#	10,384,709 km <sup>2</sup>
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G19: Population details for Common Sandpiper**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	2,600,000 - 3,200,000
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)





**Table G20: Population details for African Wattled Lapwing**

Global listing*	Least Concern
National listing*	Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

**Table G21: Population details for Spur-winged Lapwing**

Global listing*	Least Concern
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	NA
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	NA
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* IUCN (2017) # IUCN (2014)

Applying the quantitative and qualitative triggers for Criterion 3 critical habitat (Appendix B, Table B1) to the population details of each of these species in relation to the CHAA (see Table G10 to G21) and the known baseline (see Appendix C), identifies that none of them triggers Criterion 3. Therefore, these species have been assessed as not triggering critical habitat within the CHAA.

**Congregatory Bats**

■ Potential critical habitat – Criterion 3

The CHAA potentially supports populations of the migratory and congregatory African Straw-coloured Fruit-bat (*Eidolon helvum*), a Bent-wing Bat (*Miniopterus* sp.) and Large-eared Free-tailed Bat (*Otomops martiensseni*). Although these species do not migrate across vast distances and continents like birds, they can migrate hundreds of kilometres from hibernacula to breeding roosts (Monadjem et al. 2010). No roosting colonies of African Straw-coloured Bat were recorded in the CHAA, and it is more likely that individuals or small colonies of this gregarious species may occur in the CHAA. Certainly, it is doubtful that roosts would occur in the LSA. The Bent-wing Bats and the Large-eared Free-tailed Bat are obligate cavity or cave roosters (Dietz et al. 2009, Monadjem et al. 2010, Happold 2013b, Yalden and Happold 2013). Although the Large-eared Free-tailed Bat has not yet been recorded in the CHAA, a Bent-wing Bat has been recorded. This is interesting because Bent-wing Bats, being an obligate cave roosting species, which can form roosting colonies





numbering thousands of individuals (Monadjem et al. 2010), are also known to share its roosting colonies with other species of bats, such as the Large-eared Free-tailed Bat (Large-eared Free-tailed Bat). As such, it is conceivable that one or more caves occur in the CHAA where-in these species may roost. The precise locality of these caves is not known; however, these could, more than likely occur somewhere along the escarpment.

**Table G22: Population details for African Straw-coloured Fruit-bat**

Global listing*	Near Threatened
National listing	Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown, suspected to be millions
Global EOO#	12,945,414 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	Unknown
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* Mickleburgh et al. (2008a)

# IUCN (2014)

The African Straw-coloured Fruit-bat has not yet been recorded in the CHAA, and the global population of this species is unknown; therefore, the global EOO and the CHAA were used as proxies for the global and local populations, respectively. The CHAA potentially supports <0.01% of the global population.

Hence, applying the quantitative and qualitative triggers for Criterion 3 critical habitat (Appendix B, Table B1) to the population details of African Straw-coloured Fruit-bat in relation to the CHAA (see Table G22) identifies that it does not trigger Criterion 3. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

**Table G23: Population details for Large-eared Free-tailed Bat**

Global listing*	Near Threatened
National listing*	-
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	7,649,368 km <sup>2</sup>
Global AOO#	NA
Regional/national population*	Unknown
Regional/national EOO#	Unknown
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* Mickleburgh et al. (2008c)

# IUCN (2014)





The Large-eared Free-tailed Bat has not yet been recorded in the CHAA, and the global population of this species is unknown; therefore, the global EOO and the CHAA were used as proxies for the global and local populations, respectively. The CHAA potentially supports 0.01% of the global population.

Hence, applying the quantitative and qualitative triggers for Criterion 3 critical habitat (Appendix B, Table B1) to the population details of Large-eared Free-tailed Bat in relation to the CHAA (see Table G23) identifies that it does not trigger Criterion 3. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

**African Elephant (*Loxodonta africana*)**

■ Potential critical habitat – Criterion 3

The African Elephant is also identified by the CMS as a migratory species (Table G1). This species occurs in the CHAA, and individuals have been recorded in the Bugoma Central Forest Reserve (Plumptre et al. 2010, 2011). However, the records suggest that the individuals occurring in the Bugoma Central Forest Reserve are lone, or single individuals (see Plumptre et al 2010, 2011), and certainly not large breeding herd that would migrate.

**Table G24: Population details for African Elephant**

Global listing*	Vulnerable, CMS Appendix II, CITES Appendix I
National listing*	Vulnerable; Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory	Yes
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population*	Unknown
Global EOO#	3,543,323 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population*	2000 - 3000
Regional/national EOO#	103,983 km <sup>2</sup>
Regional/national AOO#	NA
Number of global discrete management units	unknown

\* Blanc (2008)

# IUCN (2014)

The African Elephant has been recorded in the CHAA, although it appears that those were lone individuals. The precise population within the CHAA is unknown, as is the global population; although the population within Uganda is estimated to be between 2000 and 3000 individuals (Table G24). Therefore, the regional EOO and the CHAA were used as proxies to estimate the regional and local populations, respectively. The CHAA potentially supports 0.8% of the regional population.

Hence, applying the quantitative and qualitative triggers for Criterion 3 critical habitat (Appendix B, Table B1) to the population details of African Elephant in relation to the CHAA (see Table G24) identifies that it does not trigger Criterion 3. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

**9.1.2 Ecosystems Potentially Triggering Criterion 4 Critical Habitat: Highly Threatened and/or Unique Ecosystems**

Highly threatened or unique ecosystems are those (i) that are at risk of significantly decreasing in area or quality; (ii) with a small spatial extent; and/or (iii) containing unique assemblages of species including assemblages or concentrations of biome-restricted species (IFC, 2012).

In addition, areas determined to be irreplaceable or of high priority/significance based on systematic conservation planning techniques carried out at the landscape and/or regional scale by governmental bodies,







recognized academic institutions and/or other relevant qualified organizations (including internationally-recognized NGOs) or that are recognized as such in existing regional or national plans, such as the NBSAP, would qualify as critical habitat per Criterion 4 (IFC, 2012).

### 9.1.2.1 Bugoma Central Forest Reserve

Bugoma Central Forest Reserve therefore triggers Critical Habitat on the basis of the following qualifying factors:

- i) It is at risk of significantly decreasing in area and quality as a result of human encroachment as a result of demand for ecosystem services by the local communities, as well as recent illegal land-grabbing activity - in 2016, the Muhangaizima Block was reportedly leased for sugar cane cultivation, with subsequent transformation activities affecting approximately 8000 ha of the CFR (ACBF, 2016);
- ii) It is conservatively assessed as Vulnerable (after Rodriguez et al. 2011) – suspected of undergoing a ≥30% decline in extent of occurrence over the last 50 years in the region (based on Plumptre 2002, Plumptre et al. 2003, 2007, 2010, 2011);
- iii) It supports a relatively unique assemblage of species including Uganda’s only population of the range-restricted Nahan’s Francolin (*Ptilopachus nahani*), as well as Endangered Eastern Chimpanzee (*Pan troglodytes schweinfurthii*), and Vulnerable African Elephant (*Loxodonta africana*) for which it is also of recognised importance as a climate change refugium; is a recognised chimpanzee conservation unit (Plumptre et al. 2010), and is a recognised area of old growth forest (Plumptre et al. 2010, 2011); and
- iv) It is recognised as an Important Bird Area by the internationally-recognised NGO BirdLife International.

### 9.1.3 Species Potentially Triggering Criterion 5 Critical Habitat (Evolutionarily Distinct Species)

As discussed in Section 3.2.3.3 and Appendix B, quantitative thresholds for distinguishing the tiers of critical habitat have been developed by the IFC for Criteria 1 to 3, and the thresholds for Criterion 5 are purely qualitative.

This criterion therefore is defined by the presence within the CHAA of subpopulations of species that are phylogenetically or morphogenetically distinct and may be of special conservation concern given their distinct evolutionary history, including ‘Evolutionarily Distinct and Globally Endangered’ (EDGE) species (ZSL, 2017).

#### 9.1.3.1 Shoebill (*Balaeniceps rex*)

The Shoebill is a globally listed Vulnerable species (Table G25) (IUCN, 2017). It has a very wide distribution, from South Sudan to Zambia, but is very locally distributed within that range and prefers large swamps (BirdLife International 2017).

This species is a true wetland specialist that is sedentary as an adult. It breeds and forages in seasonally flooded marshes where vegetation is dominated by a mixture of Papyrus (*Cyperus papyrus*), reeds (*Phragmites* spp.), cattails (*Typha* spp.) and grasses, particularly *Miscanthidium*, although it tends to avoid areas where the vegetation is taller than itself (BirdLife International 2017). Suitable habitat for breeding is very limited within the CHAA; however, it is conceivable that non-breeding individuals may occur in suitable habitat (seasonally-flooded marshes) within the CHAA.

In Uganda, this species shows a preference for feeding on Lungfish (*Protopterus aethiopicus*), although it does take a variety of species including Senegal Bichir, catfish (*Clarias* spp.) and tilapia (*Tilapia* spp.) (BirdLife International 2017), all species common in the CHAA (Appendix C).

Major threats to this species are mainly from the loss of suitable breeding wetlands, general habitat destruction and degradation, disturbance, hunting, and capture for the bird trade (BirdLife International 2017).

This species’ EOO does not overlap with the CHAA, and it has not been recorded in the CHAA (see Table G1).



**Table G25: Population details for Shoebill**

Global listing*	Vulnerable, CITES Appendix II
National listing*	Vulnerable; Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory*	No
Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population**	<10,000
Global EOO#	Unavailable at time of writing
Global AOO#	Unknown
Regional/national population**	100 – 150
Regional/national EOO#	Unknown
Regional/national AOO#	Unknown
Number of global discrete management units	Unknown

\* IUCN (2017)

\*\* BirdLife International (2017)

# IUCN (2014)

Hence, applying the qualitative triggers for Criterion 5 critical habitat (Appendix B) to the population details of Shoebills in relation to the CHAA (see Table G25) identifies that it does not trigger Criterion 5. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

### 9.1.3.2 Secretarybird (*Sagittarius serpentarius*)

The Secretarybird is a globally listed Vulnerable species (Table G26) (IUCN, 2017). It has a very wide distribution, from West Africa (Senegal and the Gambia) across to Ethiopia and Somalia, and south to South Africa (Kemp et al. 2014).

It prefers grassland or tree and grass savanna, favouring short grass with scattered *Acacia* thorn trees to provide sites for roosting and nesting; it is also found in large-scale agricultural areas and sub-desert areas; it rarely visits clearings in woodland or forest (BirdLife International 2017, Kemp et al. 2014). They tend to be sedentary (with regular seasonal breeding) in some areas, but nomadic in most areas in response to changes in rainfall, grazing and fires (Kemp et al. 2014).

Secretarybirds breed at any time of year, whenever food is abundant; build large nests on top of low trees, often flat-topped Acacias (Kemp et al. 2014).

This species can be common to rare and localised across its range, with recent evidence suggesting rapid decline; within Uganda it is never common, and is now largely confined to national parks (Kemp et al. 2014). Major threats coming from the excessive burning of grasslands, which may suppress populations of prey species, intensive grazing of livestock is also probably degrading otherwise suitable habitat (BirdLife International 2017). Disturbance by humans could affect breeding, while live trade of individuals has also been reported, along with indiscriminate poisoning at waterholes as possible threats (BirdLife International 2017).

This species' EOO overlaps substantially with the CHAA, yet it has not yet been recorded in the CHAA (see Appendix G). Suitable breeding and foraging habitat for this species occurs across the wider CHAA, and the CHAA, and includes open woodland, woodland, open wooded grassland, wooded grassland, and open grassland (Table 6).

**Table G26: Population details for Secretarybird**

Global listing*	Vulnerable, CITES Appendix II
National listing*	Protected (Uganda Wildlife Bill, 2017)
Restricted range	No
Migratory or congregatory*	No





Discrete management unit	CHAA (814 km <sup>2</sup> )
Global population**	Unknown
Global EOO#	15,137,802 km <sup>2</sup>
Global AOO#	Unknown
Regional/national population**	100 – 150
Regional/national EOO#	Unknown
Regional/national AOO#	Unknown
Global number of discrete management units	

\* BirdLife International (2013h)

\*\* BirdLife International (2014j)

# IUCN (2014b)

Hence, applying the qualitative triggers for Criterion 5 critical habitat (Appendix B) to the population details of Secretarybirds in relation to the CHAA (see Table G26) identifies that it does not trigger Criterion 5. Therefore, this species has been assessed as not triggering critical habitat within the CHAA.

### 9.1.4 Species/Ecosystems triggering other Qualitative Critical Habitat Criteria

The near-shore habitats are both important fishing grounds and nursery habitat for fisheries, that support 11 fishing villages on the Buhuka Flats and surrounds (see Chapter 18 Ecosystem Services assessment). This triggers the qualitative Criterion 13 as described in Appendix G.

Furthermore, the near-shore habitats of Lake Albert within the CHAA could potentially constitute critical habitat under Criterion 1 if the mud snail *G. candida* is confirmed present in this area.

## 9.2 Confirmed Triggers of Critical Habitat in the CHAA

Table 4 presents a summary of the confirmed triggers of critical habitat in the CHAA. The spatial representation of critical habitat within the CHAA is presented in Figure 16.





## BIODIVERSITY IMPACT ASSESSMENT

**Table 4: Triggers of critical habitat in the CHAA**

Valued Component	Potential triggering criteria*	Critical Habitat Designation**	Habitat and reasoning
Mud Snail ( <i>Gabbiella candida</i> )	1 and 2	Criterion 1 Tier 1	<ul style="list-style-type: none"> <li>■ Could occur on near-shore aquatic habitats (Bugoma Lagoon, large bays, open sandy shores, shallow river-associated water)</li> <li>■ See Appendix G for precise reasoning</li> </ul>
Nahan's Francolin ( <i>Ptilopachus nahani</i> )	2	Criterion 2 Tier 2	<ul style="list-style-type: none"> <li>■ Occurs in Bugoma Central Forest Reserve, possibly one of less than 10 DMUs globally (including DRC)</li> <li>■ Potential for CHAA to support &gt;10% of this species' known global population</li> <li>■ See Appendix G for precise reasoning</li> </ul>
Eastern Chimpanzee ( <i>Pan troglodytes schweinfurthii</i> )	1	Criterion 1 Tier 1	<ul style="list-style-type: none"> <li>■ Occurs in Bugoma Central Forest Reserve</li> <li>■ Great apes are an iconic species (see GN 74 and footnotes in PS6, IFC 2012a and b)</li> <li>■ See Appendix G for precise reasoning</li> </ul>
Bugoma Central Forest Reserve	4, 6, 7, 9, 11, 12, 13, 15, 16	Criterion 4	<ul style="list-style-type: none"> <li>■ Threatened ecosystem – over 110 km<sup>2</sup> of forest surrounding the Bugoma CFR boundary has been cleared since the mid-1980s (Plumptre 2002); and in 2016, the Muhangaizima Block was reportedly leased for sugar cane cultivation, with subsequent transformation activities affecting approximately 8000 ha of the CFR (ACBF, 2016).</li> <li>■ Therefore, conservatively assessed as Vulnerable (after Rodriguez et al. 2011) – suspected of undergoing a ≥30% decline in extent of occurrence over the last 50 years in the region (based on Plumptre 2002, Plumptre et al. 2003, 2007, 2010, 2011)</li> <li>■ Of recognised importance as a climate change refugium for Vulnerable Nahan's Francolin and Endangered Eastern Chimpanzee (Ayebare et al. 2013), and a recognised chimpanzee conservation unit (Plumptre et al. 2010)</li> <li>■ Recognised area of old growth forest (Plumptre et al. 2010, 2011)</li> <li>■ Supports a population of Eastern Chimpanzee (McLennan 2008, Plumptre et al. 2003, 2010, 2011) that is recognised as being one for the four largest in the region (Plumptre et al. 2010); apart from being an Endangered species, chimpanzees are</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

			<p>also recognised as key stone species and ecosystem engineers (Chapman et al. 2013)</p> <ul style="list-style-type: none"> <li>■ The forest is recognised for its unique biodiversity values, including biome restricted species (Plumptre et al. 2011)</li> <li>■ Local people harvest timber, fibre, fuel wood and charcoal, and non-timber forest products from the forest (Plumptre 2002)</li> <li>■ Bugoma Central Forest Reserve is recognised as an Important Bird Area (BirdLife International 1998, IUCN 2010, 2014b)</li> <li>■ Recognised as a high conservation priority area (NEMA 2010)</li> </ul>
Near-shore habitats of Lake Albert	13	Criterion 13	<ul style="list-style-type: none"> <li>■ The near-shore habitats are important fishing grounds that support 11 fishing villages on the Buhuka Flats and surrounds (see Chapter 18 Ecosystem Services Review)</li> </ul>

\* IFC (2012b)

\*\* In instances where more than one potential criterion could be triggered, only the highest-level designation is presented



# BIODIVERSITY IMPACT ASSESSMENT




Figure 16: Critical habitat in the CHAA





## 10.0 IMPACT ASSESSMENT – PRODUCTION FACILITY

This section presents an assessment of the possible interactions of biodiversity valued components with the production facility infrastructure and activities, and the resulting impacts during the construction, operation and decommissioning phases of the Project.

The biodiversity valued components for the Production Facility impact assessment are listed in Table 5 below. They include all of the species and habitats that trigger critical habitat designation within the CHAA. In addition, ecosystems of concern that will be potentially affected by the Project, and Grey Crowned Crane, were also included as valued components for impact assessment, for reasons outlined in the Table below. As mentioned in Section 10.1.2, potential impacts to other species of concern are assessed at the habitat level (ecosystems of concern).

**Table 5: Biodiversity Valued Components for Impact Assessment**

Valued Component	Confirmed CH Trigger? (see Table 4)	Reasoning (see Table 4)
Near-shore aquatic habitats of Lake Albert	<ul style="list-style-type: none"> <li>■ Yes – Criterion 13</li> <li>■ Possibly Criterion 1 and Criterion 2 (G. candida)</li> </ul>	<ul style="list-style-type: none"> <li>■ The near-shore habitats are important fishing grounds that support 11 fishing villages on the Buhuka Flats and surrounds (see Chapter 18 Ecosystem Services Review)</li> <li>■ May support the CR and range-restricted species <i>Gabbiella candida</i></li> </ul>
Wetlands	<ul style="list-style-type: none"> <li>■ No</li> </ul>	<ul style="list-style-type: none"> <li>■ Natural habitat – priority habitat according to IFC (2012)</li> <li>■ Supports Endangered Grey Crowned Crane</li> <li>■ Important in supply of ecosystem services to local communities (see Chapter 18 Ecosystem Services Review)</li> </ul>
Escarpment vegetation corridor	<ul style="list-style-type: none"> <li>■ No</li> </ul>	<ul style="list-style-type: none"> <li>■ Natural habitat – priority habitat according to IFC (2012)</li> <li>■ Forms part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor</li> <li>■ The location of caves and cavities along the escarpment that could be important for cavity-roosting bats</li> </ul>
Bugoma Central Forest Reserve	<ul style="list-style-type: none"> <li>■ Yes –</li> <li>■ Criterion 4</li> <li>■ Criterion 1</li> <li>■ Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>■ Triggers CH on the basis of being a highly threatened and unique ecosystem (Criterion 4)</li> <li>■ Triggers Criterion 1 Tier 1 CH on the basis of support of a population of Eastern Chimpanzee, that is recognised as being one for the four largest in the region; apart from being an Endangered species, chimpanzees are also recognised as key stone species and ecosystem engineers</li> <li>■ Triggers Criterion 2 Tier 2 CH on the basis of support of range-restricted Nahan’s Francolin</li> <li>■ Recognised area of old growth forest</li> </ul>





Valued Component	Confirmed CH Trigger? (see Table 4)	Reasoning (see Table 4)
		<ul style="list-style-type: none"> <li>■ The forest is recognised for its unique biodiversity values, including biome restricted species</li> <li>■ Is an important ecosystem service supply area for local people who harvest timber, fibre, fuel wood and charcoal, and non-timber forest products from the forest</li> <li>■ Bugoma Central Forest Reserve is recognised as an Important Bird Area</li> <li>■ Nationally recognised as a high conservation priority area (NEMA 2010)</li> </ul>
Mud Snail ( <i>Gabbiella candida</i> )	<ul style="list-style-type: none"> <li>■ Possibly Criterion 1 and Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>■ Could occur on near-shore aquatic habitats (Bugoma Lagoon, large bays, open sandy shores, shallow river-associated water)</li> <li>■ Has not been confirmed in LSA to date and is included on basis of precautionary principle</li> </ul>
Grey Crowned Crane	<ul style="list-style-type: none"> <li>■ No</li> </ul>	<ul style="list-style-type: none"> <li>■ Although Grey Crowned Crane is not present in numbers that would trigger CH designation, it is an Endangered species and has been confirmed present on the Buhuka Flats during baseline fieldwork in 2014 and 2017</li> <li>■ Any potential Project impacts on a globally-recognised and nationally-protected Endangered species are unacceptable and warrant addressing via the impact assessment process</li> </ul>
Nahan's Francolin ( <i>Ptilopachus nahan</i> )	<ul style="list-style-type: none"> <li>■ Yes – Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>■ Occurs in Bugoma Central Forest Reserve, possibly one of less than 10 DMUs globally (including DRC)</li> <li>■ Potential for CHAA to support &gt;10% of this species' known global population</li> </ul>
Eastern Chimpanzee ( <i>Pan troglodytes schweinfurthii</i> )	<ul style="list-style-type: none"> <li>■ Yes – Criterion 1</li> </ul>	<ul style="list-style-type: none"> <li>■ Occurs in Bugoma Central Forest Reserve</li> <li>■ Great apes are an iconic species of anthropological and evolutionary significance</li> <li>■ They generally immediately trigger CH designation (see GN 74 and footnotes in PS6, IFC 2012a and b)</li> </ul>

## 10.1 Construction and Decommissioning Phase Impacts

The anticipated impacts are expected to be similar for the construction and decommissioning phases, and will occur over a similar duration for both phases (that is, ~2 years). Therefore, for the intents of this impact assessment, the decommissioning phase impacts have been included with the construction phase impacts in their assessment.

The predicted impacts to valued components include: direct loss of habitat due to land take for the Project infrastructure (Table 6); sensory disturbances (from noise, vibration, light, and odour); changes to surface water quality and flows, air emissions and the associated potential for pollution; erosion and sedimentation; and direct mortality of species of concern from vehicle movements and site preparation.







The predicted impacts are assessed in two broad categories:

- Impacts on habitat and ecosystem integrity; including the near-shore environment of Lake Albert, the vegetation corridors along the escarpment, and wetlands in the LSA and CHAA.
- Impacts on species of concern, specifically the Mud Snail *Gabbiella candida*.

For the assessment of impacts during the construction and decommissioning phase, the key questions were divided into sub-questions that focused on individual valued components within the CHAA and LSA. In answering each question, the individual components of the Project were considered with regards to their potential to affect a valued component. These questions are presented below.

**10.1.1 What impact could the construction/decommissioning of the Project have on habitats and ecosystem integrity?**

This section presents the assessment of impacts that the construction and decommissioning of the Project could have on the habitat and ecosystem integrity within the CHAA and the LSA. These habitats either do, or could, support populations of species of concern. Therefore, the assessment of potential impacts to those species, and others, occurring in the CHAA has been assessed in this section through the determination of the impacts to potential habitat (Table 6) for those species.

The impacts of the Project on critical habitat, as triggered by species of concern, are covered under the individual assessment of those species in Section 9.1.2. Other triggers of critical habitat are discussed, as relevant, in the appropriate sections.

**Table 6: Area of vegetation communities at baseline, disturbance and loss due to Production Facility, and net change (% loss) in CHAA**

Community	Baseline CHAA (ha)*	Baseline Buhuka Flats (ha)	Loss to already permitted/constructed infrastructure	Loss to proposed infrastructure (ha)	% Loss CHAA	% Loss Buhuka Flats
Bare Ground	24.0	5.0	1.9	0.3	9.0%	43.3%
Dense Bushland	1097.6	53.3	-	0.6	0.1%	1.1%
Dense Wooded Grassland	613.2	2.2	-	-	-	-
Grassland with Thicket	1101.1	255.6	4.2	6.7	1.0%	4.3%
Lagoons	33.2	32.5	-	0.1	0.6%	0.6%
Large Bays	73.7	2.7	-	-	-	-
Open Bushland and Shrubland	2896.3	27.5	0.1	0.2	0.0%	1.2%
Open Grassland	568.5	518.2	30.3	50.8	14.3%	15.6%
Open Sandy Shores	554.8	3.3	-	-	-	-
Permanent Wetland	83.8	83.7	-	-	-	-
Riverine Bushland	640.3	35.6	-	-	-	-
Riverine Woodland and Bushland	76.8	73.9	-	-	-	-
Rocky Escarpments	101.1	0.2	-	-	-	-
Seasonal Wetland	85.3	67.8	0.8	2.8	4.2%	5.3%





Settlement	207.9	142.2	0.3	2.8	1.5%	2.1%
Shallow River-Associated Waters	37.2	21.2	-	-	-	-

**10.1.1.1 What impact could the construction/decommissioning of the Project have on the near-shore aquatic habitats of Lake Albert?**

**Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the near-shore environment of Lake Albert were changes in: regional representativeness; topography (geomorphology) and sediments; water quality; ecosystem composition; ecosystem configuration.

Loss of habitat due to direct disturbance and clearing associated with the Project was quantified by overlaying the current, baseline extent of the habitat with the Project footprint.

Additional, indirect impacts to habitat were estimated by applying a 1 km buffer to the Project footprint, forming the LSA. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by sensory disturbance, changes in water quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential succession changes in species composition that could occur. This was accomplished by examining available literature about the ecology of Lake Albert, and scientific literature about the impacts of human activities on aquatic environments.

**Impact Assessment**

■ Representativeness

The CHAA supports 16.2 km of near-shore aquatic habitats, equating to about 810 ha (based on the near-shore habitat extending 0.50 km into the lake, see Section 7.1.1). Loss and degradation of these communities as a result of the Project has already occurred through the upgrade of the jetty area. Approximately 0.12 km of open sandy shoreline, extending 20 m into the lake, has been physically lost or severely disturbed by the construction of the jetty. This represents ~1% of the near-shore habitat within the CHAA, which is, consequently, a very localised impact, which is considered to be within the expected range of natural disturbance perturbations for the shoreline of Lake Albert. For example, through extreme weather events, lake level rise and fall, and longshore drift. The loss of this proportion of sandy shoreline community is, therefore, considered negligible in relation to the regional representativeness of this community; that is, a loss of 0.12 km of ~68 km (or 0.2%) of regional community.

The impact will be long-term, but largely reversible after decommissioning. Taking all factors into account, the magnitude of loss and disturbance will be low. However, this must be weighed against the very high sensitivity of the lake ecosystem, being a fresh water body that supports a high diversity of endemic and commercially important species of fish, threatened snails (the Critically Endangered mud snail, *Gabbiella candida*, and the Near Threatened Snail *Bellamya rubicunda*), threatened reptiles (the Vulnerable African Soft-shelled Turtle *Trionyx triunguis*) and threatened birds (the Endangered Grey-crowned Crane *Balearica regulorum*), as well as a shoreline important for many species of migratory birds. As such, as identified in Section 9.0, this community triggers Tier 1 and Tier 2 critical habitat.

■ Topography (geomorphology) and sediment transport

During construction, the existing jetty - consisting of a solid concrete structure extending some 20 m into the lake to provide sufficient draught during low water periods – will be upgraded; however, no material changes in the dimensions of the structure are anticipated. Approximately 1 km north along the shoreline beside Well Pad 2, a water intake and water extraction pump station that also extends ~20 m into the lake is proposed for construction, which will see the alteration of ~0.12 km of open sandy shoreline.





The existing jetty's influence on the physical structure of the adjacent shoreline, and subsequent effects on the local geomorphology and longshore drift that maintain the shoreline, are not well understood; however, examination of recent aerial imagery suggests that some accretion of sediment on the eastern side of the jetty, and some erosion on the western side, is evident. Sediment drift is recognised as an important driver in shoreline ecosystems, contributing to the nutrient input that drives phytoplankton, zooplankton and fish communities (Parks et al. 2013). Various studies have shown that structures constructed on shorelines may disrupt hydrodynamic flow patterns, creating a barrier for sediment movement along the shore that can have a number of effects on faunal communities such as altered patterns of larval supply and food availability, and subsequent indirect effects on distribution and abundance of fish, turtles and birds through habitat modification and loss (in Walker et al. 2008). However, the zone of influence of structures constructed on shorelines is likely to be variable, depending on factors such as prevailing current direction, wave strength, and underlying substrate as well as the physical properties of the structure itself.

Given that there are not expected to be material changes in the dimensions of the jetty following upgrade works; it is not expected that its proposed upgrade will alter, in a substantial way, the geomorphological processes and sediment drift that currently govern the shoreline ecosystem of the Buhuka Flats, additional to the sediment deposition and erosion either side of the jetty that has already taken place. However, there is potential for the construction of the new water intake and water extraction pump station to affect geomorphological processes and sediment drift downshore of Well Pad 2; which in combination with the existing jetty structure, could potentially affect the sediment drift or shoreline morphodynamics between Well Pad 2 and Bugoma Lagoon.

### ■ Water Quality

#### Sediment Loads

The construction of the jetty upgrade and water intake station has the potential to alter the water quality within the immediate surrounds of the construction activities through disturbance of the lake bed, and introduction of sediment into the water column during the works. It is expected that these increased sediment loads will dissipate reasonably quickly following completion of the jetty upgrade. The sediment loads in the vicinity of the jetty are not expected to exceed those that would normally be expected during windy periods on the lake and the consequent turbid conditions caused by those winds. Furthermore, these works are not expected to permanently alter the water chemistry in the vicinity of the jetty given the large buffering capacity of the lake compared to the scale of the works (see Surface water chapter, Vol 4a). Therefore, it can be expected that the upgrade of the jetty will not significantly affect the sediment loads and water quality of the near-shore habitats during construction.

The construction of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, new in-field roads, crusher plant/spoil area A and development of existing well pads and associated infrastructure could cause increased sedimentation of near-shore habitats on the Buhuka Flats. Sediment generated during construction of the CPF itself, and other onshore infrastructure, could enter the lake during storm flows over the three-year construction period, peaking during site establishment when vegetation is being cleared and civil earthworks are ongoing. The soils of the Buhuka Flats are dispersive, and active soil erosion is evident in the LSA. With the construction of the Project's components, and the consequent exposed areas of soil, there is a potential that, without adequate erosion and sediment control measures in place, sediment loads within the watercourses draining the Project footprint could increase. Cleared areas will be prone to sheet flow and scour, and high sediment loads could be expected, particularly in River 1, which will receive the drainage from the CPF earthworks and temporary camp (Figure 17). Additional sediment loads will also be contributed from the expansion of Well Pad 2 to its full size, and the construction of the permanent camp. These sediment-laden watercourses report to Lake Albert and the Bugoma Lagoon, and, hence, there is a potential for increased sediment loads in the near-shore environments. Near-shore habitats particularly at risk include the lagoon, and to a lesser extent, the shallow river-associated habitats. Nevertheless, the watercourses draining the CHAA support dense emergent vegetation (see Section 7.1.1). Such vegetation forms an impactful filter for most sediment (IECA 2008), therefore, it can be expected that sediment loads reporting to the near-shore habitats, at least via the Kamansinig River, River 1 and Masika River, could be minimal. Sediment loads from overland (stormwater) flows may not be retarded by vegetation, and hence may report to the near-shore





habitats, contributing to measurable increased turbidity during and after storms, where River 1 discharges to the Lake.




Figure 17: Rivers and catchments on Buhuka Flats

### Oil and Chemical Pollution

Accidental spillages of small quantities of fuels and chemicals, and rain wash from oily construction equipment working on the jetty and water intake station during construction could report to the near-shore habitats of the CHAA, via stormwater drainage into River 1, and subsequently Lake Albert, south west of Well Pad 2. There is a real potential for this to occur as part of the jetty upgrade, expansion of Well Pad 1, and construction of new Well Pads. Both the jetty and the proposed expansion of Well Pad 1 are located on the lake shore, with no meaningful buffer between the facilitates and the lake; in these areas, the impact of this contamination would be more obvious than on land, and would also be harder to contain and clean up.

A further risk could result from the construction and drilling of the wells. While control systems such as bunding are proposed to manage contaminated stormwater and wash-water from the well pads, the presence of drilling crews on site for nearly a year (From 1,000 up to 2,000 personnel (including 200 day workers) are expected be employed at peak times during the construction phase. Employment will be over a 34-month period), using potentially hazardous drilling fluid and other hazardous materials; and the absence of a buffer between the





well pads and the lake (or, in the case of Well Pad 1, the seasonal wetland); makes it likely that occasionally-contaminated drainage could reach the lake, unless there is a very high level of control of day-to-day activities.

These risks should be assessed in the context of the sensitivity of the near-shore environment to oil and chemical spills. Certain invertebrate species (for example, aquatic snails (Araujo et al. 2012), mayflies (Savić et al. 2011)) and juvenile fish (for example, Agamy 2013) are highly sensitive to chemical pollutants, particularly hydrocarbons. Currently, the concentrations of hydrocarbons and other pollutants in the lake water are below levels that could cause harm to the environment (Vol 4a Surface Water); this is supported by the healthy aquatic communities observed in the lake's near-shore habitats (see Section 6.2). Therefore, impacts arising from potential changes to the water quality are predicted to be beyond the expected range of natural disturbance perturbations.

As such, without adequate spill prevention and clean-up measures in place, the entrainment of chemical and oil pollution to Lake Albert could have a detrimental impact on the near-shore habitats of the CHAA through the introduction of toxic compounds and pollutants. Such a spill could also have a detrimental impact on the aquatic invertebrate community and juvenile fish occurring in these habitats; this includes the Critically Endangered Mud Snail (*Gabbiella candida*) (if it does occur in the area).

### *Discharge of treated sewerage*

The only continuous discharge into the lake during the construction phase will be from the sewage treatment plants. The effluent will be treated to meet the Ugandan effluent quality discharge standards (see Vol 4a Surface Water). At peak construction, the discharge will be around 300 m<sup>3</sup>/day at the EPC contractor plant, and 50 m<sup>3</sup>/day at the drilling camp plant. Both of these discharges will enter the lake via River 1, just south of Well Pad 2. The currents in the near-shore area of the LSA are not well known, so dispersion of nutrients in the sewage effluent have not been modelled. However, given the point source discharge, and the quantity of effluent involved, there is a risk of localised eutrophication, causing algal growth and possibly even fish kills around the discharge point. Impacts on water quality as a result of this potential eutrophication are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Ecosystem composition

All six of the main habitat types in Lake Albert, as identified for fishes by Wandera and Balirwa (2010) (that is, shallow river-associated waters, open sandy shores, lagoons, large bays, rocky escarpments, and, open-water habitats), occur within the near-shore areas of the CHAA (see Section 7.1.1, Figure 5). Similarly, the species guilds associated with the near-shore habitats of the CHAA, in particular fish, are well represented throughout those regions of the lake that have been investigated (for example, see Wandera and Balirwa 2010, Taabu-Munyaho et al. 2012). Consequently, at baseline, the composition of these ecosystems can be said to be in good condition and reflective of the aquatic diversity of Lake Albert. Similarly, these aquatic habitats have a well-developed structure, that is, well-defined aquatic plant layers associated with underwater features and substrates.

The construction of the Project is likely to result in localised alteration of the ecosystem composition of the aquatic communities; particularly the open sandy shoreline habitat in the vicinity of the proposed water intake and water extraction pump station, and the sewage effluent discharge outfall. Although the upgrade of the existing jetty is not expected to substantially alter that section of open sandy shoreline within which it is located; a degree of sediment deposition and erosion has already taken place on either side of the structure; therefore similar effects are anticipated up-current and down-current of the proposed water intake and water extraction pump station near Well Pad 2, which would be beyond the expected range of natural disturbance.

The point source effluent discharge, and the quantity of effluent involved, presents a risk of localised eutrophication, which could cause changes in algal growth rates, and aquatic vegetation community composition, thereby changing diversity and complexity of the aquatic habitats and their ability to support associated aquatic faunal communities. Impacts from the changes to ecosystem composition as a result of this eutrophication are predicted to be beyond the expected range of natural disturbance perturbations.

Although the open sandy shoreline habitat of itself is not very complex, consisting of a gently sloping lake bed comprised of fine and medium-grained sediments (Wandera and Balirwa 2010), with occasional aquatic plants;





it could support species that trigger critical habitat including the Critically Endangered mud snail (*G. candida*) and African Soft-shelled Turtle (*T. triunguis*); therefore, any negative impacts on this habitat have the potential to be of major significance, and residual impacts will likely require offsetting.

### ■ Ecosystem configuration

The construction of the CPF is expected to last for three years, and drilling will continue on individual well pads for a period of seven years. Although the upgrading of the jetty is not expected to substantially alter the configuration of the aquatic ecosystems and habitats in the CHAA, the construction of the new water intake and pump station could interrupt the connectivity amongst the near-shore aquatic habitats in its vicinity, to an extent beyond the range of natural perturbances.

### Impact Classification

The near-shore aquatic habitat's sensitivity is high because these habitats potentially support populations of the Critically Endangered Mud Snail (*G. candida*), the Vulnerable African Soft-shelled Turtle (*T. triunguis*), and the range-restricted and Near Threatened Snail (*Bellamyia rubicunda*). Near-shore aquatic habitat constitutes Tier 2 Critical Habitat for the Mud Snail (*G. candida*) (Table 4). Impacts on this habitat are therefore classified on the basis of high sensitivity to potential effects of the proposed development.

### ■ Representativeness

Impacts to the representativeness of the habitat will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA, and will only constitute approximately 0.2% loss of sandy shore habitat in the wider area. Impact duration will be into the far future (that is, ~25 years) because the jetty will remain in place even after the project is decommissioned. The changes to the representativeness of the near-shore habitats are expected to be reversible over time as naturalised geomorphological processes re-establish in the vicinity of the jetty and water intake station following construction, and re-establish following decommissioning. The magnitude of the impacts of construction/decommissioning on representativeness of the near-shore aquatic habitats is low; however, this must be weighed against the high sensitivity of these habitats. Therefore, the magnitude and sensitivity combine to produce an overall impact of moderate significance to representativeness prior to the application of project-specific mitigation measures (Table 7).

Following the application of appropriate mitigation measures, the impact significance is expected to remain the same, that is, moderate, primarily because the magnitude will still remain the same due to the loss of habitat, and the sensitivity of the receptor will remain the same.

### ■ Topography (geomorphology), sediment load, and water quality

Impacts to the topography (geomorphology), sediment, and water quality will be adverse. The geographical extent of impacts will be regional because impacts are restricted to the CHAA. Impact duration will be short-term (that is, ~2 years) because impacts are expected to be limited to the construction phase.

The magnitude of the impact of the upgrade of the jetty and construction of the water intake pump station on topography (geomorphology) and sediment transport is medium, as although there will be no material change to the jetty's current dimensions, the water intake pump station will be constructed at a new location; and to date, the jetty appears to have caused some observable impact on the movement of sediment along the lake shore (build-up of sediment on the eastern side, and erosion on the western side). The geographical extent of any impacts will be restricted to the LSA, and the duration of impacts will be short-term (that is, ~3 years) because impacts are expected to be limited to the construction/decommissioning phase. Therefore, the significance of impact of the jetty upgrade and the construction of the new water intake pump station is considered to be **major**. Following the application of appropriate mitigation measures, the impact significance on topography (geomorphology) and sediment transport is expected to be reduced to moderate, primarily because the magnitude will become low, and the sensitivity of the receptor will remain the same.

The magnitude of the impact of construction/decommissioning of the production facility and subsequent increased sediment load on water quality is medium, because, prior to any mitigation, there is potential for pollution to alter the quality of near-shore habitats. The geographical extent of impacts will be regional, being restricted to the CHAA, because, for example, possible effects on nursery areas for fish could eventually be





felt across the entire Lake. The duration will be short-term (that is, ~3 years), that is, limited to the construction/decommissioning phase. Therefore, the medium magnitude of the impact and high sensitivity of the near-shore habitats combine to produce a major overall impact level to water quality as a result of increased sediment load during the construction/decommissioning of the production facility, pre-mitigation (Table 7). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become low, and the sensitivity of the receptor will remain the same.

In the context of the absence of a meaningful buffer between the well pads and the lake, the magnitude of the impact of the potential entrainment of small quantities of oil, chemicals and potentially hazardous drilling fluid into stormwater and wash water from the well pads on water quality is medium. This is because prior to the application of site-specific mitigation measures, the potential for pollution to alter water quality and affect highly sensitive aquatic species is substantial. As was the case for sediment load, the geographical extent of impacts could be regional. The duration will be short-term (that is, ~3 years) i.e. limited to the construction phase. As mentioned previously, the near-shore aquatic habitat's sensitivity is high, therefore, the magnitude and sensitivity combine to produce a major overall impact level to water quality as a result of potential contamination with oil and potentially hazardous chemicals during the construction/decommissioning of the production facility and drilling of the wells, pre-mitigation (Table 7). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become very low.

The magnitude of the impact of construction/decommissioning of the production facility and subsequent discharge of treated sewage effluent prior to mitigation is medium; given the quantity of effluent involved, the extent of changes to ecosystem composition brought about by increased algal growth (above that normally occurring) and eutrophication in the vicinity of the point source discharge could be significant. Therefore, the medium magnitude and high sensitivity combine to produce a major overall impact level to ecosystem composition during the construction/decommissioning phase of the Project, pre-mitigation (Table 7). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

### ■ Ecosystem composition

Impacts to the ecosystem composition of near-shore habitats will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA, in the vicinity of the jetty and proposed water intake pump station. Impact duration will be short-term (that is, ~3 years) because impacts are expected to be limited to the construction/decommissioning phase, and impacts would be largely reversible following construction, and later following decommissioning. The magnitude of the impact on ecosystem composition is low because, prior to mitigation, the potential for localised changes to ecosystem composition potentially brought about by changes in sediment erosion/deposition patterns, spread of invasive species, and changes in aquatic invertebrate, fish and plant communities around the treated sewage discharge point is possible. Therefore, the low magnitude and high sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction/decommissioning phase of the Project, pre-mitigation (Table 7).

Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

### ■ Ecosystem configuration

Adverse construction/decommissioning impacts on ecosystem configuration are predicted to occur with the construction of the new water intake pump station. The magnitude of the resultant impacts to the ecosystem configuration of near-shore habitats is predicted to be low, as the geographical extent of impacts will be local - restricted to the LSA, in the vicinity proposed water intake pump station, although impact duration will be long-term because impacts are expected to be continue beyond the construction phase and into the operation phase.





Therefore, the low magnitude and high sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction/decommissioning phase of the Project, pre-mitigation (Table 7). Following the application of the recommended mitigation measures, the impact significance is expected to remain moderate, as impacts of a low magnitude will continue for the operational lifetime of the pump station, and the sensitivity of the receptor will remain the same.

**Table 7: Potential impacts in the construction phase to near-shore habitats**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low - 2	High - 4	Moderate – 8	Low - 2	High - 4	Moderate – 8
Topography (geomorphology) and sediment transport	Medium - 3	High - 4	Major – 12	Low - 2	High - 4	Moderate – 8
Water quality – sediment loads	Medium - 3	High - 4	Major – 12	Negligible - 1	High - 4	Minor – 4
Water quality –oil and chemical pollution	Medium - 3	High - 4	Major – 12	Negligible - 1	High - 4	Minor – 4
Water quality - sewerage	Medium - 3	High - 4	Major – 12	Negligible - 1	High - 4	Minor – 4
Ecosystem composition	Low - 2	High - 4	Moderate – 8	Negligible - 1	High - 4	Minor – 4
Ecosystem configuration	Low - 2	High - 4	Moderate – 8	Low - 2	High - 4	Moderate – 8

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios with the level of Lake Albert, which has, in the recent and not so recent past, varied quite dramatically (Talbot et al. 2006), it is conceivable that level of the lake may increase or decrease thereby altering near-shore habitats.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 7.1.1). The actual extent of possible habitat may have been over-estimated.

**10.1.1.2 What impact could the construction/decommissioning of the Project have on the wetlands in the Buhuka Flats region of the CHAA?**

**Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the wetlands of the CHAA were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

Loss of habitat due to direct disturbance and clearing associated with the Project (including existing, permitted infrastructure, and proposed infrastructure not yet built) was quantified by overlaying the current, baseline extent of the vegetation communities with the current and proposed Project footprint.







Additional, indirect impacts to habitat were estimated by applying a 0.50 km buffer to the Project infrastructure. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, changes in water quantity and quality (drivers of ecosystem processes and functions), and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur, and the disruption of known corridors. This was accomplished by examining available scientific literature about the ecology of wetlands (permanent and seasonally flooded grasslands).

**Impact Assessment**

■ Representativeness

In the Buhuka Flats, the CHAA supports approximately 151 ha of wetlands (see Table 6). These are comprised of permanent wetlands (83 ha, or 55% of wetlands in the Buhuka Flats) and seasonally flooded grasslands (69 ha, or 45%).

Wetlands in the LSA are associated with drainage off the escarpment, which is seasonal to varying degrees. The shallow gradients across the Buhuka Flats encourage wetland formation, and most of the stream channels are associated with fringing wetland vegetation and seasonally flooded grasslands. The Kamansinig River is hydrologically linked to the Bugoma Lagoon, the large papyrus-fringed wetland south-west of well pad 1.

Construction/decommissioning of the Project infrastructure will affect both the permanent wetlands and the seasonally flooded grasslands of the LSA. Figure 18 shows the main areas of direct impact. Table 8 quantifies the impact, based on the area of physical disturbance during construction and the expected long-term (or permanent) impact after construction and decommissioning. It should be noted that the infield access roads have already been licensed and built (see Table 8), but are included in this discussion of impacts for completeness. The construction right-of-way for both roads and flow lines is 20 m wide.

The road and flow line to well-pad 3 cut across the permanent wetlands of the lower Masika River; the in-field road and flow line from well-pad 3 to well-pad 2 cut across seasonally flooded grassland; the extension of the airstrip has resulted in the loss of approximately 1 ha of seasonally flooded grassland; and the extension of well pad 1 will result in the direct loss of an additional 4.8 ha of seasonally flooded grassland. This loss equates to approximately 4.7 ha (3.11%) of wetlands in the CHAA.

**Table 8: Wetland areas directly affected by the construction of the production facility**

Project Infrastructure Name	Wetland Area Affected (ha)*	Propoprtionate loss (%)
	Kamansinig River	
Infield Access Road**	1.5	0.99%
Infield Flowline	0.5	0.33%
Well Pad 1	2.7	1.78%
Total	4.7	3.11%

\* Blue cells show temporary construction impact. White cells show long term / permanent impact

\*\* Licensed and built






Figure 18: Wetlands in the Buhuka Flats directly impacted by construction/decommissioning of the Production Facility

The loss of this quantity of wetlands in relation to their representation in the wider CHAA and the Buhuka Flats is not insubstantial (~4% - 5%); furthermore, because this loss is brought about by the construction of linear features, there is a potential for the downstream wetland habitat to be affected if proper management controls are not implemented.

Impacts from the changes to representation of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Wetland Structure and Ecosystem Composition

#### Roads

The construction of linear infrastructure, such as roads through wetland communities is known to have detrimental impacts on the functioning, processes and species composition of these communities (Roise et al. 2004). Road construction may result in significant loss of biodiversity at both local and regional scales due to restricted movement between populations, increased mortality, habitat fragmentation and edge effects,





invasion by exotic species, or increased human access to wildlife habitats, all of which are expected to increase local extinction rates or decrease local recolonisation rates (Findlay and Bourdages 2000). Consequently, it could be expected that, particularly the construction of the airstrip and in-field roads through the seasonally flooded grassland would have already altered the ecosystem processes and functions driving these wetlands, especially downstream. This could have occurred if flow paths for water, both surface and sub-surface, were not maintained (see Vol 4a Surface Water). If these flow paths are not maintained on an ongoing basis, there is a potential that the wetland community downstream of the obstruction could become altered – typically via redirection of flows, and/or flow concentration and channel incision in downstream wetlands that were previously supplied by diffuse, dispersed flows. In an environment like the Buhuka Flats, where soils are clayey and dispersive, the risk of incised drainage and associated loss of wetland function due to concentration of water flows is high. The vegetation within these seasonally flooded grasslands is adapted to seasonal inundation, and, therefore, is dependent upon that cycle of wet and dry for survival. Additional, associated impacts that could occur in tandem with channel incision and subsequent wetland desiccation include exotic species invasion.

As part of standard construction methods, wetland/drainage line crossings have been installed as part of the in-field road and airstrip construction. Nevertheless, even with such measures in place, there is still a potential for changes to the wetland character to occur. For example, there is the potential for erosion downstream of the crossings, backwater upstream of the crossings, and erosion at the entrance to the crossing structures. The airstrip is one area in particular where construction across a drainage line might lead to decreased flows and erosion downstream of the airstrip. This could lead to changes in the ecosystem functions and processes in the downstream wetlands.

There is an existing example of this on the Kamansing River, where a road crossing upstream of the airfield leads to a quarry near the base of the escarpment (Figure 19). The damage was done by inappropriate culvert design (size, numbers and spacing), mostly in the wrong place, which has resulted in an altered flow regime, and subsequent desiccation of the wetland downstream of the impeding feature (the road), and increased wetness upstream of the road. The long-term impacts of the flow impediment created by the road include encroachment of terrestrial and exotic plant species to areas of the wetland that have become desiccated, and changes in vegetation community upstream – from seasonally flooded grassland communities to plants more characteristic of permanently saturated conditions, such as *Phragmites* sp. and *Typha* sp. During times of peak rainfall, outflow to the downstream areas is likely to be concentrated at a single spill point, and could contribute to channel erosion and further desiccation of the downstream wetland areas over time.

In addition, these wetland communities are already under pressure from livestock grazing, and harvesting of fibre for house construction. It is possible that these communities may change in the long-term as grazing pressure increases, and the human population of the Buhuka Flats increases. These changes could alter the habitat structure and composition, which, in turn, could affect the utilisation of these wetlands by the currently resident species guilds.

Impacts to ecosystem composition of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.



*Figure 19: Impact of a poorly-designed road crossing on wetland in Buhuka flats*

### Flowlines

Buried pipelines crossing rivers and wetlands will not (of themselves) obstruct surface water flow. The backfill into the trench is not normally cemented or compacted, which, for small diameter pipelines, minimises the risk of impact on subsurface flow. The risk to wetland function is mainly due to the disruption of wetland vegetation and soils by heavy machinery, particularly when tracked vehicles are used that have greater impact on soil structure and the soil profile is overturned due to careless construction management. Disruption of flows and ecosystem composition may also occur if fill material is imported into the wetland to provide stability for excavators and pipe layers, and is not completely removed and replaced with the natural soils after construction. This may result in long-term changes in vegetation composition and changes in flow patterns, with similar downstream consequences to those described under 'roads'.

Impacts on wetland fauna will depend on the changes in wetland vegetation as a result of construction; in addition, the noise and sensory disturbances created by the construction equipment could alter the behaviour of species frequenting the wetlands during construction/decommissioning. For example, wading birds and Grey Crowned Cranes are unlikely to tolerate construction nuisance within 500 m and could avoid these areas entirely during the construction period. If construction occurs during the breeding season of the Grey Crowned Crane (that is, the dry season (Archibald et al. 2013)), which could be breeding in the permanent wetland associated with the Masika River, then disturbance from the construction activities could cause nest abandonment (Strasser et al. 2013).

### Wellpad 1 expansion

The extension of well pad 1 will impact directly on wetland functioning in the seasonally-flooded grassland associated with the lower reaches of the Kamansinig River. The existing well pad is within the northern edge of these seasonally flooded grasslands. The expanded well pad will extend the impact on the wetland into the centre of the floodplain. The darker colour of the wetland in the satellite image in Figure 20 shows its position in relation to the well pad. The magnitude of this impact is considered to be high from legal and functional perspectives – the location is prohibited by Ugandan legislation (Uganda Wildlife Act, 2000), and contrary to the natural habitat conservation guidelines of IFC PS6 - and the impact on wetland function could be material, interfering with subsurface flow and surface flow during peak flow events. The vegetation within these seasonally flooded grasslands is adapted to seasonal inundation, and, therefore, is dependent upon that cycle of wet and dry for survival. The noise and sensory disturbances created by the construction equipment could alter the behaviour of species frequenting the wetlands, particularly Grey Crowned Crane. Coupled with the



very high sensitivity of this system, with its hydrological interconnectivity to the Bugoma lagoon, the impact significance will be high.

### Increased runoff from construction areas

The majority of the area affected by construction of the CPF will drain north into River 1 (Figure 20). Perimeter drains upslope of the CPF will divert clean stormwater flow around the platform and discharge directly to River 1. The first 15 minutes of stormwater draining from potentially contaminated areas such as the platforms will be diverted to a testing and treatment tank, before being discharged to River 1 once acceptable standards have been reached. After the first 15 minutes, stormwater from potentially contaminated areas will be discharged directly to River 1 via the perimeter drains. Stormwater from the construction of most of the permanent camp, the extension of well pad 1 and the southern section of the flow line linking well pad 1 to the CPF will drain southward into the Kamansing River, and will follow the same standard clean and dirty water practises. Figure 20 shows the most likely direction of flow from the construction sites.

Drainage volumes during storm events are expected to be larger (due to the removal of vegetation and the compaction of ground surfaces, and hardstand areas), and, consequently, peak-flow volumes will be larger. Additionally, concentration of stormwater flows into the drainage lines will significantly increase the risk of channel incision, which in the flat environment between the escarpment and the lake, may result in drying out of the associated wetlands due to more rapid drainage of the area. The magnitude of this impact will be exacerbated by the soils, which are dispersive (see Vol 4a Soils and Landuse), prone to gulley erosion, and, therefore, highly likely to form incised channels.



# BIODIVERSITY IMPACT ASSESSMENT




Figure 20: Direction of storm flows from construction sites





### ■ Water Quality and Ecosystem Composition

Impacts to ecosystem composition of the wetlands in the CHAA are predicted to be well beyond the expected range of natural disturbance perturbations; factors affecting wetland water quality that could impact ecosystem composition, including erosion and sedimentation, fuel and chemical spills, discharge of hydrotest water, disposal of treated effluent, overturning of acid sulphate soils and population increases are discussed below.

#### **Erosion and Sedimentation**

The construction of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, new in-field roads, crusher plant/spoil area A, and associated infrastructure could cause increased erosion and sediment-laden run-off to report to the wetlands surrounding the Project footprint. The soils of the Buhuka Flats are dispersive, and active soil erosion is evident in the LSA. With the construction of the Project's components, and the consequent exposed areas of soil, there is a potential that, without adequate erosion and sediment control measures in place, sediment loads within the watercourses draining the Project footprint could increase. Hence, there is a potential for increased sediment loads in the wetland habitats. The watercourses and associated wetlands draining the LSA support dense emergent vegetation (see Section 6.1.1.1). Such vegetation can form an impactful filter for most sediment (IECA 2008); therefore, it can be expected that sediment loads reporting to downstream wetland habitats could be minimal. Nevertheless, if sediment loads are substantial, there is a potential for that sediment to smother wetland vegetation and interfere with aquatic invertebrates. If this occurs, it could detrimentally affect the wetland processes and functions, which, in turn, could alter wetland composition, albeit on a localised scale.

As part of standard construction methods, appropriate drainage line crossings will be installed as part of the in-field road and airstrip construction. Nevertheless, even with such measures in place, there is still a potential for changes to the wetland character to occur. For example, there is the potential for erosion downstream of the crossings, backwater upstream of the crossings, and erosion at the entrance to the crossing structures. The airstrip is one area in particular where construction across a drainage line might lead to decreased flows and erosion downstream of the airstrip. This could lead to changes in the ecosystem functions and processes in the downstream wetlands.

#### **Hydrocarbon and Chemical Spills**

Accidental spillage of fuels and chemicals during the construction/decommissioning of the CPF are likely, where the most complex construction activities will take place over a period of 3 years, involving a wide range of potentially toxic materials. On the well pads, the presence of drilling crews for around a year, using large quantities of potentially hazardous drilling fluid and other hazardous materials, increases the risk of escape of contaminated water into watercourses and subsequently the wetlands of the LSA. The highest risk is most likely at Well Pad 1, most of which is being extended into the seasonally-flooded grassland wetland associated with the Kamansing River.

Certain invertebrate species (for example, aquatic snails (Araujo et al. 2012), mayflies (Savić et al. 2011)) and juvenile fish (for example, Agamy 2013) are highly sensitive to chemical pollutants, particularly hydrocarbons. Currently, the concentrations of hydrocarbons and other pollutants in the water of the Bugoma Lagoon are below levels that could cause harm to the environment; this is supported by the healthy aquatic communities observed in the lagoon's and Masika River wetland habitats (see Section 6.2). The proximity of the main areas of construction to wetlands on the Flats provides little buffer in the event that minor day-to-day spillages escape from the work areas, which increases the risk of impact.

#### **Discharge of Hydrotest Water**

The commissioning of pressure vessels and flow lines involves hydrotesting, in which the vessels are filled with water and pressurised to verify their integrity. On occasions, biocides and corrosion inhibitors are added to the water, depending on the residence time, before it is discharged. After hydrotesting, the water will be treated and tested to confirm it complies with the Ugandan effluent standards and World Bank Effluent Standards, and disposed back into Lake Albert. Treatment typically involves sand filtration on sand bed screens to remove scale and other solids. It is not expected that biocides and corrosion inhibitors will be added to the hydrotest water in the flowlines but, in case that any additives are used, the hydro test water will be





subjected to a neutralizing treatment process typically by neutralizer dosing and bio-inoculation at the end of the hydro test exercise. The treated water will then be tested to demonstrate that the additives have fully been neutralized to acceptable prior to discharge into the environment. The solids will be filtered and disposed to a certified waste disposal site.

### Disposal of Treated Sewage Effluent

Treated sewage effluent in excess of 300 m<sup>3</sup>/day will be discharged for much of the construction phase. The effluent will be treated to meet the Ugandan effluent quality discharge standard (see Vol 4a Surface Water). It is proposed that the effluent from the temporary camp will be discharged into River 1, north of the camp. The effluent from the drilling camp will be discharged into River 1, within Buhuka flats.

Wetlands are efficient nutrient sinks and have been used in both controlled and uncontrolled conditions to polish sewage effluent. Depending on the point of discharge, River 1 is likely to be tolerant of the additional daily flow and the addition of nutrients, which will promote the growth of emergent wetland vegetation.

### Overturning of Acid Sulphate Soils

The potential for the occurrence of acid sulphate soils in the permanent wetland associated with the Masika River should not be discounted. Typically, submerged sediments in inland aquatic ecosystems have very little oxygen below the first few millimetres, and can, therefore, be sites where sulphur compounds are formed, which can lead to potential acid sulphate soils (EPHC 2011). When these soils are disturbed and exposed to oxygen, such as through the construction of a trench through a wetland, the sulphur compounds react with the oxygen in the air and the water to produce sulphuric acid. If the amount of acidity produced by this process is greater than the system's ability to absorb, or neutralise that acidity, then the acidity, or pH, of the system falls (EPHC 2011). The formation and release of this acid can lead to adverse impacts to aquatic habitats. Assuming the wetland does not have a high capacity to neutralise the acid, the process of forming the acid can consume oxygen, and, in extreme cases, can remove all of the oxygen from the water column, resulting in the death of aquatic organisms. The acid formed by this process may also lead to the mobilisation and availability of heavy metals (such as, cadmium and lead), aluminium, iron, and arsenic into the environment. These elements can be toxic to aquatic life (EPHC 2011).

Within Uganda, these soils are typically associated with papyrus peats and other permanent wetlands. The permanent wetland associated with the lower reaches of the Masika River, has a potential to harbor potential acid sulphate soils. Therefore, if these soils are disturbed, for example, during the construction of trenches for flowlines, there is a potential for acid formation and the consequent toxic impacts to the ecosystem. If severe enough, such toxic impacts can cause the permanent dieback of wetland vegetation and alter wetland functioning and processes (EPHC 2011).

However, while acid sulphate soils may be present in the permanent wetlands along the Masika River to the south and in the Buhuka Lagoon, they are not expected in the seasonal wetlands that will be affected by construction of the production facility and the toxic effect of acid generation associated with their disturbance should not arise.

### Increase in Population on Buhuka Flats

Indirect impacts on wetlands as a result of water quality impacts could occur as a result of the migration of people onto the Buhuka Flats in search of work. Existing sanitary conditions on the Flats are poor, with all of the streams that drain across the Flats being contaminated with faecal waste from both animals and humans (Vol 4c Social Impacts). Increasing population pressures on the flats will exacerbate these conditions. Increased grazing pressure and erosion from denuded areas around expanding settlements will increase erosion and sedimentation in the wetlands.

#### ■ Ecosystem configuration

Roads are known to be significant barriers to, or can alter behaviours of, a range of wetland wildlife, from: amphibians (for example, Pontoppidan et al. 2013); to turtles (for example, Langen et al. 2012). Depending on the species, the presence of the existing in-field roads may affect individuals in many direct and indirect ways. For example, roads may inhibit seasonal migration and may cause an impactful loss of habitat due to







avoidance. The presence of the in-field roads through the wetlands of the Buhuka Flats are not expected to be major barriers to movement for those species inhabiting them. These roads are only expected to be 5 m wide and unsealed, and, once construction is complete, they will convey limited traffic volumes. Therefore, they are not expected to be major barriers.

The construction of the existing and proposed in-field roads and airstrip has/will cut currently contiguous wetlands in the LSA. If not managed correctly during the construction process, that is, the installation of appropriate drainage connections, these Project components could cause permanent barriers between the two newly separated wetland habitats – as has already been demonstrated at the road crossing adjacent to the airstrip (Figure 19). The process of clearing the wetlands for the construction of the roads and airstrip will create edge impacts, and result in the fragmentation of the wetland habitats. As already discussed, vegetation clearing creates edges or boundaries where habitat meets a disturbance. These edges open up habitat in areas where it was previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges also often result in changes in species composition along the edge, with the edges typically becoming dominated by pioneer and weedy species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Fragmentation of wetland habitat is known to reduce the viability of many species and the wetland as a whole, with the viability of the particular fragment dependent on its size, proximity and, hence, connectivity to other wetland habitats (Uzarski et al. 2009).

What long-term impacts the construction of these roads and the airstrip could have on the wetland communities' configuration are unknown; in particular their resilience. What is known is that these wetland communities are already under pressure from livestock grazing, and harvesting of fibre for house construction.

Impacts to ecosystem configuration of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The wetland's sensitivity is high because these habitats, particularly the permanent wetlands, are potential breeding habitat for Grey Crowned Cranes (a species of concern). The wetlands are also already under stress from livestock grazing and harvesting of fibre. This high sensitivity is weighed against the magnitude of each of the impacted indicators as described in the paragraphs below, in order to derive the overall impact level for each indicator.

#### ■ Representativeness

Impacts to the representativeness of the habitat will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA on the Buhuka Flats, and will constitute loss of approximately 4.2% of seasonally flooded grassland communities in relation to the CHAA. The flowline construction impacts will be generally short-term (subject to appropriate construction management); and while permanent wetland habitat loss will be caused by well pad 1 and its proposed expansion, and by the access roads, the relatively small area covered results in an overall impact of low magnitude. In the context of high wetland sensitivity, this results in impacts of moderate significance.

Following the application of appropriate mitigation measures, the impact significance is expected to remain minor, as some permanent wetland loss to the roads and expansion of well pad 1 will remain in place even after the project is decommissioned. The changes to the representativeness of the wetland communities, although probably irreversible, are expected to be amendable via offsetting (see Section 13.0).

#### ■ Wetland Structure and Ecosystem Composition

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because impacts are expected last as long as the in-field roads and airstrip are in place. The magnitude of the impact on ecosystem composition is medium because, prior to the implementation of the recommended site-specific mitigation, the potential for changes to wetland structure and ecosystem composition potentially brought about by the roads, flowlines, well pad 1 extension, potentially contaminated runoff from construction areas and population/livestock head increases are likely, which could result in edge effects, changed flow





regimes, and erosion and sedimentation of affected wetlands. Therefore, the magnitude and high sensitivity of the wetlands combine to produce a major overall impact level to ecosystem composition during construction/decommissioning of the Project, pre-mitigation (Table 9).

Following the application of appropriate mitigation measures, including appropriate construction management methods, the impact significance is expected to remain moderate, because although the magnitude will become low, the sensitivity of the receptor will remain the same.

### ■ Water Quality and Ecosystem Composition

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be for the duration of construction, which is expected to take approximately 3 years. The magnitude of the different project-specific impacts on ecosystem composition varies, as discussed under each of the subheadings below.

#### **Erosion and Sedimentation**

Clearing of vegetation will result in increased sediment loads in stormwater flows. These impacts will be of short duration, most likely occurring as a result of significant rain events, and will be most frequent in the early stages of construction when bulk earthworks are in progress and large areas of exposed earth are available. Impacts will be local (mainly in the catchment of River 1 and, to a lesser degree, in the Kamansinig River catchment). They will generally be reversible, and, consequently, have a medium magnitude. Wetland plant species are generally tolerant of occasional increases in sediment load in stormwater flows, and can serve as an effective sediment filter. In cases of severe and/or ongoing sediment loading, detrimental impacts on wetland vegetation and macro-invertebrates could be likely, which could detrimentally affect wetland processes and functions and, in turn, wetland composition at a localised scale. Overall wetland sensitivity to sediment increase is considered to be high; therefore the overall significance of this impact is moderate.

Following the application of appropriate mitigation measures, the impact significance is expected to reduce to minor, because the majority of sediment would not reach the rivers, lagoon or lake in one flush.

#### **Hydrocarbon and Chemical Spills**

The concentrations of hydrocarbons and other industrial pollutants in the wetlands of the LSA are presently below levels that cause harm in the aquatic environment. Some invertebrate species (such as aquatic snails, described in Araujo et al. 2012), mayflies (Savić et al. 2011) and juvenile fish (Agamy 2013) are particularly sensitive to these pollutants. The proximity of the main areas of construction to wetlands on the flats provides little buffer in the event that spillages escape from the work areas, which increases both the risk of impact, and the potential magnitude of such an event. In the absence of daily monitoring and management of site activities by competent personnel, the potential impact magnitude could be high, resulting in an overall impact of major significance.

Following the application of appropriate mitigation measures, the impact magnitude is expected to reduce to medium, which in combination with the high sensitivity of the wetland habitat, results in an overall impact of moderate significance.

#### **Discharge of Hydrotest Water**

Details on the use of biocides and corrosion inhibitors in the hydrotest water are not presently available, but for the purposes of the assessment it is assumed that they are present, and the hydrotest water will be discharged into the nearest drainage lines. Without management, this one-off release can present a severe risk to the aquatic environment, resulting in possible mortality and/or degradation of downstream species and ecosystems. Unmitigated impacts will be long term, only partly reversible, local in geographic extent, and, therefore, of low magnitude. Combined with high receptor sensitivity, this will result in impacts of major significance, prior to mitigation.

Following the application of appropriate mitigation measures, the impact magnitude is expected to reduce to low, which in combination with the high sensitivity of the wetland habitat, results in an overall impact of moderate significance.





### Disposal of Treated Sewage Effluent

Assuming that the point of discharge is located further upstream where the permanent wetland vegetation is established in that area and has the capacity to act as a buffer, River 1 is likely to be tolerant of the additional daily flow and the addition of nutrients, a change in species composition due to the promotion of the growth of emergent wetland vegetation could occur, with some species that adapt well to nutrient-enrichment (persistent emergent plants such as sedges (eg. *Scirpus sp.*), rushes (*Juncus sp.*), common reed (*Phragmites sp.*) and cattails (*Typha sp.*) could proliferate. The magnitude of this impact is considered low, which, combined with high receptor sensitivity, will result in impacts of moderate significance.

### Overturning of Acid Sulphate Soils

Acid sulphate soils are not expected in the seasonally-flooded grassland wetlands that will be affected by construction of the production facility; therefore, the toxic effect of acid generation associated with their disturbance should not arise.

The permanent wetland associated with the lower reaches of the Masika River, has a potential to harbor potential acid sulphate soils. Therefore, if these soils are disturbed during the construction of the flowlines between the well pads, there is a potential for acid formation and the consequent toxic effects to the ecosystem.

It is assumed that the potential acid sulphate soils in the permanent wetlands in the CHAA will be adequately managed during construction, and no lasting effects will occur. The magnitude of this impact is thus considered negligible, which, combined with high receptor sensitivity, will result in impacts of minor significance.

### Increase in Population on Buhuka Flats

In the absence of project interventions, the impact of increased population density (and associated increases in grazing livestock) on the Buhuka Flats is expected to be long-term, and to have a material effect on water quality in the wetlands across the Flats through reductions in sanitary water quality, exacerbation of wetland erosion, increased harvest of plant species used for traditional home construction, increased fire frequency and increased grazing pressure. The effects will be irreversible, and of high magnitude. In the context of the high wetland sensitivity, the predicted impact significance is major, prior to mitigation.

Following the application of appropriate mitigation measures, including appropriate construction management methods, the impact significance is expected to remain moderate, as although the magnitude will become low, the sensitivity of the receptor will remain the same.

### ■ Ecosystem Configuration

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will extend into the far future (that is, ~25 years) because impacts are expected to continue as long as the in-field roads and airstrip are in place. The magnitude of the impact on ecosystem configuration is medium because, prior to the application of the site-specific mitigation measures (Section 9.0), the potential for changes to ecosystem configuration is possible, especially from fragmentation.

As mentioned, the wetland's sensitivity is high because these habitats, particularly the permanent wetlands, are potential breeding habitat for Grey Crowned Cranes. The wetlands are also already under stress from increased fire frequency, livestock grazing and harvesting of fibre. Therefore, the magnitude and sensitivity combine to produce a major overall impact level to ecosystem configuration during the construction phase of the Project, pre-mitigation (Table 11).

Following the application of site-specific mitigation measures (Section 12.2), the impact significance is expected to remain moderate, although the magnitude will become low, yet the sensitivity of the receptor will remain the same.



**Impact Significance Rating**

**Table 9: Potential impacts in the construction phase to the wetlands of the CHAA**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Medium – 3	High – 4	Major – 12	Low – 2	High – 4	Moderate – 8
Ecosystem composition – wetland structure	Medium – 3	High – 4	Major – 12	Low – 2	High – 4	Moderate – 8
Ecosystem composition – water quality						
Erosion and sedimentation	Medium – 3	High – 4	Major – 12	Low - 2	High – 4	Moderate - 6
Hydrocarbon and chemical spills	High – 4	High – 4	Major – 16	Low - 2	High – 4	Moderate - 8
Discharge of hydrotest water	Medium – 3	High – 4	Major – 12	Low - 2	High – 4	Moderate - 8
Disposal of treated sewage effluent	Low - 2	High – 4	Moderate – 8	Negligible - 1	High – 4	Minor – 4
Overturning of acid sulphate soils	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Population increase	High – 4	High – 4	Major – 16	Low – 2	High – 4	Moderate – 8
Ecosystem configuration	Medium – 3	High – 4	Major – 12	Low – 2	High – 4	Moderate – 8

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the wetlands of the Buhuka Flats region of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.1). The actual extent of possible habitat may have been over-estimated.

**10.1.1.3 What impact could the construction/decommissioning of the Escarpment Road have on the escarpment vegetation corridors?**

While the escarpment road has been licensed and built on the basis of an earlier impact assessment (AWE, 2014c), the loss of escarpment habitat is considered and included here for completeness, and because more accurate post-construction information is now available about the area disturbed.

**Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the vegetation corridors on the escarpment were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.





Loss of habitat due to direct disturbance and clearing associated with the Project was quantified by overlaying the current, baseline extent of the vegetation communities with the Project footprint. This included the escarpment road (which, it is recognised, is covered by a separate ESIA process, see AWE (2014c); however, the inclusion of this road in this assessment is important because of its cumulative impacts on the vegetation of the escarpment) (ref. Chapter 17 Cumulative Impacts).

Additional, indirect affects to habitat were estimated by applying a 0.50 km buffer to the temporary camp and quarry at the top of the escarpment, and the footprint of the escarpment road. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, barriers to movement, changes in water quantity and quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur, and the disruption of known movement corridors. This was accomplished by examining available literature about the ecology of the vegetation communities on the escarpment, and scientific literature about the impacts of human activities on corridors, and consultation with experts.

**Impact Assessment**

■ Representativeness

The CHAA supports approximately 2443 ha of escarpment corridor vegetation communities (see Section 6.3.1.1.2), which are bounded on the east by agriculturally modified landscapes, and the Buhuka Flats on the west (Figure 11). As mentioned, these vegetation communities form part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2003) (see Section 7.1.2). Therefore, their continuous extent represents an important landscape feature in the CHAA.

The escarpment road traverses for approximately 5.5 km through the escarpment vegetation communities before it arrives on the Buhuka Flats. The total area of escarpment habitat that has been permanently lost as a result of the road’s construction is 12.8 ha; this comprises: 1.9 ha of open wooded bushland; 4.6 ha of open wooded grassland; 2.9 ha of dense bushland; 3.1 ha of dense wooded grassland; 0.1 ha of riverine woodland; and 0.1 ha of open grassland (Table 10). A further 3.6 ha has been temporarily disturbed along the edges of the road. Approximately 4.5 ha of open wooded bushland was lost to the temporary camp on top of the escarpment, this area has since been rehabilitated. All these vegetation communities are widely represented on the escarpment, and the CHAA.

The total loss within the CHAA equates to: 1% of open wooded bushland; 0.6% of open wooded grassland; 0.4% of dense bushland; 0.2% of dense wooded grassland; 0.1% of riverine woodland; and 0.02% of open grassland.

**Table 10: Permanent loss of escarpment vegetation due to construction of escarpment road**

Vegetation Type	Total area in the CHAA (ha)	Total area in Escarpment corridor (ha)	Area lost to Road (ha)	% loss in CHAA	% loss in escarpment corridor
Cultivation and settlement	31860.9	0.1	0.1	0.0%	100.0%
Dense bushland	1097.6	337.7	2.9	0.3%	0.9%
Dense wooded grassland	613.2	589.0	3.1	0.5%	0.5%
Open grassland	568.5	31.5	0.1	0.0%	0.3%
Open wooded bushland	523.0	214.0	1.9	0.4%	0.9%
Open wooded grassland	1900.9	552.2	4.6	0.2%	0.8%





Vegetation Type	Total area in the CHAA (ha)	Total area in Escarpment corridor (ha)	Area lost to Road (ha)	% loss in CHAA	% loss in escarpment corridor
Riverine woodland	74.8	69.8	0.1	0.2%	0.2%

The loss of this quantity of vegetation in relation to the total amount in the CHAA is not substantial. Nevertheless, the loss of this vegetation does open up and cut a previously contiguous tract of vegetation with a linear corridor that introduces edge effects and the concomitant aspects associated with those, as discussed below.

Impacts from the changes to representativeness are predicted to be beyond the expected range of natural disturbance perturbations.

■ Ecosystem composition

The construction of the road is expected to have contributed to losses of biodiversity at both local and regional scales due to it restricting movement between populations, increasing mortality, habitat fragmentation and edge effects, facilitating invasion by exotic species, or increased human access to wildlife habitats, all of which are expected to increase local extinction rates or decrease local recolonisation rates (Findlay and Bourdages 2000). Indeed, the construction of any linear corridor (such as the road) through an area of relatively intact vegetation, like that on the escarpment, creates edge effects that could, in the long-term, alter the composition of the ecosystem through which the road traverses. Vegetation clearing creates edges or boundaries where habitat (for example, riverine woodland) meets a disturbance (for example, the road). Edges associated with disturbance are different than transition areas, or ecotones, amongst vegetation communities, because disturbance edges tend to be abrupt with a high degree of contrast between two areas (for example, road and open wooded grassland). Edges open up habitat in areas where it was previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges often result in changes in species composition along the edge, with the edges typically becoming dominated by pioneer species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Often, these microclimates are favoured by weed species. However, certainly for fauna species, and depending on the species, edges can have either a positive or negative impact on habitat quality and quantity (Prevedello et al. 2013, Wellicome et al. 2014). Given the length of the escarpment road (approximately 5.8 km) traversing the escarpment vegetation communities, this equates to approximately 11.6 km of edges in the escarpment vegetation corridor, which would have otherwise not existed.

The escarpment road is sealed for its length down the escarpment. Sealing of the road presents other aspects that may affect the ecosystem composition of the communities on either side of the road. For example, concentrating water run-off from the sealed surface, which could carry contaminants, such as fuel, heavy metals and polycyclic aromatic hydrocarbons, to watercourses not otherwise influenced by such run-off. This could possibly lead to changes in water quality of local watercourses and erosion regimes.

The construction of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, pipeline, new in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure will not directly lead to changes in the composition of vegetation communities on the escarpment; however, indirect impacts from the changes to ecosystem composition are predicted to be well beyond the expected range of natural disturbance perturbations.

■ Ecosystem configuration

Roads, and especially sealed roads are known to be significant barriers, or alter behaviours, of a range of wildlife, from: small ground-dwelling mammals, insects, reptiles and amphibians (for example, Brehme et al. 2013, Pontoppidan et al. 2013, Rotholz and Mandelik 2013); to bats (for example, Berthinussen and Altringham 2012); to birds (for example, Kociolek et al. 2011); to primates (for example, Mammides et al. 2009); to large ungulates (for example, Leblond et al. 2013, Meisingset et al. 2013). Depending on the species, the presence





of roads may affect individuals in many direct and indirect ways. For example, roads may inhibit seasonal migration and may cause an impactful loss of habitat due to avoidance.

The construction of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, pipeline, new in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure will not directly lead to changes in the configuration of vegetation communities on the escarpment. The construction of the escarpment road is also unlikely to have had any direct impact on the configuration of vegetation communities on the escarpment. As in the case of ecosystem composition, the most significant changes in ecosystem configuration on the escarpment are already occurring as a result of increased migration into the area facilitated by the new ease of access, and associated degradation of escarpment vegetation, creating barriers to species movement. Ecosystem configuration impacts being caused by ongoing habitat degradation due to increased population and road traffic during operation are discussed in Section 10.2.1.

Direct impacts from the changes to ecosystem configuration on the escarpment as a result of the road are beyond the expected range of natural disturbance perturbations.

**Impact Classification**

The vegetation community of the escarpment’s sensitivity is medium because these habitats are already under stress from livestock grazing and harvesting of fuel wood and non-timber forest products. They also form part of a wider wildlife corridor, which is recognised for its regional importance.

Impacts to the representativeness of the habitat will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA, and will only constitute approximately 0.02 % loss of vegetation communities in relation to the CHAA. Impact duration will be into the far future (that is, ~25 years) because the road down the escarpment will remain in place even after the project is decommissioned. The changes to the representativeness of the vegetation communities, although possibly irreversible, are expected to be amendable via offsetting (see Section 13.0). The magnitude of the impacts of the road on representativeness of the vegetation communities of the escarpment is medium. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to representativeness during the construction phase of the Project, pre-mitigation (Table 11). Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as the magnitude will become low, while the sensitivity of the receptor will remain the same.

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because impacts are expected last as long as the road is in place. The magnitude of the impact on ecosystem composition is medium because, prior to any mitigation, the potential for changes to ecosystem composition potentially brought about by edge impacts is possible. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction phase of the Project, pre-mitigation (Table 11). Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as although the magnitude will become low, the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be regional because impacts are restricted to the CHAA. Impact duration will extend into the far future (that is, ~25 years) because impacts are expected to continue as long as the road is in place. The magnitude of the impact on ecosystem configuration is medium because, prior to mitigation, the potential for changes to ecosystem configuration is possible, especially inference with wildlife movement corridors. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction phase of the Project, pre-mitigation (Table 11).

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as although the magnitude will become low, the sensitivity of the receptor will remain the same.

**Table 11: Potential impacts in the construction phase to the vegetation communities of the escarpment**

	Pre-mitigation	Post-mitigation (pre-offsets)
--	----------------	-------------------------------





Indicator of potential impact	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6
Ecosystem composition	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6
Ecosystem configuration	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the vegetation communities of the escarpment, and the wildlife corridors of which they form part. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain. Indications are that they will increase in importance (Ayebare et al. 2013), provided human pressures do not overwhelm them.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 7.1.2). The actual extent of possible habitat may have been over-estimated.

**10.1.1.4 What impact could the construction/decommissioning of the Project have on the Bugoma Central Forest Reserve?**

**Impact Indicators**

The Ugandan Government will be responsible for upgrading the roads that the oil industry will need for access. Scheduled ‘oil industry’ road upgrades to support the Kingfisher Field Development Area include those shown in Figure 21. Although it is recognised that CNOOC will not be upgrading the roads, and the Ugandan Government will be responsible for the upgrades; any effects associated with, and stemming from the proposed road upgrades can be seen to be induced impacts arising as a consequence of the Project’s development.







## BIODIVERSITY IMPACT ASSESSMENT




Figure 21: Proposed Government road upgrades in the region

A formal impact classification based on indicators was developed for induced and cumulative impacts to the Bugoma Central Forest Reserve; the impacts are discussed, and their significance assessed through a **reasoned narrative**. An overall impact significance classification is then developed. This was accomplished by examining available literature about the ecology of the Bugoma Central Forest Reserve (BCFR), and scientific literature regarding the effects of migration and human population pressure on forests in Africa.

The impacts were assessed in light of the guidance provided by IFC (2013), and in consideration of other known projects being developed in the wider area. In particular, the development of the oil processing facility at Kabaale, the oil developments around the Kaiso-Tonya area (AECOM 2012), the Hoima-Mputa-Fort Portal-Nkenda power line, and the potential for regional population increases in the wider area.

### Impact Assessment

The Bugoma Central Forest Reserve (Bugoma CFR) is identified as a valued component for this impact assessment, certainly in terms of biodiversity (see Section 6.3.1.1.4). As identified in that section, apart from being one of the last stands of tropical semi-deciduous forests in the region, it also supports known populations of the Endangered Eastern Chimpanzee and range-restricted Nahan's Francolin (Plumptre et al. 2011), potential non-breeding habitat for the Endangered Madagascar Pond Heron (see Section 6.3.3.1), as well as elephants and a host of other threatened and irreplaceable species.

The R5 passes through the centre of Bugoma CFR in a north-south direction for approximately 9.7 km, and the P1 road runs along part of the main south-western and south-eastern boundaries of the Bugoma CFR, passing through the reserve in an east-west direction for approximately 3.5 km near Kisaru. Both roads are currently unsealed and relatively narrow, and become impassable from time to time in the wet season. No data traffic data are available for the roads.



The construction of the roads will cause the direct loss of forest species in the area of the road widening. Depending on the final road width (assuming a width of 10 m), the construction methodologies used and the need to accommodate traffic, this could result in the permanent removal of around 9.7 ha of forest habitat along the R5 section of road, and 3.5 ha along the P1 section of road from Kiziranfumbi and Nsozi.

The Bugoma Central Forest Reserve is home to populations of threatened species and an array of other species (see Plumptre et al. 2010, 2011). Many of these species will move within the forest and between sections of the forest. Although the negative effects of roads on wildlife in tropical rainforests, like Bugoma Central Forest Reserve, are poorly understood, indications are that: (1) many species avoid roads altogether (especially, medium-sized mammals, diurnal, solitary and group living animals, and ungulates); and (2) high vegetation cover on the road verges (Figure 22) increases crossing probability substantially (van der Hoeven et al. 2010). Currently, the road side vegetation on the R5 and P1 roads would encourage wildlife to cross (Figure 22). This could place them in the direct paths of traffic.

During construction, Project-generated traffic will consist of 65 trucks per day over a 2.5 year period, amounting to approximately one truck every 10 minutes during daylight hours. This, combined with a general increase in vehicular traffic to the area facilitated by the improving road surfaces, is predicted to cause increased disturbance to faunal species within the Bugoma CFR, as well as increase the risk of direct mortality of wildlife due to traffic collisions. Impacts to species of concern associated with Bugoma CFR (Eastern Chimpanzee and Nahan's Francolin) due to increased risk of traffic collisions during the construction phase are addressed in Section 9.1.2.



*Figure 22: The P1 road through Bugoma Central Forest Reserve*

### **Impact Classification**

Impacts from the upgrade of the R5 and the P1 roads and the resultant increased traffic (reducing adjacent habitat integrity as a result of noise, vibration etc) along that road during the construction of the Project will be adverse. The geographical extent of impacts will be regional because effects are restricted to the R5 and the P1 road corridors in the CHAA. Impact duration will be short-term (that is, limited to the road upgrade



construction, and the construction phase of the Project, that is, ~2 years). The magnitude of the effects of construction, in the context of habitat loss, on the Bugoma Central Forest Reserve is low.

The sensitivity of the Bugoma Central Forest Reserve is high because it is a threatened ecosystem that is already under pressure. Therefore, the intensity and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 12).

The mitigation hierarchy is an important process that has been used to minimise impacts to the Bugoma Forest Reserve. The focus for the continued use of the mitigation hierarchy during the road upgrade and construction will be continued development and implementation of mitigation measures through monitoring and adaptive management; including suggested measures for the Ugandan Government to apply in the management of the upgrade of the P1 Road (see Section 11.0).

Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the intensity could become negligible, although the sensitivity of the receptor will remain the same (Table 12).

**Table 12: Potential impacts in the construction phase to the Bugoma Central Forest Reserve**

	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	significance
Habitat and ecosystem integrity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4

As noted above, the focus for reducing impacts to the Bugoma Central Forest Reserve is mitigation to lessen various types of disturbance that may occur (Section 11.0). Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for residual impacts to the forest have been identified and are discussed in Section 10.0.

**10.1.2 What impact could the construction/decommissioning of the Project have on species of concern?**

This section presents the assessment of the impacts that the Project construction/decommissioning could have only on those species of concern that potentially trigger critical habitat, as identified in Section 6.3.3; that is, the Mud Snail (*Gabbiella candida*), Grey Crowned Crane (*Balearica regulorum*), Nahan’s Francolin (*Ptilopachus nahani*) and Eastern Chimpanzee (*Pan troglodytes*). Potential impacts to other species of concern are assessed at the habitat level (see Section 7.3.1.1).

**10.1.2.1 What impact could the construction/decommissioning of the Production Facility have on the Mud Snail (*Gabbiella candida*)?**

The Mud Snail (*Gabbiella candida*) is a Critically Endangered and range-restricted species. Currently, the only known populations occur around Butiaba (see Section 6.3.3.1), which is on the eastern shore of Lake Albert approximately 90 km north of the LSA. Although this species was not confirmed within the CHAA during the course of baseline studies, there is a potential that this species could occur in the near-shore habitats of the CHAA, based on its known habitat preferences, and those of other Mud Snail species (*Gabbiella* spp.), which have previously been recorded in both the same locality as this species at Butiaba, as well as in the LSA. Hence, a precautionary approach has been adopted, and *G. candida* is assumed to occur in the near-shore habitats of the CHAA.





### Impact Indicators

Indicators used to assess impacts of the construction of the Project on the Mud Snail were: habitat quantity and quality; and habitat connectivity, because no individuals were recorded, yet potential habitat is present.

Habitat loss due to direct disturbance and clearing of habitat was assessed by calculating the loss of suitable habitat from the CHAA as a result of the construction of the Project. Changes to habitat quality were assessed by the prediction of sediment loads and changes to water quality in the water column from construction activities.

Changes in habitat connectivity were assessed by identifying potential barriers to genetic movement, and source populations. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the Project footprint, to qualitatively identify areas where critical habitat becomes fragmented.

### Impact Assessment

#### ■ Habitat Quality and Quantity

Very little information is available for this species. Information about the genus *Gabbiella* indicates that this group is generally found in lakes, rivers and, less commonly, in small water bodies, and rarely in habitats that dry out (Kristensen and Stensgaard 2010). Two sister species (*G. humerosa* and *G. walleri*) were recorded in the LSA (see Section 3.3.2.4, Appendix D). These two species have also been recorded in the same locality and habitat as *G. candida* (GBIF, 2014). Therefore, the habitat preferences of those two species were used as a model to predict the potential impacts of changes to habitat quantity and quality for this species.

The two sister Mud Snail species recorded from the CHAA were collected from bottom substrates in the open sandy shore habitats in the LSA. These habitats are characterised by a gently sloping lake bed extending from the shore line to deeper water. The substrate is typically comprised of sand and finer sediments (Wandera and Balirwa 2010). This habitat constitutes approximately linear (lake shore) 10.5 km of the CHAA, most of which is located within the LSA.

The jetty will be upgraded, although there are not expected to be material changes in its dimensions. Currently, it consists of a solid concrete structure extending some 20 m into the lake to provide sufficient draught during low water periods. Although the existing jetty structure appears to have caused some accretion of sediment on the eastern side of the jetty, and some erosion on the western side, the proposed jetty upgrade is not expected to alter the geomorphological processes and sediment drift that currently govern the shoreline ecosystem of the Buhuka Flats, additional to the sediment deposition and erosion either side of the jetty that has already taken place.

The new water intake and pump station will extend a similar distance (~20 m) into the lake. The construction works will affect, through direct disturbance, approximately 0.04 ha or 0.005% of potential habitat (810 ha of near-shore aquatic habitats) for the Mud Snail in the CHAA. Additionally, there is potential for this proposed structure to affect geomorphological processes and sediment drift down-shore of Well Pad 2; which in combination with the existing jetty structure, could potentially affect the sediment drift or shoreline morphodynamics between Well Pad 2 and Bugoma Lagoon. Since sediment drift is recognised as an important driver in shoreline ecosystems, contributing to the nutrient input that drives phytoplankton, zooplankton and fish communities (Parks et al. 2013); there is potential for the Mud Snail habitat to be affected, beyond the area of primary disturbance.

The new water intake and pump station also has the potential to alter the water quality within the immediate surrounds of the construction activities through disturbance of the lake bed, and introduction of sediment into the water column over the short-term during the works. It is expected that these increased sediment loads will dissipate reasonably quickly following completion of the construction works. The sediment loads in the vicinity of the new water intake and pump station are not expected to exceed those that would normally be expected during windy periods on the lake, and the consequent turbid conditions caused by those winds. Furthermore, these construction works are not expected to permanently alter the water chemistry in the vicinity of the new water intake and pump station given the large buffering capacity of the lake compared to the scale of the works, and the short-term duration of the works. Therefore, it can be expected that the construction of the new water





intake and pump station will not affect the sediment loads and water quality of the near-shore habitats following completion of works.

Accidental spillages of small quantities of fuels and chemicals during the construction of the Production Facility components (not including significant/catastrophic spillages, which are described in Chapter 07 of the main ESIA report) could end up in River 1 and, ultimately report to the near-shore habitats of the CHAA south-west of well pad 2. There is also a real potential for accidental spillages to occur as part of the jetty upgrade works, and during the construction of the new water intake and pump station. A further risk will result from the construction and drilling of the wells. While control systems are proposed to manage contaminated stormwater and wash-water from the well pads, the presence of drilling crews on site for nearly a year, using potentially hazardous drilling fluid and other hazardous materials; and the absence of a buffer between the well pads and the lake (in the case of well pad 1, the seasonal wetland); makes it likely that occasionally contaminated drainage will reach the lake unless there is a very high level of control of day-to-day activities. Aquatic snails are highly sensitive to chemical pollutants, particularly hydrocarbons (Araujo et al. 2012). Currently, the concentrations of hydrocarbons and other pollutants in the lake waters of the near-shore habitats are below levels that could cause harm to the environment; this is supported by the healthy aquatic communities observed in the lake's near-shore habitats (see Section 6.2). As such, without a very high level of control of day-to-day activities and adequate spill prevention and clean-up measures in place during construction, accidental spillages of fuels and chemicals could, depending on the volume spilt, have a detrimental impact on the near-shore habitats of the LSA through the introduction of toxic compounds and pollutants. Such a spill could have a detrimental impact on the Mud Snail.

Discharge of treated sewerage during construction could affect water quality and algal growth rates, potentially affecting the quality of the Mud Snail's preferred habitat. However, *Gabbiella* sp. are detritivores/omnivore living on muddy lake bottoms and plants, and *Gabbiella humerosa*, the sister species of *Gabbiella candida*, appears to benefit from increased eutrophication (Van Damme & Lange, 2017). Therefore, the discharge of treated sewerage during construction may not have a detrimental impact on the Mud Snail.

The construction of the KFDA permanent and temporary camps/parking lots/materials yards, airstrip extension, CPF, new in-field roads, crusher plant/spoil area A, expansion of exploration well pads to production well pads (particularly the expansion of well pad 2 to its full size), and associated infrastructure could cause increased sedimentation of near-shore habitats on the Buhuka Flats. The soils of the Buhuka Flats are dispersive and active soil erosion is evident in the LSA. With the construction of the Project's components, and the consequent exposed areas of soil, there is a potential that, without adequate erosion and sediment control measures in place during construction, sediment loads within the watercourses (particularly River 1, which will receive the drainage from the CPF earthworks and temporary camp) draining the Project footprint could increase. These sediment-laden watercourses report to Lake Albert, and, hence, there is a potential for increased sediment loads in the near-shore habitats. Near-shore habitats particularly at risk include the lagoon, and to a lesser extent, the shallow river-associated habitats, both potential habitat for the Mud Snail. Nevertheless, the watercourses draining the LSA support dense emergent vegetation (see Section 6.1.1.1). Such vegetation forms an impactive filter for most sediment (IECA 2008), therefore, it can be expected that sediment loads reporting to the near-shore habitats, at least, via the Kamansing River, River 1 and Masika River, could be minimal. Sediment loads from overland flows may not be as retarded by vegetation, and hence may report to the near-shore habitats and affect them detrimentally. However, little construction storm water will flow into the Kamansing River and the seasonal wetlands upstream of the Bugoma Lagoon, since only the expansion of Well Pad 1 and roughly half of the construction area of the permanent camp fall within its catchment (Figure 20). The seasonal wetland will provide efficient attenuation of sediment, and a significant increase in sediment concentrations in the lake or in Bugoma lagoon are unlikely.

Impacts on habitat quality and quantity for *G. candida* arising from direct disturbance, changes in sediment dynamics and potential accidental spillages of small quantities of fuels and chemicals during construction are predicted to be beyond the expected range of natural disturbance perturbations. Potential loss of critical habitat for the Mud Snail requires the consideration of offsets to meet IFC requirements.

### ■ Habitat Connectivity





The construction of the Project is expected to last for three years. Besides the upgrading of the jetty, it is unlikely that construction activities could substantially alter the habitat connectivity of the near-shore habitats in the CHAA. No structures are being put in place that will alter the natural connectivity of the aquatic habitats of the lake. It is expected that the connectivity amongst the aquatic habitats will remain the same during construction as they were during baseline.

**Impact Classification**

The Mud Snail’s sensitivity is high because this species is Critically Endangered, and potentially triggers a Tier 1 critical habitat designation.

Impacts to the Mud Snail’s habitat quantity and quality will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA, with approximately 0.04 ha or 0.005% of potential habitat affected. Impact duration will be short-term (that is, limited to the construction phase of ~3 years) because disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. Although the jetty structure will remain in place into the far future (that is, ~25 years) after the project is decommissioned, the changes to the open sandy shore habitat in the vicinity of the jetty are expected to be reversible over time as natural geomorphological processes re-establish the open sandy shoreline. The magnitude of the physical impacts of construction on the habitat quantity and quality of the Mud Snail is expected to be low. Following the application of appropriate mitigation measures, the impact significance for physical impacts on habitat quality and quantity is expected to remain the same, that is, moderate, primarily because the magnitude will still remain the same due to habitat disturbance during construction (Table 13).

Adverse impacts on habitat quality are expected as a result of accidental spillages of small quantities of fuels and chemicals (including potentially hazardous drilling fluid) during the construction of the Project components and installation of the wells, ultimately reporting to the near-shore habitats of the Lake. Although the impact duration will be short-term, and should be reversible with time as the Lake waters dilute and disperse the contaminants, the magnitude of the impacts of accidental entrainment of contaminants to *G. candida*’s habitat could be high. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact on physical habitat quality and quantity during the construction phase of the Project, and a major overall impact on habitat quality and quantity as a result of contamination, pre-mitigation (Table 13). The strict application of the recommended mitigation measures (Section 11.2.1) will reduce the risk, and thus the likely magnitude, of potential habitat contamination for the Mud Snail, reducing the overall impact significance to moderate.

Impacts to the Mud Snail’s habitat connectivity will be neutral. The geographical extent of impacts will be local because impacts are restricted to the LSA, with approximately 0.04 ha or 0.005% of potential habitat affected. Impact duration will be short-term (that is, limited to the construction phase of ~3 years) because disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities, and no barrier to habitat connectivity will be constructed. Although the existing jetty structure will remain in place into the far future after the project is decommissioned (that is, ~25 years), the proposed upgrade works will not result in changes to current jetty footprint and subsequently the open sandy shore habitat in the vicinity of the jetty. The application of the recommended site-specific mitigation measures in the construction of the new water intake and pump station (Section 11.0) are expected to minimise any detrimental effects on longshore sediment drift and morphodynamics, and any subsequent negative effects on the habitat connectivity of the Mud Snail, to a point where the magnitude of the potential impact is negligible. Therefore, the magnitude and sensitivity combine to produce a minor overall impact on habitat connectivity for *G. candida* during the construction phase of the Project, pre-mitigation (Table 13).

**Table 13: Potential impacts in the construction phase to the Mud Snail**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance





Habitat quality and quantity (physical impacts)	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8
Habitat quality and quantity (contamination)	High – 4	High – 4	Major - 16	Low – 2	High - 4	Moderate – 8
Habitat connectivity	Negligible – 1	High – 4	Minor – 4	Negligible – 1	High – 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA, and hence potential local populations of Mud Snail. However, there is some uncertainty with regard to this species actually occurring in the CHAA. As such, the above assessment has been undertaken based on a precautionary approach.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**10.1.2.2 What impact could the construction/decommissioning of the Production Facility have on Grey Crowned Crane?**

The Grey Crowned Crane is an Endangered species. Up to 14 individuals were regularly recorded on the Buhuka flats during baseline surveys, and a pair of Grey Crowned Crane was observed on the Flats during social baseline survey work conducted in November 2017. More than three-quarters of the world’s Grey Crowned Cranes occur in Uganda and Kenya in East Africa, leading Uganda to develop a species action plan for their conservation in-country. The population of Grey Crowned Crane using the Buhuka flats is thus regarded as highly sensitive, and the wetlands of the Flats are considered an important conservation unit for the species.

The construction of the Production Facility could directly impact Grey Crowned Crane breeding and foraging habitat on, given that construction impacts on wetlands in the Buhuka Flats are expected (Section 9.1.1.2). In addition, although it is recognised that the escarpment road has already been licensed and built on the basis of an earlier impact assessment (AWE, 2014c), the induced effects of its operation on the Buhuka Flats wetland habitats used by Grey Crowned Crane are expected to exacerbate predicted construction phase impacts on this species in the Buhuka Flats locality, and this is included for completeness.

The potential direct, indirect and induced impacts of the construction of the Production Facility to the Grey Crowned Crane are presented below.

**Impact Indicators**

Indicators used to assess effects of the construction of the Production Facility on Grey Crowned Crane were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to direct disturbance and clearing of habitat was assessed by calculating the loss of suitable habitat from the CHAA as a result of the construction of the various Project components and infrastructure. Changes to habitat quality due to indirect disturbance were estimated by applying a 200m buffer to the infrastructure footprint. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance arising from light, noise, vibration, and edge effects.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat for Grey Crowned Crane in relation to the infrastructure to qualitatively identify areas where habitat becomes fragmented.





Potential changes in abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration) and site clearing activities. These disturbances were considered in light of known or inferred effects to the survival and reproduction of Grey Crowned Crane, where data on these types of effects are available.

### **Impact Assessment**

#### ■ Habitat Quality and Quantity

Grey Crowned Crane is distributed across eastern and southern Africa. Populations in many areas including Uganda have experienced rapid declines during the past 45 years (BirdLife International, 2018) for reasons including habitat loss to farming, human presence causing disturbance, collection of chicks for domestication, disruption of breeding activity, loss of roosting sites (large trees located remotely from areas frequented by humans) and dry-season fires in wetland habitats (Olupot, 2014). Its habitat preferences are diverse, including wetlands with tall emergent vegetation, open riverine woodland, shallowly flooded plains and temporary pools with adjacent grasslands, open savannas, croplands, pastures, fallow fields and irrigated areas (Archibald et al, 2018). It prefers short to medium height open grasslands adjacent to wetlands for foraging, which is consistent with the seasonally flooded grassland wetlands in the Buhuka Flats. For breeding, it prefers marshes with water 1 m deep and emergent vegetation 1 m above the water (Archibald et al, 2018); habitat which corresponds to the permanent wetlands of the Buhuka Flats.

Direct loss of suitable foraging habitat (seasonally flooded grassland) to the Project footprint will consist of 5.8 ha or 8.4% of a total of 69 ha of seasonally flooded grassland in the Buhuka Flats area. No direct losses of permanent wetlands are anticipated on the Flats. Some large trees with cultural importance have reportedly been removed from the Buhuka Flats during construction activities. Such trees are important roosting resources for Grey Crowned Crane and their loss could influence their presence in/use of the area for foraging and breeding (Olupot, 2014).

Although Grey Crowned Crane can tolerate a low degree of anthropogenic disturbance in foraging habitat (e.g. subsistence and commercial farming practises), in Ugandan wetlands it has been observed to be intolerant of human proximity within 100-200m (Olupot, 2014), flying away on approach; a factor which also affects breeding success as breeding birds flush from nests on approach, causing increased rates of predation, reduced time at the nest (either incubating or feeding), and ultimately nest abandonment. How tolerant foraging/roosting/breeding Grey Crowned Crane may be to indirect disturbances, such as noise, light, vibration and edge effects, is not known. The application of a 200m buffer around the Project infrastructure footprint indicates that approximately 4.64 ha of seasonally flooded wetland habitat will be reduced in quality as a result of sensory disturbance.

This equates to a total potential habitat loss in the Buhuka Flats region of the CHAA from direct losses from vegetation clearing, and indirect losses from sensory disturbances and edge effects associated with the Production Facility of up to 10.44 ha (0.09% of 11,579 ha of wetlands in CHAA; 15% of 69 ha seasonally flooded grassland in the Buhuka Flats).

Effects from loss of habitat are predicted to be beyond the expected range of natural disturbance perturbations (for example, fire), although not beyond the human-induced rate of degradation via cattle grazing pressure etc.

#### ■ Habitat Connectivity

The effect of the Project construction as a barrier to the movement of Grey Crowned Crane is likely to be adverse. The construction activity will create sensory disturbances in the short-term, which can elicit reduced use or complete avoidance of affected areas, thereby creating movement barriers (for example, see Kolowski and Alonso 2009, Gleeson and Gleeson 2012). It is noted that the construction of power lines between the CPF and development wells will present a significant barrier to movement for Grey Crowned Crane, as this species is susceptible to in-flight collisions with overhead powerlines (BirdLife International, 2018); however, the construction of powerlines does not form part of this impact assessment and is instead discussed in the cumulative impact assessment (ref. Chapter 17 Cumulative Impact Assessment). Construction of linear infrastructure (roads and flowlines) through wetlands has the potential to create temporary barriers to movement as a result of the associated disturbance due to human presence in the area.







Effects from loss of habitat are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

Up to 14 individuals of Grey Crowned Crane were frequently observed on the Flats during the baseline fieldwork conducted in 2014. Given the extent of their tolerance of human presence (approx. 100m – 200m), the large-scale changes in the human population on the Flats that have occurred since the construction of the escarpment road are likely to have affected Grey Crowned Crane occurrence on the Buhuka Flats.

For the purposes of this assessment, it is assumed that Grey Crowned Crane remains present in suitable habitat (permanent and seasonal wetlands) in the Buhuka Flats. The construction of the Production Facility is likely to exacerbate existing levels of sensory disturbance in the locality, with resultant effects on habitat quality and the distribution of the species. The loss of foraging habitat to the Project footprint is also likely to affect the abundance and distribution of Grey Crowned Crane in the CHAA.

Effects on the abundance and distribution of Grey Crowned Crane are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

During the baseline fieldwork conducted in May 2014, the formation of breeding pairs of observed Grey Crowned Crane was indicated. Grey Crowned Crane nests are typically constructed within or on the edges of marshes with water of 1m depth and emergent vegetation 1m high (BirdLife International, 2018). Loss and degradation of wetland habitat on the Buhuka Flats is expected to decrease the likelihood of Grey Crowned Crane selecting these areas for breeding.

As mentioned above, direct disturbance via human proximity within 100-200m causes breeding birds to flush from nests on approach (Olupot, 2014), which may result in increased rates of predation, reduced time at the nest, either incubating or feeding, and ultimately nest abandonment, affecting reproductive success. It is assumed for this assessment that indirect disturbances arising from noise, light, vibration and edge effects are also likely to affect the breeding success of Grey Crowned Crane on the Buhuka Flats.

Removal of large trees which have importance as night-time roosts and day-time shelter from the midday sun has been indicated in localised declines in Grey Crowned Crane in Uganda (Olupot, 2014); the loss of such trees is therefore expected to have implications for the survival of flocks and individuals in the area.

The installation of powerlines as part of the Project infrastructure presents a serious risk of in-flight collisions, resulting in mortalities of individual birds – these impacts are addressed in the cumulative impact assessment (ref Chapter 17 Cumulative Impact Assessment).

These effects to the survival and reproductive success of Grey Crowned Crane are predicted to be well beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Grey Crowned Crane sensitivity is high because it is Endangered, and triggers Tier 2 critical habitat.

Impacts to Grey Crowned Crane habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas within 200 m of the Production Facility infrastructure footprint, with approximately 5.8 ha (8.4%) of potential habitat in the Buhuka Flats being permanently lost, and 10.44 ha (6.6 %) of potential habitat in the Buhuka Flats indirectly affected. Impact duration will be short-term (that is, limited to the construction phase of the Project, that is, ~3 years). Changes to the habitat quality and quantity from sensory disturbances arising from construction activities are expected to be reversible after completion of the works. The magnitude of the effects of construction on Grey Crowned Crane habitat quantity and quality is considered medium. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to



moderate, because the magnitude of effects could be reduced to low, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to Grey Crowned Crane habitat connectivity will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas where wetlands will be crossed by linear infrastructure i.e. access roads. Impact duration will be long-term, as the presence of access roads will commence during construction and remain in place throughout the operational lifetime of the Project, although sensory disturbances arising from construction activities are expected to dissipate within a short time after completion of the construction phase. The magnitude of the effects of construction on Grey Crowned Crane habitat connectivity is low, as the species is mobile and capable of flight to preferred areas. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the magnitude will remain low as long as the roads are present, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to Grey Crowned Crane abundance and distribution will be adverse. The geographical extent of impacts will be local because effects are restricted to wetlands within 200m of the proposed Production Facility infrastructure on the Buhuka Flats. Impact duration will be short-term (that is, limited to the construction phase of ~3 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. There is a possibility that individuals could be killed or severely disturbed during the construction phase; however, in the context of overall species abundance, these losses or disturbances are expected to be reversible after completion of the works. The magnitude of the effects of construction on Grey Crowned Crane abundance and distribution is thus considered medium. Therefore, the magnitude and sensitivity combine to produce an impact of major significance during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to moderate, because the magnitude of effects could be reduced to low, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to the survival and reproduction of Grey Crowned Crane will be adverse. The geographical extent of impacts will be local because effects are restricted to wetlands within 200m of the proposed Production Facility infrastructure on the Buhuka Flats. Impact duration will be short-term (that is, limited to the construction phase of ~3 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. A threshold of 10% for this species' survival and reproduction in the CHAA is reasonable, and it is considered probable that at least this number of individuals in the local population within the Buhuka Flats will be affected. The magnitude of the effects of construction on the survival and reproduction of the Grey Crowned Crane is therefore medium. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to moderate, primarily because the intensity could become low, although the sensitivity of the receptor will remain the same (Table 14).

**Table 14: Potential construction phase impacts to Grey Crowned Crane**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Medium - 3	High - 4	Major - 12	Low - 2	High - 4	Moderate - 8
Habitat connectivity	Low - 2	High - 4	Moderate - 8	Low - 2	High - 4	Moderate - 8





Abundance and distribution	Medium - 3	High – 4	Major - 12	Low – 2	High – 4	Moderate - 8
Survival and reproduction	Medium - 3	High – 4	Major - 12	Low – 2	High – 4	Moderate - 8

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts on the Grey Crowned Crane and its habitat in the CHAA.

The spatial extent of the wetland habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**10.1.2.3 What impact could the construction/decommissioning of the Production Facility have on Nahan’s Francolin?**

Nahan’s Francolin is a Vulnerable and range-restricted species. It has a very restricted distribution, being found only in north-east DRC and western and south-central Uganda; in particular, the forests of Budongo, Bugoma and Mabira (McGowan and de Juana 1994). It has been recorded within Bugoma Central Forest Reserve within the CHAA.

This species triggers Tier 2 critical habitat (Section 6.3.2.2, Appendix G) under Criterion 2. Tier 2 habitats are considered to be sensitive, and, therefore, if a project is located in such a habitat, the IFC considers that compliance with the provisions of paragraph 17 of PS 6 (IFC 2012a) would be difficult. In summary, a project will not be developed in Tier 2 habitat unless: no other viable alternatives exist; and, the project does not lead to measurable and irreversible adverse impacts to the valued component that triggered critical habitat; and, the project does not lead to a net reduction in the global and/or national/regional population of the triggering species (such as Nahan’s Francolin) over a reasonable period of time; and, a robust, appropriately designed, and long-term biodiversity monitoring and evaluation programme is part of the project’s Environmental and Social Management System (ESMS). A Biodiversity Action Plan (BAP) will then be developed to achieve net gain for the affected species.

Although it is expected that the construction of the Production Facility will not have direct impacts to Nahan’s Francolin, the proposed upgrade of the R5 and P1 roads by the Ugandan Government for the benefit of the Project (Figure 21) could have induced and cumulative impacts to this species and Bugoma Central Forest Reserve (see also Section 7.3.1.1.4). The IFC’s PS1, paragraph 2, recognises that certain effects and impacts arising from a Project may be “the responsibility of the government or other third-parties over which the client does not have control or influence” (IFC 2012c). Nevertheless, although CNOOC cannot control the actions of the government, the ESMS for the Project must identify the Ugandan Government’s role, and the corresponding risks they present to CNOOC and the Project (IFC 2012c). Furthermore, the ESMS must identify opportunities for CNOOC to collaborate with the Ugandan Government in order to achieve outcomes that are consistent with PS6.

The proposed Government upgrade of R5 and P1 roads is one such opportunity. The potential induced and cumulative effects of that proposed development to this species are presented below.

**Impact Indicators**

Indicators used to assess effects of the upgrade of the R5 and P1 on Nahan’s Francolin were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to direct disturbance and clearing of habitat was assessed by calculating the loss of suitable habitat from the CHAA as a result of the upgrade of the R5 and P1 roads, approximately 9.7 km and 3.5 km respectively of which passes through Bugoma CFR. Direct habitat loss due to vegetation clearance for road widening was estimated using a footprint of 10 m width. Habitat loss due to indirect disturbance and edge





effects was estimated by applying a 200 m buffer to the road corridor. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance like light, noise, vibration, and edge effects.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the road corridor to qualitatively identify areas where habitat becomes fragmented.

Potential changes in abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration) and site clearing activities. These disturbances were considered in light of known or inferred effects to the survival and reproduction of other francolin/buttonquail species for which data on these types of effects are available.

### **Impact Assessment**

#### ■ Habitat Quality and Quantity

This species is confined to dense, mature, moist, sometimes swampy medium-altitude forest below 1,500 m (McGowan and de Juana 1994, BirdLife International 2014i); and is reasonably common in Budongo Central Forest Reserve (Plumptre et al. 2010, 2011). Suitable habitat was predicted to cover an area of ~35,201 ha (352 km<sup>2</sup>) in the CHAA, principally in the Bugoma Central Forest Reserve. Direct habitat loss of suitable habitat as a result of clearing to widen the road corridor will be 9.7 ha of forest habitat along the R5 section of road, and 3.5 ha along the P1 section of road (based on approximately 9.7 km of the R5 and 3.5 km of the P1 traversing dense forest habitat (Figure 21), and a conservative clearing width of 10 m either side of the current road), representing 13.2 ha or 0.03% of suitable habitat present at baseline.

The Nahan's Francolin's degree of vulnerability to disturbance is unknown. Birds are known to be sensitive to land use and habitat alteration (Lussier et al. 2006). The behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions such as the availability of alternative foraging sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the population. How tolerant the Nahan's Francolin may be to indirect disturbances, such as noise, light, vibration and edge effects, is not known. However, assuming it is sensitive to such disturbance because it is a shy, forest-dependent species (Sande et al. 2009a), then with a 200 m buffer applied to the road upgrade's footprint, approximately 264 ha of potential Nahan's Francolin habitat will be lost or reduced in quality as a result of edge effects, and possibly sensory disturbance. This equates to a potential habitat loss in the CHAA from vegetation clearing, sensory disturbances and edge effects of up to 277 ha (0.8%).

Effects from loss of habitat are predicted to be well beyond the expected range of natural disturbance perturbations (for example, fire), although not beyond the human induced deforestation rate.

#### ■ Habitat Connectivity

The effect of the road upgrade as a barrier to the movement of Nahan's Francolin is unknown. Roads are recognised as creating sensory disturbances, which can elicit reduced use or complete avoidance of affected areas, thereby creating movement barriers (for example, see Kolowski and Alonso 2009, Gleeson and Gleeson 2012). However, given that the R5 and P1 already exist, if the Nahan's Francolin is sensitive to such effects, then it is reasonable to assume the existing roads are already a barrier to a greater or lesser degree. The upgrade of the road will bring increased sensory disturbance in the short-term, during upgrade construction, and then a potential for long-term effects as traffic along the road increases. Certainly, for the duration of the construction of the Project and the movement of components to the Buhuka Flats, traffic volumes on the road are expected to increase significantly (ref. Section 9.1.1.4).

Effects from loss of habitat connectivity are predicted to be well beyond the expected range of natural disturbance perturbations (for example, fire), although not beyond the human induced deforestation rate.

#### ■ Abundance and Distribution



Nahan's Francolin is reported to be relatively common in the Bugoma Central Forest Reserve (Plumptre et al. 2011). What effects the upgrade of the road, and the predicted increased traffic levels along the road during the construction of the Project, could have on the abundance and distribution of this species is unknown. Although this species within the forest is reported to be relatively common (*viz.*, Plumptre et al. 2011), the distribution of individuals within this habitat is unknown.

It is conceivable that the current road could act as a barrier and sensory disturbance to this species, and, as such, its distribution and abundance around the road corridor could be affected. That is, the local population along the road could be less than the surrounding forest. However, the actuality of this scenario is unknown.

Consequently, it is assumed that the current distribution of this species is evenly spread within suitable habitat in the Bugoma Central Forest Reserve, including along the current road. With the construction of the road upgrade, and the increased traffic along the road during the construction of the Project, it is reasonable to assume that the distribution of this species may be altered. Individuals may avoid the resultant sub-optimal habitat brought about by the upgrade construction activities and increased traffic volumes. Additionally, the increased traffic on the road could lead to an increase in direct mortality of individuals, with individuals potentially being killed by that traffic.

Effects from the upgrade construction and increased traffic volumes during construction of the production facility are predicted to be well beyond the expected range of natural disturbance perturbations, although not beyond the disturbance from the human-induced deforestation rate.

### ■ Survival and Reproduction

The Nahan's Francolin's degree of vulnerability to direct disturbance, particularly during the breeding season, is not well understood. Nahan's Francolin is reliant upon large trees, with appropriate buttress formation, for breeding sites (Sande et al. 2009a). The reduction of the number of suitable breeding sites, through the removal of such large trees, can, therefore, be expected to reduce the breeding success of this species. Generally, for birds, the behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions such as the availability of alternative breeding sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the local population of Nahan's Francolin.

How tolerant the Nahan's Francolin may be to indirect disturbances, such as noise, light, vibration and edge effects, during the breeding season, are also not completely understood. What is known is that disturbance of mature forest generally appears to reduce the home range of this species (Sande et al. 2009b), and hence its potential to find suitable mates. Assuming it is sensitive to sensory disturbance because it is a shy, forest-dependent species (Sande et al. 2009a), the construction of the road upgrade, and the disturbance arising from the increased traffic levels on the road during the construction of the Project could affect the breeding success of those individuals living in close proximity to the road. In addition, if the birds attempt to cross the upgraded road in order to forage within their usual range, they will risk collision with Production Facility construction-related traffic, resulting in mortality.

These effects to the survival and reproductive success of Nahan's Francolin are predicted to be well beyond the expected range of natural disturbance perturbations, although not beyond the human induced deforestation rate.

### **Impact Classification**

The Nahan's Francolin's sensitivity is high because it triggers Tier 2 critical habitat.

Impacts to the Nahan's Francolin's habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas within 200 m of the R5 and P1 road corridors in the CHAA, with approximately 277 ha (0.8%) of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade construction, and the construction phase of the Project, that is, ~2 years).



Changes to the habitat quality and quantity from sensory disturbances arising from construction activities are expected to be reversible after completion of the works. The magnitude of the effects of construction on the habitat quantity and quality of the Nahan's Francolin is low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 15). Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 15).

Impacts to the Nahan's Francolin's habitat connectivity will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas within the 10 m R5 and P1 road clearance footprint in the CHAA, with approximately 13.2 ha or 0.03% of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade and construction phase of ~2 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. Changes to the habitat connectivity from sensory disturbances arising from construction activities are expected to be reversible after completion of the works. The magnitude of the effects of construction on the habitat connectivity of the Nahan's Francolin is Low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 15).

Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 15).

Impacts to the Nahan's Francolin's abundance and distribution could be adverse. The geographical extent of impacts will be local because effects are restricted to those within 200 m of the R5 and P1 road corridors in the CHAA, with approximately 277 ha (0.8%) of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade and construction phase of ~2 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. There is a possibility that individuals could be killed or severely disturbed during the construction phase; however, these losses or disturbances are expected to be reversible after completion of the works. The magnitude of the effects of construction on the abundance and distribution of the Nahan's Francolin is Low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 15).

Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 15).

Impacts to the survival and reproduction of Nahan's Francolin could be adverse. The geographical extent of impacts will be local because effects are restricted to those areas within the 10 m R5 and P1 road clearance footprint in the CHAA, with approximately 13.2 ha or 0.03% of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade and construction phase of ~2 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. No large, buttressed trees, utilised as breeding sites, are expected to be removed during the road upgrade. There is a possibility that individuals could be killed or severely disturbed during the construction phase; however, those losses or disturbances are expected to be reversible after completion of the works. A threshold of 10% for this species' survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve will not be affected. The magnitude of the effects of construction on the survival and reproduction of the Nahan's Francolin is therefore low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 15).

Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 15).

### **Table 15: Potential impacts in the construction phase to the Nahan's Francolin**



Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Habitat connectivity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Abundance and distribution	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Survival and reproduction	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential effects that the upgrade works associated with the R5 and P1 roads, and traffic associated with the construction of the Project, could have on the Nahan’s Francolin and its habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**10.1.2.4 What impact could the construction/decommissioning of the Production Facility have on Eastern Chimpanzee?**

The Eastern Chimpanzee is an Endangered species. The population of Eastern Chimpanzees in the Bugoma Central Forest Reserve is recognised as being one for the four largest in the region (Plumptre et al. 2010). Hence, the forest is recognised as an important chimpanzee conservation unit by the IUCN (Plumptre et al. 2010).

This species triggers Tier 1 critical habitat (Section 6.3.2.2, Appendix G). Tier 1 habitats are considered to be very sensitive, and, therefore, if a project is located in such a habitat, the IFC considers it unlikely that the client will be able to comply with the provision of PS 6, in particular paragraphs 17, 18 and 19 (IFC 2012a, as presented in Appendix C). In summary, a project will not be developed in Tier 1 habitat unless: no other viable alternatives exist; and, the project does not lead to measurable and irreversible adverse impacts to the valued component that triggered critical habitat; and, the project does not lead to a net reduction in the global and/or national/regional population of the triggering species (i.e. Eastern Chimpanzee) over a reasonable period of time; and, a robust, appropriately designed, and long-term biodiversity monitoring and evaluation programme is part of the project’s Environmental and Social Management System (ESMS). A Biodiversity Action Plan (BAP) will be developed to achieve net gain for the affected species.

Although it is expected that the construction of the Production Facility will not have direct impacts to Eastern Chimpanzees, the proposed upgrade of the R5 by the Ugandan Government (Figure 21) for the benefit of the Project could have induced and cumulative impacts to this species and Bugoma Central Forest Reserve (also see Section 6.1.1.4). The IFC’s PS1, paragraph 2, recognises that certain effects and impacts arising from a Project may be “the responsibility of the government or other third-parties over which the client does not have control or influence” (IFC 2012c). Nevertheless, although CNOOC cannot control the actions of the government, the ESMS for the Project must identify the Ugandan Government’s role, and the corresponding risks they present to CNOOC and the Project (IFC 2012c). Furthermore, the ESMS must identify opportunities for CNOOC to collaborate with the Ugandan Government in order to achieve outcomes that are consistent with PS6.





The proposed Government upgrade of the R5 is one such opportunity. The potential induced and cumulative effects to from this upgrade to the Eastern Chimpanzee are presented below.

### *Impact Indicators*

Indicators used to assess effects of the upgrade of the R5 on Eastern Chimpanzee were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to direct disturbance and clearing of habitat was assessed by calculating the loss of suitable habitat from the CHAA as a result of the upgrade of the road. Habitat loss due to indirect disturbance and edge effects was estimated by applying a 500 m buffer to the road corridor. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance like noise, vibration and traffic. The buffer width was determined based known chimpanzee sensitivities to noise disturbance (Parren and Byler 2003, Rabanal et al. 2010). The amount of loss or degradation of habitats within the buffer was evaluated across a range of possibilities, including that habitats become completely unavailable to chimpanzees during the construction phase. Evaluating the potential for complete avoidance of the buffer area is a conservative approach, which addresses uncertainty about the attenuation distance of sensory disturbances for chimpanzees, even though the likelihood of strict avoidance throughout the entire buffer may be low.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the road corridor to qualitatively identify areas where habitat becomes fragmented.

Potential changes in abundance and distribution were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration) and site clearing activities. These disturbances were considered in light of known or inferred effects to the survival and reproduction of other populations of chimpanzees for which data on these types of effects are available.

To assess effects to survival and reproduction as a result of in-migration and potential associated increases in poaching and disease spread, in-migration rates were predicted based on the predictions in the baseline survey report. A literature review of the impact of contact with humans was also conducted

### *Impact Assessment*

#### ■ Habitat Quality and Quantity

Eastern Chimpanzees appear to range throughout the Bugoma Central Forest Reserve, both on the northern southern sides of the existing P1 road (Plumptre et al. 2010). Chimpanzees build nests to sleep in each night (Thompson and Wrangham 2013). Based on the data presented in Plumptre et al. (2010), the distribution of nesting sites appears to be widely distributed through the forest; the exception being within the vicinity of the existing Nsozi-Kabwoya road. Here the frequency of nest encountered by those workers was a lot lower than elsewhere in the forest. This suggests that the Eastern Chimpanzees within the Bugoma Central Forest Reserve are actively avoiding the road, preferring to sleep some distance away from the disturbances arising from the road. The observation supports the findings of Parren and Byler (2003) that chimpanzees actively avoid environments where they will be disturbed at night.

Based on the above, the habitat along the current P1 and R5 roads could be considered foraging, or non-core, habitat (after Parren and Byler 2003). The entire Bugoma Central Forest Reserve is considered suitable habitat for Eastern Chimpanzees, as well as the areas beyond the boundaries of the forest reserve (see McLennan 2008). However, for the intents of this impact assessment, the habitat within and immediately surrounding the Bugoma Central Forest Reserve was only considered.

Suitable habitat was predicted to cover an area of ~40,200 ha (402 km<sup>2</sup>) in the CHAA, principally in the Bugoma Central Forest Reserve. Direct loss of suitable habitat as a result of clearing to widen the road corridor will be approximately 26.4 ha (based on ~9.7 km of the proposed R5 upgrade, and 3.5 km of the proposed P1 upgrade, traversing suitable habitat (Figure 21), and a conservative clearing width of 10 m either side of the current road for upgrade purposes), representing 0.07% of critical habitat present at baseline.







The degree of vulnerability to disturbance experienced by chimpanzees is reasonably well known (for example, see Parren and Byler 2003, Rabanal et al. 2010, Thompson and Wrangham 2013). The chimpanzees living in and around the Bugoma Central Forest Reserve most likely experience sensory disturbances at present from human activities, given the high human populations living around the reserve. Indeed, groups are known to forage in the agricultural fields surrounding the forest, and hence, would more than likely be used to human noises and disturbances (McLennan 2008). As such, the potential sensory disturbances arising from the road upgrade construction and increased traffic during the Project's construction are likely to be minimal. Nevertheless, the magnitude of noise may not be the most important determinant of chimpanzee response. Instead, chimpanzees may respond to 'new' noises or may associate particular noises with other occurrences (for example, machine noise may be associated with human presence, which chimpanzees may, in turn, associate with the presence of danger). Where humans pose a threat, chimpanzees generally avoid them (Hockings and Humle 2009, Parren and Byler 2003). Therefore, the degree of avoidance may depend on the behaviour of people, highlighting the importance of managing contractor activity.

Avoidance of industrial activity, like earth moving, by chimpanzees also varies. Chimpanzees have been recorded to leave their range as a result of logging activities heard from a distance of 5 to 10 km, and there are suggestions that this could cause lasting avoidance of disturbed areas (Parren and Byler 2003). Such avoidance may explain why chimpanzee densities were consistently lower in logged areas in Kibale National Park, compared to unlogged areas, although avoidance of hunting as a result of logging activity may also have been a factor (Chapman and Lambert 2000). However, Rabanal et al. (2010) did not find large-scale spatial responses to oil and gas related noise disturbance, in Loango National Park, Gabon; although chimpanzees avoided sites where explosions were used for exploration for a period of four months after the activity had ceased (Rabanal et al. 2010). The chimpanzees within the Bugoma Central Forest Reserve are, therefore, likely to show localised patterns of avoidance, particularly near the immediate road corridor during upgrade construction phase when noise and human activity will be most intense.

The introduction and spread of invasive species during the road upgrade is not expected to cause a change in habitat quality and quantity for the Eastern Chimpanzees. Populations of invasive species currently occur along the road corridor. The construction activities could create new sites for the colonisation of invasive species present in the area, although it is doubtful that the current populations of these species will increase dramatically.

With the above in mind, then with a 500 m buffer applied to the road upgrade's footprint, approximately 660 ha of Eastern Chimpanzee habitat will be reduced in quality as a result of sensory disturbance. This equates to a total potential habitat loss in the CHAA from vegetation clearing, sensory disturbances and edge effects of up to 686.4 ha (1.7%). The reality of this quantity is doubtful given that upgrade construction works may not occur along the entire length at once; however, there is a potential that the sensory disturbance arising from the increased traffic associated with the Project's construction may affect the habitat quality along the length of the road through the forest, as there is presently very little traffic on the roads through the centre of the reserve.

Although the effects on habitat quantity are probably not beyond the human induced deforestation rate, the disturbance arising from the Project construction traffic and subsequent effects on habitat quality is predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Habitat Connectivity

The upgrade of the R5 and P1 Roads will directly remove suitable habitat for the Eastern Chimpanzees. It may also affect that habitat within the 500 m buffer through indirect impacts like edge effects and sensory disturbance. However, the chimpanzees within Bugoma Central Forest Reserve are assumed to be accustomed to human activity, and they may, more than likely, regularly cross roads within their range. Therefore, it is possible that they will not avoid the road corridor when construction activity is not occurring. However, during upgrade construction works, they may avoid those sections where construction activity is occurring.

While the Bugoma Central Forest Reserve chimpanzees are accustomed to some human activity, including occasional road traffic, and they are known to regularly cross roads within their range, the magnitude of the



impact caused by construction activities (both road building and particularly, the CNOOC production facility construction traffic) is likely to be significantly greater than is currently experienced. The probable split between construction traffic on the R5 and the P1 is not known, nor is the extent of avoidance behaviour by the chimpanzees to increasing degrees of nuisance and perceived threat. However, it is reasonable to assume that the order of magnitude increase in traffic will materially affect the behaviour of the animals, and will discourage regular road crossings. As such, the impact of the increased Project construction traffic as a barrier to chimpanzee movements is predicted to be of moderate magnitude.

### ■ Abundance and Distribution

The Bugoma Central Forest Reserve supports one of the top four Eastern Chimpanzee populations in Uganda, with a population of between 450 and 850 individuals (Plumptre et al. 2010). The construction works associated with the upgrade of the R5 Road are not expected to detrimentally affect the abundance of chimpanzees in the forest. Their localised distribution may be altered temporarily as they avoid sensory disturbances associated with the construction works. However, these distributions are expected to return to baseline conditions when the works cease.

Construction traffic will increase to frequencies where interaction between vehicles and animals crossing the road could be likely on occasions. How the chimpanzees would behave in the face of an oncoming vehicle is unknown. The probability of collisions and the potential magnitude of this impact on abundance and distribution is still thought to be low, but it is no longer negligible, as is the case at present where traffic volumes are limited.

Effects from the increased traffic volumes arising from the construction of the Production facility are predicted to be well beyond the expected range of natural disturbance perturbations, although not beyond the disturbance from the human-induced deforestation rate and habitat loss, and bush meat hunting in Bugoma Central Forest Reserve.

### ■ Survival and Reproduction

As mentioned, the chimpanzees within the forest appear to currently avoid the road corridor for night-time nesting and other activities (after Plumptre et al. 2011). Furthermore, given that the chimpanzees within the forest are more than likely used to human activities in and around the forest, they are predicted to adapt to most of the sensory disturbance arising from the construction activities. The survival and reproduction of the Eastern Chimpanzees within the Bugoma Central Forest Reserve are not expected to be affected as a result of the road upgrade construction works; however, the increased traffic associated with the Project's construction presents an increased collision risk and risk of injuries and mortalities.

The increased traffic on the upgraded road, associated with the Project construction, has the potential to cause direct mortality of individuals, should such individuals cross the road. What effect the loss of individuals from the population could have is unknown. It is doubtful that the population could be reduced by 10% and, hence, reach that critical population threshold due to road mortalities alone; however, any mortality or injury to individuals of Eastern Chimpanzee as a result of collisions with Project construction vehicles is considered unacceptable.

These effects to the survival and reproductive success of Eastern Chimpanzees are predicted to be beyond the expected range of natural disturbance perturbations, although not beyond the human induced deforestation rate, habitat degradation and bush meat hunting. In the short-term, the survival and reproduction of individual chimpanzees within the forest could be detrimentally affected by the significant increase in construction traffic associated with the construction of the Production Facility.

### **Impact Classification**

The Eastern Chimpanzee's sensitivity is high because it is Endangered.

Impacts to the Eastern Chimpanzee's habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 and P1 Road corridor in the CHAA, with approximately 504 ha (1.2%) of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade construction, and the construction phase of the Project, that is, ~3 years). Changes to the habitat quality and quantity from sensory disturbances arising from construction



activities are expected to be reversible after completion of the works. The magnitude of the effects of construction on the habitat quantity and quality of the Eastern Chimpanzee is low. Therefore, the intensity and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 16). Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 16).

Impacts to the Eastern Chimpanzee's habitat connectivity are expected, primarily as a result of the significant increase in traffic associated with the trucks being used during construction of the Production Facility. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to road corridor in the CHAA, with approximately 504 ha, or 1.2%, of potential habitat affected. Impact duration will be short-term (that is, limited to the construction phase of ~3 years) because physical and sensory disturbances arising from the road upgrade works, and the physical barrier to movement presented by the Production Facility construction traffic are expected to dissipate within a short time after cessation of upgrade activities, and so are expected to be reversible after completion of the works. The magnitude of the effects of construction of the Production Facility on the habitat connectivity of the Eastern Chimpanzee is thus considered medium. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 16). Following the application of recommended mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude would become negligible, although the sensitivity of the receptor will remain the same (Table 16).

Impacts to the Eastern Chimpanzee's abundance and distribution, primarily as a result of the increased traffic volumes during construction of the Production Facility, could be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 Road corridor within the Bugoma CFR. Impact duration will be short-term (that is, limited to the road upgrade and Production Facility construction phase of ~3 years) because physical and sensory disturbances arising from construction activities and traffic are expected to dissipate within a short time after cessation of activities. Although there is a possibility that individuals could be killed or severely disturbed during the construction phase, this is unlikely to have a material effect on the species abundance or distribution of Eastern Chimpanzee. The magnitude of the potential construction impacts on the abundance and distribution of the Eastern Chimpanzee is low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 16). Following the application of the recommended mitigation measures (Section 11.0), the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (Table 16).

Impacts to the survival and reproduction of population of Eastern Chimpanzee in Bugoma Central Forest Reserve could be adverse. The geographical extent of physical and sensory disturbances will be local because effects are restricted to those areas immediately adjacent to the R5 Road corridor in the CHAA, with approximately 504 ha, or 1.2%, of potential habitat affected. Impact duration will be short-term (that is, limited to the road upgrade and construction phase of ~3 years) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. Indications are that the chimpanzees in Bugoma Central Forest Reserve nest away from the current road, and therefore, it can be expected that effects to their survival and reproduction due to physical and sensory disturbances will be minimal.

However, in the event that the chimpanzees do need to cross the upgraded road, the increased traffic associated with the Project's construction could present an unacceptable collision risk and subsequent injuries/mortality of chimpanzees. Therefore, in the short-term, the survival and reproduction of chimpanzees within the forest could be detrimentally affected by the increased vehicular traffic that will be present during the construction phase of the Project. A threshold of 10% for this species' survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve will not be affected through direct mortality or severe sensory disturbance; nevertheless, a single incidence of mortality or injury to any individual of this Endangered species is considered unacceptable. Therefore, the magnitude of the effects of increased traffic on the P1 associated with the



construction of the Project on the survival and reproduction of the Eastern Chimpanzee is medium. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 16). Following the application of the recommended mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, and the sensitivity of the receptor will remain the same (Table 16).

**Table 16: Potential impacts in the construction phase to the Eastern Chimpanzee**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Habitat connectivity	Medium - 3	High – 4	Major – 12	Negligible – 1	High – 4	Minor – 4
Abundance and distribution	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Survival and reproduction	Medium - 3	High – 4	Major – 12	Medium - 3	High – 4	Moderate - 12

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential effects that the upgrade works associated with the R5 and P1 Roads, and traffic associated with the construction of the Project, could have on the Eastern Chimpanzees and their habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.1). The actual extent of possible habitat may have been over-estimated.

**10.2 Operation Phase Impacts**

For the assessment of impacts during the operations phase, the key questions were divided into sub-questions that focused on individual valued components within the CHAA and LSA. In answering each question, the individual components of the Project were considered with regards to their potential to affect a valued component. These questions are presented below.

**10.2.1 What impact could the operation of the Project have on habitats and ecosystem integrity?**

This section presents the assessment of impacts that the operation of the Project could have on the habitat and ecosystem integrity within the CHAA and the LSA. These habitats either do, or could, support populations of species of concern. Therefore, the assessment of potential impacts to those species, and others, occurring in the CHAA has been assessed in this section through the determination of the impacts to potential habitat for those species.

The impacts of the Project on critical habitat, as triggered by species of concern, are covered under the individual assessment of those species in Section 7.2. Other triggers of critical habitat are discussed as relevant in the appropriate sections, and in Appendix G.





### 10.2.1.1 *What impact could the operation of the Project have on the near-shore environment of Lake Albert?*

#### **Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the near-shore environment of Lake Albert were changes in: regional representativeness; topography (geomorphology), sediments, water quality; ecosystem composition; ecosystem configuration.

Additional, indirect impacts to habitat were estimated by applying a 1 km buffer to the Project footprint, forming the LSA. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by sensory disturbance, changes in water quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential succession changes in species composition that could occur. This was accomplished by examining available literature about the ecology of Lake Albert, and scientific literature about the impacts of human activities on aquatic environments.

#### **Impact Assessment**

##### ■ Representativeness

The operation of the KFDA camps/parking lots/materials yards, airstrip, CPF, pipeline, in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure will not cause the loss of additional near-shore aquatic habitat beyond that already removed during the construction phase.

The potential for influx and in-migration of people onto the Buhuka Flats seeking opportunities from the Project, and because of the easier access provided by the escarpment road, could place increased pressure on the near-shore habitats. This increased pressure could arise from increased fishing activities and pollution (such as, fuels, human and livestock waste, fish waste, and litter). The resultant increased pressures could lead to a change in the current drivers of change to the near-shore aquatic habitats, although it is doubtful that these changes would lead to a change in the representativeness of habitats.

Impacts from the changes to representativeness are predicted to be within the expected range of natural disturbance perturbations.

##### ■ Topography (geomorphology) and sediment transport

The operation of the KFDA camps/parking lots/materials yards, airstrip, CPF, pipeline, in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure could cause increased sedimentation of near-shore habitats on the Buhuka Flats. The soils of the Buhuka Flats are dispersive, and active soil erosion is evident in the LSA; which could be exacerbated by increased areas of bare ground and deterioration of land capability (due to overgrazing and more extensive presence of hardstanding areas) in the LSA. Erosion around cleared areas around the site could lead to the accumulation of sediment upstream of the points where the infrastructure crosses the drainage paths (ref. Vol 4A Physical Environment). It is expected that most areas that were cleared of vegetation during the construction phase and where no infrastructure is located, will have been revegetated by the operations phase (see Vol 1b, 2 & 3 ESMPs). Dust generation, erosion and sedimentation is, therefore, likely to be limited to the drainage associated with the main access road. Sediment-laden stormwater runoff from this road will enter the near-shore environment of the Lake via the Kamansinig River, to the north of the lagoon.

If adequate erosion and sediment control structures are not maintained as part of the prescribed stormwater management system during the operation of the Project, there is a potential that sediment loads within the watercourses draining the Project footprint could increase. These sediment-laden watercourses report to Lake Albert, and, hence, there is a potential for increased sediment loads in the near-shore environments. Near-shore habitats particularly at risk include the lagoon, and to a lesser extent, the shallow river-associated habitats. Nevertheless, the watercourses draining the LSA support dense emergent vegetation (see Vol 4a). Such vegetation forms an impactive filter for most sediment (IECA 2008), therefore, it can be expected that sediment loads reporting to the near-shore habitats, at least, via the Kamansinig River, River 1 and Masika River, could be minimal. Sediment loads from overland flows may not be retarded by vegetation, and hence





may report to the near-shore habitats, contributing to measurable increased turbidity during and after storms, where River 1 discharges to the Lake.

Impacts from the changes to topography (geomorphology) and sediment are predicted to be beyond the expected range of natural disturbance perturbations if the mitigation measures fail or are inadequate.

### ■ Water quality

Impacts from the changes to water quality are predicted to be beyond the expected range of natural disturbance perturbations, if the mitigation measures fail or are inadequate. Contributing factors to potential changes in water quality are discussed below:

#### *Discharge of poor quality sewerage effluent*

The discharge of treated sewerage effluent from the plant to Lake Albert presents an increased risk of eutrophication of the near-shore habitats on the Buhuka Flats. Sewage from the CPF will be routed via conservancy tanks to a regulating tank at the permanent camp, from where it will be treated in a Membrane Bioreactor sewage treatment works (ref, Vol1b Chapter 2). Options for final disposal of treated sewage effluent include the base case (discharge into perimeter drains around the CPF, which discharge into small drainage lines leading to Lake Albert), irrigation onto land in the buffer area around the CPF and at the personnel camp lawns and gardens, and/or discharge into an artificial wetland or sustainably managed plantation.

#### **Oil and Chemical Pollution**

Accidental, minor spillage of fuels and chemicals during the day-to-day operation of the Project components (other than catastrophic spillages, which are addressed in Vol 1b) could report to the near-shore habitats of the CHAA, via stormwater drainage into River 1, and subsequently Lake Albert, south west of Well Pad 2. In this way, minor spillages during operation could ultimately report to the near-shore habitats of the CHAA. Further sources of contamination could occur around the jetty when barges are being loaded and unloaded. This Project component is located right on the lake shore, and the impact of an accidental spill would be more obvious than a land-based spill; it would also be harder to contain and clean up.

These risks must be assessed in the context of the high sensitivity of the near-shore environment to oil and chemical spills. Certain invertebrate species (for example, aquatic snails (Araujo et al. 2012), mayflies (Savić et al. 2011)) and juvenile fish (for example, Agamy 2013) are highly sensitive to chemical pollutants, particularly hydrocarbons. Currently, the concentrations of hydrocarbons and other pollutants in the lake water are below levels that could cause harm to the environment (Vol 4a); this is supported by the healthy aquatic communities observed in the lake's near-shore habitats. As such, without adequate spill prevention and clean-up measures in place, a chemical spill could, depending on the volume spilt, have a detrimental impact on the near-shore habitats of the CHAA through the introduction of toxic compounds and pollutants. Such a spill could have a detrimental impact on the aquatic invertebrate community and juvenile fish occurring in these habitats; including the Critically Endangered Mud Snail (*G. candida*) (if it does occur in the area).

#### **Discharge of poor quality storm water**

Potentially Oil Contaminated (POC) stormwater generated in the defined hazardous areas of the plant will be collected in the open drain system for delivery to an API oil separator. These API separators are designed to separate gross amounts of oil and suspended solids from the water. The first 15 minutes of any storm will be captured and routed through the API separator, before being delivered to the secondary treatment section of the produced water treatment system for further treatment and disposal with produced water. A maximum 15-minute stormwater runoff value of 120 m<sup>3</sup> (equivalent to runoff of 478 m<sup>3</sup>/hr) is provided for. The balance of any stormwater will be captured in a stormwater pond, tested and released into the environment, if it meets the discharge specification. All stormwater from designated non-hazardous areas of the plant will be released directly from the open drains, without testing.

While control systems are proposed to manage contaminated stormwater and wash water from the well pads, the absence of a buffer between the well pads and the lake (or, in the case of well pad 1, the seasonally-flooded grassland wetland); makes it likely that occasionally-contaminated drainage could reach the near-shore habitats of the lake, unless there is a very high level of control of day-to-day activities.





### **Population increases**

The potential for influx and in-migration of people onto the Buhuka Flats seeking opportunities from the Project, and because of the easier access provided by the escarpment road, could place increased pressure on the near-shore habitats. This increased pressure could arise from increased pollution (such as fuels, human and livestock waste, fish waste, and litter), particularly nutrient enrichment from sewerage and livestock. The resultant increase in pollution levels could lead to a dramatic change in the current drivers of change to the near-shore aquatic habitats, especially the Bugoma Lagoon which is already nutrient-enriched to a degree.

#### ■ Ecosystem composition

The operation of the jetty, KFDA camps/parking lots/materials yards, airstrip, CPF, pipeline, in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure is not expected to result any noticeable alternation of the ecosystem composition of the aquatic communities.

As mentioned above, pollution and erosion and sedimentation derived from the Project's footprint could alter the composition of the communities if that pollution is toxic to aquatic life, or sediment loads smother aquatic organisms if mitigation measures fail. This notwithstanding (with the exception of catastrophic pollution events, which are addressed in Vol 1b) it is considered unlikely that operation activities would alter the ecosystem composition of the near-shore aquatic communities during the expected ~25-year operation period for the Project.

The potential for influx and in-migration of people onto the Buhuka Flats seeking opportunities from the Project, and because of the easier access provided by the escarpment road, could place increased pressure on the near-shore habitats. This increased pressure could arise from increased fishing activities, as well as additional sources of pollution (such as, fuels, human and livestock waste, fish waste, and litter). The resultant increase in the population on the Buhuka Flats could lead to a dramatic change in the current drivers of change to the near-shore aquatic habitats. These pressures could alter the ecosystem composition of these habitats.

Provided that the appropriate management measures are in place, in line with CNOOC's in-house alien invasive species management policy, increases in populations of invasive and exotic species are not expected to result directly from the operation of the project. The increased population of people on the Buhuka Flats could alter ecosystem processes and functions and lead to an increased susceptibility of the natural ecosystems to invasion by exotic species. However, certainly for the near-shore aquatic environment, this is not expected.

Impacts from the changes to ecosystem composition are predicted to be beyond the expected range of natural disturbance perturbations, if the mitigation measures fail or are inadequate.

#### ■ Ecosystem configuration

The operation of the Project is expected to last for 25 years. The operation of the jetty, KFDA camps/parking lots/materials yards, airstrip, CPF, pipeline, in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure is not expected to result any noticeable alteration of the ecosystem configuration of the aquatic ecosystems and habitats in the CHAA. It is expected that the connectivity amongst the aquatic habitats will remain the same during operation as they were during baseline.

The potential for influx and in-migration of people onto the Buhuka Flats seeking opportunities from the Project, and because of the easier access provided by the escarpment road, could place increased pressure on the near-shore habitats. This increased pressure could arise from increased fishing activities, as well as additional sources of pollution (such as, fuels, human and livestock waste, fish waste, and litter), and introduction/spread of invasive plant species. The resultant increase in the population on the Buhuka Flats could lead to a dramatic change in the current drivers of change to the near-shore aquatic habitats. These pressures could alter the ecosystem composition of these habitats.

Impacts from the changes to ecosystem configuration are predicted to be within the expected range of natural disturbance perturbations.





### Impact Classification

The near-shore aquatic habitat's sensitivity is high because these habitats potentially support populations of the Critically Endangered Mud Snail (*Gabbiella candida*), and the range restricted and Near Threatened Snail (*Bellamya rubicunda*).

Impacts to the representativeness of near-shore habitats will be neutral. The geographical extent of impacts will be still be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because the jetty and water intake and pump station will remain in place for the duration of the operation phase, and natural ecosystem processes (i.e. longshore sediment drift) are expected to become re-established in the vicinity of the water intake and pump station (assuming that the recommended construction-phase mitigation measures were successfully applied). Although increased human populations and activity (fishing activity – wastes, fuels, litter; increased cattle grazing) on the Buhuka Flats could contribute to increased pressure on Lake Albert, no changes to the representativeness of the near-shore habitats during operations are expected. The magnitude of operation phase impacts of on representativeness of the near-shore aquatic habitats is therefore considered negligible. Therefore, the negligible magnitude of impact and high sensitivity of the receptor combine to produce a minor overall impact level to representativeness during the operations phase of the Project, pre-mitigation (Table 17). Following the application of appropriate mitigation measures (Section 11.0), the impact significance is expected to remain the same, that is, minor, primarily because the magnitude will remain negligible, and the sensitivity of the receptor will remain the same.

Impacts to the topography (geomorphology) and sediment transport in the near-shore habitats during operation will be adverse. The geographical extent of impacts will be local, because impacts are restricted to the LSA. Impact duration will be medium-term (that is, ~25 years operation duration). The magnitude of the impact on topography (geomorphology) and sediment transport is expected to be low prior to mitigation, because wetland vegetation within the Kamansinig River, Masika River and River 1 are expected to provide efficient attenuation of sediment, and a significant increase in sediment concentrations in the lake or in Bugoma Lagoon as a result of Project operations are unlikely. However, sediment loads from overland (stormwater) flows may not be retarded by vegetation (particularly in a scenario where human-induced pressures such as cattle overgrazing and land deterioration occurs over the lifetime of the Project), and hence may report to the near-shore habitats, contributing to measurable increased turbidity during and after storms, where River 1 discharges to the Lake; potentially resulting in impacts of low magnitude. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact significance to topography (geomorphology), sediment, and water quality during the operational phase of the Project, pre-mitigation (Table 17). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

Impacts to the water quality as a result of discharge of treated sewerage effluent, and stormwater that could contain oil and chemical pollution will be adverse. The geographical extent of impacts will be local, because impacts are restricted to the LSA, in the immediate vicinity of the discharge points to the Lake. Impact duration will be medium-term (that is, ~25 years). The baseline concentration of hydrocarbons and other pollutants in the lake water is currently below levels that could cause harm in the lake environment (ref. Vol 4a Surface Water). The magnitude of the potential impacts on water quality varies; with high magnitude assigned to any impacts associated with stormwater contaminated with oil and potentially-hazardous/toxic chemicals, and medium magnitude impacts predicted for potential discharge of treated sewerage. Combined with the high sensitivity of the near-shore habitats, both impacts on water quality could be of major significance, prior to mitigation (Table 17). Following the application of appropriate mitigation measures, the impact significance associated with contaminated stormwater reaching the Lake is expected to be reduced to moderate, primarily because the magnitude will become low, and the sensitivity of the receptor will remain the same. Consideration of other options for final disposal of treated sewerage effluent, such as irrigation onto land in the buffer area around the CPF and at the personnel camp lawns and gardens, or discharge into an artificial wetland or sustainably managed plantation (ref. Section 11.0), could further reduce the magnitude of effects on water quality associated with disposal of treated sewerage effluence to negligible, resulting in overall impacts of minor significance on water quality.

Impacts to the water quality as a result of increased numbers of people and livestock on the Buhuka Flats are likely, and will be adverse. The geographical extent of impacts will be local, because impacts will be restricted





to the LSA. Impact duration will be medium-term (that is, ~25 years). In the context of existing nutrient enrichment in the Bugoma Lagoon, the magnitude of additional nutrient input is predicted to be medium, which combined with the high sensitivity of the habitat, results in an overall impact of major significance, prior to mitigation. Following the application of appropriate mitigation measures, the impact significance could be reduced to moderate, primarily because the magnitude will become low, but the sensitivity of the receptor will remain the same (Table 17).

Impacts to the ecosystem composition will be adverse. The geographical extent of any impacts will be local because impacts are restricted to the LSA. Impact duration will be long-term (that is, ~25 years). The magnitude of the impact on ecosystem composition is low because, prior to any mitigation, the potential for changes to ecosystem composition potentially brought about by increased fishing intensity, pollution and smothering is possible. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the operational phase of the Project, pre-mitigation (Table 17). Following the application of appropriate mitigation measures, the impact significance is expected to remain minor, because the magnitude will remain negligible, and the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be neutral. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be long-term (that is, ~25 years). The magnitude of the impact on ecosystem configuration is negligible because, prior to any mitigation, the potential for changes to ecosystem configuration is very remote. Consequently, even post-mitigation, the significance of the impact will remain minor (Table 17).

**Table 17: Potential impacts in the operation phase to near-shore habitats**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance
Representativeness	Negligible – 1	High - 4	Minor – 4	Negligible – 1	High - 4	Minor – 4
Topography (geomorphology) and sediment	Low – 2	High - 4	Moderate – 8	Negligible – 1	High - 4	Minor – 4
Water quality – sewerage effluent	Medium - 3	High - 4	Major - 12	Negligible – 1	High – 4	Minor - 4
Water quality – contaminated stormwater	High - 4	High - 4	Major - 16	Low – 2	High - 4	Moderate – 8
Water quality – increased population and livestock	Medium - 3	High - 4	Major - 12	Low – 2	High - 4	Moderate – 8
Ecosystem composition	Low – 2	High - 4	Moderate – 8	Negligible – 1	High - 4	Minor – 4
Ecosystem configuration	Negligible – 1	High - 4	Minor – 4	Negligible – 1	High - 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios with the level of Lake Albert, which has, in the recent and not so recent past varied quite dramatically (Talbot et al. 2006), it is conceivable that level of the lake may increase or decrease thereby altering near-shore habitats.





The extent of influx and in-migration of people onto the Buhuka Flats specifically seeking opportunities is uncertain. Indications are that additional large numbers of people have moved onto the flats, or are present daily in large numbers, primarily because access is now easier (insert ref to updated 2017 social study). This increase in population is expected to exacerbate existing anthropogenic pressures on the Buhuka Flats and the near-shore habitats of Lake Albert.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

### **10.2.1.2 What impact could the operation of the Project have on the wetlands in the Buhuka Flats region of the CHAA?**

#### **Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the wetlands of the CHAA were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

Indirect affects to habitat were estimated by applying a 0.50 km buffer to the Project infrastructure. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge impacts, fragmentation, sensory disturbance, changes in water quantity and quality, air emissions and dust, and population increases.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur, and the disruption of known corridors. This was accomplished by examining available scientific literature about the ecology of wetlands (permanent and seasonally flooded grasslands).

#### **Impact Assessment**

##### ■ Representativeness

Operation of the Project infrastructure is unlikely to cause ongoing loss of representativeness of wetlands in the CHAA, or ongoing impacts to the permanent wetlands and the seasonally flooded grasslands of the CHAA, assuming the mitigation measures devised for the construction phase are impactful.

Impacts to representativeness of wetlands during the operation of the Project are predicted to be within the expected range of natural disturbance perturbations.

##### ■ Ecosystem composition

The operation of the Project infrastructure is unlikely to cause changes to the composition of wetlands in the CHAA, or ongoing impacts to the permanent wetlands and the seasonally flooded grasslands of the CHAA, assuming the mitigation measures devised for the construction phase are impactful. Similarly, during the operation phase, noise and sensory disturbances created by the equipment is not expected to alter the behaviour of species frequenting the wetlands. For example, wading birds and Grey Crowned Cranes could become accustomed to the ongoing operational noises.

It is assumed, as part of standard operational methods, that appropriate drainage-line crossings will be maintained as part of the in-field road and airstrip management. Nevertheless, even with such measures in place, there is still a potential for long-term changes to the wetland character to occur. For example, there is the potential for erosion downstream of the crossings, backwater upstream of the crossings, and erosion at the entrance to the crossing structures. The airstrip is one area in particular where construction across a drainage line might lead to decreased flows and erosion downstream of the airstrip in the long-term. This could lead to changes in the ecosystem functions and processes in the downstream wetlands, if not maintained during the operational phase.

The operation of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, in-field roads, crusher plant/spoil area A, well pads, and associated infrastructure could cause increased erosion and sediment-laden run-off to report to the wetlands surrounding the Project footprint. The soils of the Buhuka Flats are dispersive





(Vol 4a), and active soil erosion is evident in the LSA. There is a potential that, without adequate erosion and sediment control measures in place during operations, sediment loads within the watercourses draining the Project footprint could increase. Hence, there is a potential for increased sediment loads in the wetland habitats. The watercourses and associated wetlands draining the LSA support dense emergent vegetation (see Section 6.1.1). Such vegetation can form an impactful filter for most sediment (IECA 2008); therefore, it can be expected that sediment loads reporting to downstream wetland habitats could be minimal. Nevertheless, if sediment loads are substantial, there is a potential for that sediment to smother wetland vegetation and interfere with aquatic invertebrates. If this occurs, it could detrimentally affect the wetland processes and functions, which, in turn, could alter wetland composition, albeit on a localised scale.

It is assumed that the potential acid sulphate soils in the permanent wetlands in the CHAA were adequately managed during construction, and no lasting impacts occur.

All of the above direct impacts to ecosystem composition of the wetlands in the CHAA are predicted to be within the expected range of natural disturbance perturbations during operations.

However, indirect impacts on wetlands will occur as a result of the migration of people into the CHAA both as a result of easier access provided by the operation of the newly constructed escarpment road, and in search of work opportunities, and associated increase in head of livestock grazing in the Buhuka Flats. Increased stocking is likely to result in greater pressure on vegetation communities supplying wood and fibre resources, overgrazing, and expansion of subsistence crops, causing greater runoff, channel incision, heightened fire risk and the loss of wetland function over large areas. In the dispersive soils of the Buhuka Flats, the risk of catastrophic soil loss and subsequent effects on wetland ecosystem composition will be high.

Indirect effects on the ecosystem composition of wetland communities are also likely, as a result of ongoing discharge of treated sewage effluent and stormwater in excess of the first 15 minutes of rainfall, to the receiving surface water systems (primarily River 1 and the Kamansinig River). Long-term discharges of this nature are likely to affect the vegetation species composition as a result of nutrient enrichment from the sewerage discharge, as well as the fluctuations in wetness regimes brought about by both the sewerage discharge and occasional (seasonal) stormwater peak flow inputs.

In addition, minor spillages of fuels and chemicals during the day-to-day operation of the Project components (other than catastrophic spillages, which are addressed in see Vol 1b, 2 & 3 ESMPs) could end up in the wetlands of the LSA; potentially affecting some invertebrate species (for example, aquatic snails (Araujo et al. 2012), mayflies (Savić et al. 2011)) and juvenile fish (for example, Agamy 2013) which are highly sensitive to chemical pollutants, particularly hydrocarbons. Currently, the concentrations of hydrocarbons and other pollutants in the water of the Bugoma Lagoon are below levels that could cause harm to the environment (Vol 4a); this is supported by the healthy aquatic communities observed in the lagoon's and Masika River wetland habitats (see Section 5.2). The entrainment of small quantities of oil and potentially hazardous chemicals in stormwater runoff from the Project infrastructure and subsequent entry to wetlands (particularly the seasonally flooded grassland adjacent to well pad 1) is considered possible as the proximity of well pad 1 to the wetlands provides little buffer for potential contamination.

### ■ Ecosystem configuration

The long-term impacts of the construction of roads, the airstrip and the flowlines through the wetlands of the CHAA are unknown. Although the roads could potentially act as barriers to certain wetland species, they are only expected to be 5 m wide and unsealed, and, during operations, they will convey reduced Project-related traffic volumes; however, they are likely to be preferentially used by members of the increased local communities of the Buhuka Flats. Therefore, they could become barriers to species movement, particularly for species such as Grey Crowned Crane, should that species become accustomed to the increased human disturbance brought about by the population influx to the Buhuka Flats during operation.

The operation and management of the in-field roads and airstrip should not affect ecosystem processes driving the wetlands, assuming the installation of appropriate drainage connections was successful during construction. What the long-term impact of the edge impacts, and fragmentation of the wetland habitats, in the CHAA caused by the construction of roads, is unknown. As already discussed, vegetation clearing creates edges or boundaries where habitat meets a disturbance. These edges open up habitat in areas where it was





previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges also often result in changes in species composition along the edge, with the edges typically becoming dominated by pioneer and weedy species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Fragmentation of wetland habitat is known to reduce the viability of many species and the wetland as a whole, with the viability of the particular fragment dependent on its size, proximity and, hence, connectivity to other wetland habitats (Uzarski et al. 2009).

What long-term impacts that the operation these roads and the airstrip could have on the wetland communities' configuration are unknown; in particular their resilience. What is known is that these wetland communities are already under pressure from livestock grazing, and harvesting of fibre for house construction.

The operation of the KFDA camps/parking lots/materials yards, CPF, crusher plant/spoil area A, well pads, and associated infrastructure will not directly lead to changes in the ecosystem configuration of wetlands in the CHAA.

The long-term impacts to ecosystem configuration of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations during operations, if mitigation measures are inadequate or fail.

### **Impact Classification**

Wetland sensitivity is high because these habitats, particularly the permanent wetlands, are potential breeding habitat for Grey Crowned Cranes (a species of concern). The wetlands are also already under stress from fires, livestock grazing and harvesting of fibre.

Impacts to the representativeness of the habitat may be adverse during operations. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because the roads (which may affect the hydrological character of the wetlands, and subsequently wetland extent and condition) will be permanent features, and will remain in place even after the project is decommissioned. The magnitude of the impacts of operation of the Project on representativeness of the wetland communities is low. Therefore, the magnitude and high sensitivity of the wetlands combine to produce a moderate overall impact level to representativeness during the operations phase of the Project, prior to the implementation of site-specific mitigation measures (Section 11.2). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced minor because the magnitude will be negligible, while the sensitivity of the receptor will remain the same.

Impacts to the ecosystem composition as a result of the operation of Project infrastructure, and presence of wetlands crossings, will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because impacts are expected last as long as the in-field roads and airstrip are in place, and the potential changes to ecosystem composition are unknown. The magnitude of the impact on ecosystem composition is low because, prior to any mitigation, the potential for changes to ecosystem composition potentially brought about by edge impacts, changed flow regimes, and sedimentation, is possible. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the operation phase of the Project, pre-mitigation (Table 18). Following the application of appropriate mitigation measures, the impact significance is expected to become minor because the magnitude will become negligible, yet the sensitivity of the receptor will remain the same.

Impacts to the ecosystem composition as a result of the increased human population living on the flats, and the associated increase in head of livestock grazing in the Buhuka Flats, will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because impacts are expected to be permanent, and the potential changes to ecosystem composition are unknown. The magnitude of the impact on ecosystem composition is medium because, prior to mitigation, changes to ecosystem composition potentially brought about by overgrazing, cattle trampling and subsequent effects on wetlands soils could be near the limits of wetland capacity to adapt. Therefore, the magnitude and sensitivity combine to produce a major overall impact level to ecosystem composition during the operation phase of the Project, pre-mitigation (Table 18). Following the application of



appropriate mitigation measures, the impact significance is expected to become moderate because the magnitude will become low, yet the sensitivity of the receptor will remain the same.

The entrainment of small quantities of oil and potentially hazardous chemicals in stormwater runoff from the Project infrastructure and subsequent entry to wetlands (particularly the seasonally flooded grassland adjacent to well pad 1) will result in adverse impacts of local extent, that could occur intermittently throughout the lifetime of the Project. The magnitude of the potential impacts on wetland water quality is high, as pollution events could significantly alter aquatic species communities, and therefore ecosystem composition. Combined with the high sensitivity of the wetland habitats, impacts on wetland water quality could be of major significance prior to mitigation (Table 18). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will extend into the far future (that is, ~25 years) because impacts are expected to continue as long as the in-field roads, flowlines and airstrip are in place. The magnitude of the impact on ecosystem configuration is low because, prior to any mitigation, the potential for changes to ecosystem configuration is possible, especially from fragmentation. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem configuration during the operation phase of the Project, pre-mitigation (Table 18). Following the application of appropriate mitigation measures, the impact significance is expected to become minor because the magnitude will become negligible, yet the sensitivity of the receptor will remain the same.

**Table 18: Potential impacts in the operation phase to the wetlands of the CHAA**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Ecosystem composition – project infrastructure, wetland crossings	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Ecosystem composition – increased population	Medium - 3	High – 4	Major – 12	Low - 2	High – 4	Moderate – 8
Ecosystem composition – contaminated stormwater	High – 4	High – 4	Major - 16	Negligible – 1	High – 4	Minor – 4
Ecosystem configuration	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the wetlands of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain.





The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

### 10.2.1.3 *What impact could the operation of the Project have on the escarpment vegetation corridors?*

#### **Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the vegetation corridors on the escarpment were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

It is recognised that the escarpment road has already been permitted and is currently operational, and was covered by a separate ESIA process, see AWE (2014c); however, the inclusion of this road in this assessment is important because of its induced (and cumulative – Chapter 17 CIA) impacts on the vegetation of the escarpment. Literature was reviewed to understand the long-term impacts of roads through natural habitats, during operation.

Additional, indirect effects to habitat were estimated by applying a 0.50 km buffer to the temporary camp and quarry at the top of the escarpment, and the footprint of the escarpment road. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge impacts, fragmentation, sensory disturbance, changes in water quantity and quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur over the life of the Project, and the disruption of known corridors. This was accomplished by examining available literature about the ecology of the vegetation communities on the escarpment, and scientific literature about the impacts of human activities on corridors in the long-term.

#### **Impact Assessment**

##### ■ Representativeness

The CHAA supports approximately 2443 ha of vegetation communities on the escarpment (see Section 7.1.2), which are bounded on the east by agriculturally modified landscapes, and the Buhuka Flats on the west. As mentioned, these vegetation communities form part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2003). Therefore, their continuous extent represents an important landscape feature in the CHAA.

The initial loss of habitat from the construction of the temporary camp on top of the escarpment is expected to be reverted during operation, because the camp site will have been rehabilitated. Rehabilitation measures should return the site to a vegetation cover similar to that that was there before the camp was constructed, that is, open wooded bushland. The loss of approximately 12.8 ha of vegetation communities, because of the construction of the escarpment road, will persist during the operational phase.

The operation of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, pipeline, new in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure will not directly lead to the loss of vegetation communities on the escarpment.

Impacts from the changes to representativeness are predicted to be the less than those experienced during the construction phase because temporary construction areas should have been rehabilitated. Nevertheless, the loss of vegetation to the escarpment road footprint will remain for operation, representing a change in representativeness is predicted to be beyond the expected range of natural disturbance perturbations.

##### ■ Ecosystem composition

What the long-term impact of the escarpment road on the ecosystem composition of the escarpment in the CHAA is unknown. The long-term consequences of a linear corridor through an area of relatively intact vegetation, like that on the escarpment, create edge impacts that could, in the long-term, alter the composition





of the ecosystem through which the road traverses (Findlay and Bourdages 2000). Edges associated with roads tend to be abrupt, with a high degree of contrast between the two areas (for example, road and open wooded grassland). The edges open up habitat in areas where it was previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges often result in changes in plant species composition along the edge, with the edges typically becoming dominated by pioneer species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Often, these microclimates are favoured by weed species. However, certainly for fauna species, and depending on the species, edges can have either a positive or negative impact on habitat quality and quantity (Prevedello et al. 2013, Wellicome et al. 2014).

Apart from the possible changes brought about by the edge impacts, the road may also: restrict movement of certain less mobile faunal species between populations; increase mortality of individuals due to collision with vehicles; fragment habitat; present a possible path for invasive species to enter the area; or increase human and livestock access to otherwise less accessible habitats (Findlay and Bourdages 2000). All these predicted impacts, combined, could be expected to increase the potential for local extinction rates or decrease local recolonisation rates.

The escarpment road is sealed for its length down the escarpment. Sealing of the road presents other aspects that may affect the ecosystem composition of the communities on either side of the road in the long-term. For example, concentrating water run-off from the sealed surface, which could carry contaminants such as fuel, heavy metals and polycyclic-aromatic hydrocarbons, to watercourses not otherwise influenced by such run-off, and possibly lead to changes in water quality and erosion regimes. Polluted run-off could lead to detrimental impacts on aquatic species downstream from the escarpment road.

What long-term effects that these edge impacts could have on the vegetation communities' composition throughout the lifetime of the road operation are unknown; in particular their resilience<sup>4</sup>. The most significant project-related impact on the ecosystem composition of the escarpment is likely to be the indirect effect of migration into the area, facilitated by the escarpment road. Without mitigation, it is likely that both the project and the new ease of access to the flats will cause significant additional in-migration (refer to Vol 4c Social Assessment for detail), with people settling both on the Flats and in the villages above the escarpment. The escarpment vegetation communities are already under pressure from harvesting of fuel wood and charcoal manufacture, and livestock grazing. Cattle have been shown to substantially increase the edge impacts in savannah habitats (Porensky et al. 2013). It is conceivable that the road could afford people, cattle and other livestock easier access to resources and grazing on the escarpment, and facilitate the spread of weed and invasive plant species, which could place increased pressure on these communities in the long-term. It is likely that these communities may change in the long-term as grazing pressure increases, and large trees are removed for charcoal manufacture. These changes could substantially alter the habitat structure and composition, which, in turn, could affect its utilisation by the current species guild. To some extent, this may already be occurring. For example, very few medium sized mammals were recorded in the CHAA, and those that were recorded tended to be thicket and dense bushland specialists, such as bushbuck and duiker. The low populations and diversity of these species could also be a reflection of increased pressure for bush meat from the local human population, which has increased markedly over the last ten years (AECOM 2012).

The operation of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, pipeline, new in-field roads, crusher plant/spoil area A, new well pads, and associated infrastructure will not directly lead to changes in the composition of vegetation communities on the escarpment.

Impacts from the long-term changes to ecosystem composition are predicted to be well beyond the expected range of natural disturbance perturbations if mitigation measures fail or are inadequate.

### ■ Ecosystem configuration

Roads, and especially sealed roads are known to be significant barriers, or alter behaviours, of a range of wildlife, from: small ground-dwelling mammals, insects, reptiles and amphibians (for example, Brehme et al.

<sup>4</sup> For this study, indirect effects, such as increased harvesting pressures due to migration into the area, either because of better access to existing resources or because of opportunities provided by the oil industry, are considered to be operational impacts which are driven by the access provided by the road





2013, Pontoppidan et al. 2013, Rotholz and Mandelik 2013); to bats (for example, Berthinussen and Altringham 2012); to birds (for example, Kociolek et al. 2011); to primates (for example, Mammides et al. 2009); to large ungulates (for example, Leblond et al. 2013, Meisingset et al. 2013). Depending on the species, the presence of roads may affect individuals in many direct and indirect ways. For example, roads may inhibit seasonal migration and may cause an impactful loss of habitat due to avoidance.

The wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor is recognised as being important for threatened species in the face of climate change adaptation (Ayebare et al. 2013), and as part of a much broader set of corridors running the length of the Albertine Rift. This corridor in the southern portion of the eastern shores of Lake Albert, in the vicinity of the Project, is very narrow compared to its extent elsewhere, and is recognised as being important for savannah species (Plumptre et al. 2010). The escarpment road to the Buhuka Flats is the only major road on the south-eastern portion of Lake Albert from the southern end of the lake to the Kabwoya Wildlife Reserve, that is, the only major road within the identified wildlife corridor. This area is otherwise devoid of notable roads and tracks, adding to its value as a wildlife movement corridor.

Indications are that populations of highly mobile wildlife within the area, which may depend on the corridor, are not substantial, and potentially severely depleted (see Section 7.1.2). In particular, most of those species that could utilise this corridor, such as large ungulates, predators, and primates are very rare in the escarpment area surrounding the Buhuka Flats, with the exception of birds. However, the road is not expected to be a barrier for birds; for example, White-backed Vultures were seen moving south down the escarpment corridor during the surveys (see Section 6.1.4). What impact the road could have on the movement on those terrestrial individuals and species that may utilise this corridor in the long-term is unknown. The corridor is recognised as an important climate change refugium for a range of threatened species, which may become increasingly important for those species in the future (Ayebare et al. 2013), that is, within the life time of the Project.

Impacts from the changes to ecosystem configuration on the escarpment as a result of the long-term operation of the road are predicted to be beyond the expected range of natural disturbance perturbations, if mitigation measures fail or are inadequate.

### **Impact Classification**

The vegetation community of the escarpment's sensitivity is medium because, although the integrity of these habitats is already under stress from livestock grazing and harvesting of fuel wood and non-timber forest products, they do form part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor.

During operation, the impacts to the representativeness of the habitat will be neutral, because areas used temporarily during construction will be rehabilitated and the vegetation on those areas will be establishing. The geographical extent of impacts will be local because impacts are still restricted to the LSA, and the loss of vegetation communities because of the road's construction will persist. Impact duration will be into the far future (that is, ~25 years) because the road down the escarpment will remain in place even after the project is decommissioned. The magnitude of the impacts of operation on representativeness of the vegetation communities of the escarpment is medium. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to representativeness during the operational phase of the Project, pre-mitigation (Table 19). Following the application of appropriate mitigation measures during operation, including the assumption that offset mechanisms are in place and working, the impact significance is expected to remain moderate, although the magnitude will be low, primarily because it is assumed that offsets will be achievable, while the sensitivity of the receptor will remain the same.

Impacts to the ecosystem composition will be adverse in the long-term. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be into the far future (that is, ~25 years) because impacts are expected last as long as the road is in place. The magnitude of the impact on ecosystem composition is medium because, prior to any mitigation, the potential for changes to ecosystem composition potentially brought about by edge impacts, and the consequences of easier access to the adjacent escarpment areas for livestock grazing and natural resource harvest, is possible. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the operational phase of the Project, pre-mitigation (Table 19). Following the application of appropriate mitigation







measures, including the assumption that offset mechanisms for the losses suffered during construction are in place and working, the impact significance is expected to remain moderate, although the magnitude will become low, yet the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be regional because impacts are restricted to the CHAA. Impact duration will extend into the far future (that is, ~25 years) because impacts are expected to continue as long as the road is in place. The magnitude of the impact on ecosystem configuration is medium because, prior to any mitigation, the potential for changes to ecosystem configuration is possible, especially inference with wildlife movement corridors. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the operational phase of the Project, pre-mitigation (Table 19).

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate; although the magnitude will become low, the sensitivity of the receptor will remain the same.

**Table 19: Potential impacts in the operation phase to the vegetation communities of the escarpment**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6
Ecosystem composition	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6
Ecosystem configuration	Medium – 3	Medium – 3	Moderate – 9	Low – 2	Medium – 3	Moderate – 6

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the vegetation communities of the escarpment, and the wildlife corridors of which they form part. However, there is some uncertainty with regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain. Indications are that they will increase in importance (Ayebare et al. 2013), provided human pressures associated with the presence of the escarpment road do not overwhelm them.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**10.2.1.4 What impact could the operation of the Project have on the Bugoma Central Forest Reserve?**

**Impact Analysis – Methods**

As discussed in Section 6.1.6, it is recognised that CNOOC will not be responsible for the management of the R5 and P1 Roads during the operation of the Project. It is expected that the Ugandan Government will be responsible for that management and upkeep. Nevertheless, as discussed, the road will have been upgraded during the construction phase, and will remain in place throughout the operational phase and beyond. Therefore, any effects associated with, and stemming from the existence of the upgraded road can be seen to be induced impacts arising as a result of the Project’s development.

A formal impact classification based on indicators was developed for induced and cumulative impacts to the Bugoma Central Forest Reserve; the impacts are discussed, and their significance assessed through a





**reasoned narrative.** An overall impact significance classification is then developed. This was accomplished by examining available literature about the ecology of the Bugoma Central Forest Reserve, and scientific literature regarding the effects of migration and human population pressure on forests in Africa.

The impacts were assessed in light of the guidance provided by IFC (2013), and in consideration of other known projects being developed in the wider area. In particular, the development of the oil processing facility at Kabaale, the oil developments around the Kaiso-Tonya area (AECOM 2012), the Hoima-Mputa-Fort Portal-Nkenda power line, and the potential for regional population increases in the wider area.

### **Impact Analysis Results**

As discussed, the Bugoma Central Forest Reserve is identified as a valued component for this impact assessment, in terms of biodiversity (see Section 6.3.1.1.4). As identified in that section, apart from being one of the last stands of tropical semi-deciduous forests in the region, it also supports known populations of the range-restricted Nahan's Francolin and Endangered Eastern Chimpanzee (Plumptre et al. 2011), potential non-breeding habitat for the Endangered Madagascar Pond Heron (see Section 6.3.3.1), as well as elephants and a host of other threatened and irreplaceable species.

The Bugoma Central Forest Reserve is under severe pressure from the human population surrounding it. Only recently were more than 1,500 people evicted illegally after settling within the forest (Mugerwa 2013). The land cover study (see Appendix D and Figure 14) clearly shows that the areas surrounding the forest have largely been transformed for agricultural and subsistence purposes. This trend of encroachment of protected areas, like the Bugoma Central Forest Reserve, is not unique. Protected areas are known to be particularly vulnerable to changes in human demographics and deforestation; and Wittemyer et al. (2008) identified that human population growth and encroachment around protected areas is significantly higher than the average population growth in rural areas. This difference was largely due to the immigration of people into these areas because of the perceived increased availability of opportunities, natural resources and potential jobs (Wittemyer et al. 2008). This is supported by research conducted in the forests of the Albertine Graben, and the wider CHAA. For example, in a study of Budongo Central Forest Reserve, Zommers and MacDonald (2012), identified that of the local communities that hunted bush meat in the forest, nearly 73% were immigrants to the area. Furthermore, these workers identified that the households of immigrants were also more likely to be involved with deforestation.

The upgrade and improvement of roads in rural areas can influence immigration rates into those areas (Wennergren and Whitaker 1976, Godar et al. 2012). It is highly likely that the upgraded R5 and P1 roads could induce population influx into the Bugoma Central Forest Reserve, as well as areas adjacent to the road itself throughout its operational lifetime, because access would be made easier. Furthermore, the proximity of the Bugoma Central Forest Reserve in the area could also make the area more attractive to immigrants seeking opportunities (*viz.*, Wittemyer et al. 2008). The current influx of refugees into the CHAA is also not expected to dissipate with the continued instability in the DRC (UNHCR 2014). Therefore, there is a potential that the upgrade of the road could place increased pressure on the Bugoma Central Forest Reserve, the natural resources it offers (such as, timber, non-timber forest products), and bush meat.

An Influx Management Strategy and Framework Plan has been developed to manage the potential influx of people into the LSA. However, this plan can only focus on those measures over which CNOOC has control, and to support the range of government and donor projects in Uganda aimed at socio-economic development and environmental conservation. How this translates to the potential influx of people along the improved R5 and P1 roads is unknown.

The improved road could also allow for an increase in vehicular traffic into the area, and faster speeds associated with those vehicles. Therefore, there is also the potential that direct mortality of wildlife along the road could increase. The Bugoma Central Forest Reserve is home to populations of threatened species and an array of other species (see Plumptre et al. 2010, 2011). Many of these species will move within the forest and between sections of the forest. Although the negative effects of roads on wildlife in tropical rainforests, like Bugoma Central Forest Reserve, are poorly understood, indications are that: (1) many species avoid roads altogether (especially, medium-sized mammals, diurnal, solitary and group living animals, and ungulates); and (2) high vegetation cover on the road verges (see Figure 20) increases crossing probability substantially (van



der Hoeven et al. 2010). Currently, the roadside vegetation on the P1 road would encourage wildlife to cross (Figure 22); presumably this is also the case for the R5 road. This could place them in the direct paths of traffic during the operational phase of the Project.

**Impact Classification**

Impacts from the presence of the R5 and P1 roads, and the resultant increased traffic and ease of forest access along that road during the operation of the Project will be adverse. The geographical extent of impacts will be regional because effects are restricted to the R5 and P1 Road corridor in the CHAA. Impact duration will be long-term (extending for the lifetime of the Project, and beyond, as the roads are government-managed and will remain in place). The magnitude of the effects of operation on the Bugoma Central Forest Reserve is low.

The sensitivity of the Bugoma Central Forest Reserve is high because it is a threatened ecosystem that is already under pressure. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Project, pre-mitigation (Table 20).

The mitigation hierarchy is an important process that has been used to minimise impacts to the Bugoma Forest Reserve. The focus for the continued use of the mitigation hierarchy during the road upgrade and construction will be continued development and implementation of mitigation measures through monitoring and adaptive management; including suggested measures for the Ugandan Government to apply in the management of the upgrade of the P1 Road, and recommendations regarding the intended use of the R5 road running north to south through Bugoma CFR (see Section 11.0).

Following the application of appropriate mitigation measures, including management support of the forest to limit influx of people, the impact significance is expected to remain major, although the intensity could become medium, yet the sensitivity of the receptor will remain the same (Table 20).

**Table 20: Potential impacts in the operation phase to the Bugoma Central Forest Reserve**

	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance
Habitat and ecosystem integrity	High – 4	High – 4	Major – 16	Medium – 3	High – 4	Major – 12

As noted above, the focus for reducing impacts to the Bugoma Central Forest Reserve is mitigation to lessen various types of disturbance that may occur (Section 12.0). Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to the forest have been identified and are discussed in Section 13.0.

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential effects that the operation of the upgraded R5 and P1 roads, could have on the Bugoma Central Forest Reserve.

The spatial extent of the forest reserve in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**10.2.2 What impact could the operation of the Project have on species of concern?**

This section presents the assessment of the impacts that the operation of the Project could have only on those species of concern that trigger critical habitat, as identified in Section 9.2, and other species of concern that





were included as valued components for the impact assessment (Table 5). Potential impacts to other species of concern are assessed at the habitat level (see Section 3.3.2).

### 10.2.2.1 What impact could the operation of the Project have on the Mud Snail?

As previously discussed, the Mud Snail (*Gabbiella candida*) is a Critically Endangered and range restricted species, which could occur in the CHAA.

#### Impact Indicators

Indicators used to assess impacts of the operation of the Project on the Mud Snail were: habitat quantity and quality, and habitat connectivity.

Habitat loss due to indirect disturbance was assessed by calculating the area of disturbance of suitable habitat from the CHAA as a result of the operation of the Project. Changes to habitat quality were assessed by the prediction of sediment loads and changes to water quality in the water column from operational activities.

Changes in habitat connectivity were assessed by identifying potential barriers to genetic movement, and source populations. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the Project footprint to qualitatively identify areas where critical habitat becomes fragmented.

#### Impact Assessment

##### ■ Habitat Quality and Quantity

Very little information is available for this species. The two Mud Snail species recorded from the CHAA were collected from bottom substrates in the open sandy shore habitats. These habitats are characterised by a gently sloping lake bed extending from the shore line to deeper water.

The operation of the Project, and, in particular, the jetty, could affect the open sandy shoreline habitat within the vicinity of this infrastructure. However, the operation of the jetty itself is unlikely to change the physical structure of the open sandy shoreline habitat. Rather, the wash created by berthing and departing barges could affect the habitat quality through the disturbance of bed sediments. It is expected that these increased sediment loads will dissipate reasonably quickly. The sediment loads in the vicinity of the jetty are not expected to exceed those that would normally be expected during windy periods on the lake, and the consequent turbid conditions caused by those winds. In addition, with the opening of the escarpment road, the use of the jetty is expected to be minimised.

Accidental spillages of small quantities of fuels and chemicals during the operation of the Project components, such as loading and unloading barges, or refuelling vehicles, and drilling fluids, could end up directly in the lake, or in a watercourse, which, ultimately, report to the near-shore habitats of the CHAA. There is a real potential for this to occur as part of the jetty operation, and as a component of stormwater discharge beyond the first 15 minute rainfall events. The jetty and stormwater discharge points are located right on the lake shore, and the impact of an accidental spill would be more obvious than a land-based spill; it would also be harder to contain and clean up. Aquatic snails like *G. candida* are highly sensitive to chemical pollutants, particularly hydrocarbons (Araujo et al. 2012). Currently, the concentrations of hydrocarbons and other pollutants in the lake waters of the near-shore habitats are below levels that could cause harm to the environment; this is supported by the healthy aquatic communities observed in the lake's near-shore habitats (see Section 6.2). As such, without adequate spill prevention and clean-up measures in place during operation, a chemical spill could, depending on the volume spilt, have a detrimental impact on the near-shore habitats of the CHAA through the introduction of toxic compounds and pollutants. Such a spill could have a detrimental impact on the Mud Snail *G. candida*.

The operation of the KFDA camps/parking lots/materials yards, airstrip extension, CPF, in-field roads, crusher plant/spoil area A, well pads, and associated infrastructure could cause increased sedimentation of near-shore habitats on the Buhuka Flats. The soils of the Buhuka Flats are dispersive, and active soil erosion is evident in the LSA. With the operation of the Project's components, there is a potential that, without adequate and maintained erosion and sediment control measures in place, sediment loads within the watercourses draining the Project footprint could increase. These sediment-laden watercourses report to Lake Albert (particularly River 1, which will receive stormwater runoff from the CPF and the airstrip; and the Kamansing, which will





receive stormwater runoff from the escarpment road, part of the airstrip and Well Pad 1), and, hence, there is a potential for increased sediment loads in near-shore habitats in the vicinity of the drainage discharge points south of Well Pad 2 and north of Bugoma Lagoon. Near-shore habitats particularly at risk include the Bugoma Lagoon, and to a lesser extent, the shallow river-associated habitats of the Kamansinig and River 1, both potential habitat for the Mud Snail. Nevertheless, the watercourses draining the CHAA support dense emergent vegetation (see Section 6.1.1.1). Such vegetation forms an impactful filter for most sediment (IECA 2008), therefore, it can be expected that sediment loads reporting to the near-shore habitats, at least, via the Kamansinig River, River 1 and Masika River, could be minimal. Sediment loads from overland flows may not be retarded by vegetation, and hence may report to the near-shore habitats and affect them detrimentally.

The potential for influx and in-migration of people onto the Buhuka Flats seeking opportunities from the Project, and because of the easier access provided by the escarpment road, could place increased pressure on the near-shore habitats. This increased pressure could arise from increased pollution (such as, fuels, human and livestock waste, fish waste, and litter), as well as physical disturbance from drinking cattle. The resultant increase in pollution levels could lead to a dramatic change in the current drivers of change to the near-shore aquatic habitats, and, ultimately affect the Mud Snail.

Impacts from the alteration quality of, or loss of, habitat are predicted to be beyond the expected range of natural disturbance perturbations.

■ **Habitat Connectivity**

The operation of the Project is expected to last for ~25 years. Besides the operation of the jetty, it is unlikely that operational activities could substantially alter the habitat connectivity of the near-shore habitats in the CHAA. No structures are being put in place that will alter the natural connectivity of the aquatic habitats of the lake. It is expected that the connectivity amongst the aquatic habitats will remain the same during operations as they were during baseline.

**Impact Classification**

The Mud Snail's sensitivity is high because this species is Critically Endangered.

Impacts to the Mud Snail's habitat quantity and quality could be adverse during operations. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be long-term (that is, 25 years). The magnitude of the impacts during operation - particularly long-term effects of trampling by drinking cattle and changes influenced by the jetty - on the habitat quantity and quality of the Mud Snail is predicted to be moderate. Therefore, the moderate magnitude and high sensitivity combine to produce a major overall impact level during the operation phase of the Project, pre-mitigation (Table 24). Following the application of appropriate mitigation measures, the impact significance is may be reduced to moderate, primarily because the magnitude of habitat disturbance arising from human and livestock will at best be reduced to low during operation, and the sensitivity of the receptor will remain the same (Table 21).

Impacts to the Mud Snail's habitat connectivity will be neutral. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be long-term, and no barrier to habitat connectivity will be constructed. The magnitude of the impacts of operation on the habitat connectivity of the Mud Snail is negligible, therefore the magnitude and sensitivity combine to produce a minor overall impact level during the operation phase of the Project, pre-mitigation (Table 21). Following the application of appropriate mitigation measures, the impact significance is expected to remain the same, that is, minor, primarily because the magnitude will still remain the same during operation, and the sensitivity of the receptor will remain the same (Table 21).

**Table 21: Potential impacts in the operation phase to the Mud Snail**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the	Significance





					Recept or	
Habitat quality and quantity	Medium - 3	High – 4	Major - 12	Low – 2	High – 4	Moderate – 8
Habitat connectivity	Negligible – 1	High – 4	Minor – 4	Negligible – 1	High – 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA, and hence potential local populations of Mud Snail. However, there is some uncertainty in regard to this species actually occurring in the CHAA. As such, the above assessment has been undertaken based on a precautionary approach.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**10.2.2.2 What impact could the operation of the Project have on Grey Crowned Crane?**

Grey Crowned Crane is an Endangered species. In the CHAA, it is associated with permanent and seasonal wetlands habitats.

The operation of the Production Facility could indirectly impact Grey Crowned Crane breeding and foraging habitat, given that operational impacts on wetlands in the Buhuka Flats are expected (Section 9.1.1.2); largely as a result of the induced impacts brought about by the escarpment road on the Buhuka Flats wetland habitats used by this species.

The potential direct, indirect and induced impacts to this species due to the operation of the Production Facility are presented below.

**Impact Indicators**

Indicators used to assess effects of the operation of the Production Facility on Grey Crowned Crane were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to indirect disturbance and edge effects was estimated by applying a 200 m buffer to the Project infrastructure. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance like light, noise, vibration, and edge effects.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat for Grey Crowned Crane (permanent and seasonal wetlands) in relation to the Project infrastructure to qualitatively identify areas where habitat remains fragmented during operation.

Potential changes in abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration). These disturbances were considered in light of known or inferred effects to the survival and reproduction of Grey Crowned Crane for which data on these types of effects are available.

**Impact Assessment**

■ **Habitat Quality and Quantity**

No direct losses of wetland habitat are expected during the Production Facility’s operation, however, additional indirect losses of suitable foraging habitat (seasonally flooded grassland) may occur in the event that wetlands





being crossed or encroached by Project infrastructure incur changes in hydrological flow patterns if construction design mitigation is not successful, potentially resulting in desiccated or permanently flooded habitats.

Although Grey Crowned Crane can tolerate a low degree of anthropogenic disturbance (e.g. agricultural cultivation), in Ugandan wetlands it has been observed to be intolerant of human proximity within 100-200m (Olupot, 2014), flying away on approach; a factor which also affects breeding success as breeding birds flush from nests on approach, causing increased rates of predation, reduced time at the nest, either incubating or feeding, and ultimately nest abandonment. How tolerant foraging/roosting/breeding Grey Crowned Crane may be to indirect disturbances, such as noise, light, vibration and edge effects, is not known. The application of a 200m buffer around the Project infrastructure footprint indicates that approximately 4.64 ha of seasonally flooded wetland habitat will be reduced in quality as a result of sensory disturbance, throughout the operational lifetime of the Project.

The escarpment road to the Project is expected to facilitate an influx of people into the area, seeking opportunities. It can reasonably be expected that the increased human population in the Buhuka Flats will exacerbate existing pressures on the wetlands of the area. These pressures are likely to manifest in increased rates of habitat loss and degradation, resulting in long-term reduction of habitat quality and quantity for Grey Crowned Crane.

Impacts from loss of habitat and reduction of habitat quality are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Habitat Connectivity

The effect of the Project operation as a barrier to the movement of Grey Crowned Crane is likely to be adverse. The operation activity is likely to create noise and visual sensory disturbances, which could elicit reduced use or complete avoidance of affected areas, thereby creating movement barriers. The presence of power lines between the CPF and development wells will present a significant barrier to movement for Grey Crowned Crane, as this species is susceptible to in-flight collisions with overhead powerlines (BirdLife International, 2018). The presence of roads crossing the wetlands has the potential to create barriers to movement as a result of the associated disturbance due to human presence and traffic in the area. The increased human population, and the associated increase in head of livestock grazing in the Buhuka Flats, is expected to further degrade wetland habitats and reduce habitat connectivity for Grey Crowned Crane.

Effects from disruption of habitat connectivity are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

Grey Crowned Crane was relatively common in the Buhuka Flats during baseline surveys, with up to 14 individuals frequently observed. Given the extent to which they reportedly tolerate human presence (approx. 100m – 200m; Olupot, 2014), the large-scale changes in the human population on the Flats that have occurred with the operation of the escarpment road are likely to have affected Grey Crowned Crane occurrence on the Buhuka Flats. The ongoing population influx to the Flats following development of the escarpment road is likely to exacerbate existing levels of sensory disturbance in the locality, which in combination with anthropogenic degradation of suitable foraging, roosting and breeding habitat is expected to adversely affect the abundance and distribution of the species.

Effects on the abundance and distribution of Grey Crowned Crane are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

Although Grey Crowned Crane were observed forming breeding pairs during the baseline fieldwork, the widespread degradation of suitable breeding habitat (permanent wetlands) is expected to become exacerbated throughout the operational lifetime of the Project, largely due to the population increase on the Buhuka Flats. This habitat degradation and increased human presence is expected to significantly reduce the likelihood of Grey Crowned Crane continuing to select these areas for breeding.





As mentioned above, direct disturbance via human proximity within 100-200m causes breeding birds to flush from nests on approach (Olupot, 2014), which may result in increased rates of predation, reduced time at the nest, either incubating or feeding, and ultimately nest abandonment, affecting reproductive success. The increased human population on the flats is also expected to result in increased hunting of cranes for bush meat and capture for domestication, and taking of eggs for food.

It is assumed that indirect disturbances arising from noise, light, and vibration would continue to affect the breeding success of Grey Crowned Crane on the Buhuka Flats, should they nest there during operation.

Removal of large trees which have importance as night-time roosts and day-time shelter from the midday sun has been indicated in localised declines in Grey Crowned Crane in Uganda (Olupot, 2014); the absence of such trees during the operational phase is therefore expected to have implications for the survival of flocks and individuals in the area.

These effects to the survival and reproductive success of Grey Crowned Crane are predicted to be well beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Grey Crowned Crane's sensitivity is high because it is an Endangered species and it triggers Tier 2 critical habitat.

Impacts to Grey Crowned Crane habitat quantity and quality may be adverse during operation. The geographical extent of impacts will be local because effects will be restricted to the LSA, with approximately 4.64 ha (6.72 %) of potential habitat on the Buhuka Flats indirectly affected. Impact duration will be into the far future (that is, ~25 years) because changes to the habitat quality and quantity from sensory disturbances are expected to continue throughout the operational lifetime of the Production Facility. The magnitude of the effects of operation on Grey Crowned Crane habitat quantity and quality is considered high (ref. Section 9.2.1.2). Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the operation phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain major, because the magnitude of effects can at best be reduced to medium as the pressures on wetland integrity associated with human influx are expected to be difficult to mitigate, and the sensitivity of the receptor will remain the same (Table 22).

Impacts to Grey Crowned Crane habitat connectivity will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas where wetlands will be crossed by access roads. Impact duration will be long-term, as the presence of the access roads will remain in place throughout the operational lifetime of the Project, with ongoing sensory disturbances arising from traffic and the increased human population using the access roads continuing into the far-future; the increased human population living on the flats, and the associated increase in head of livestock grazing in the Buhuka Flats is likely to further compromise habitat connectivity for Grey Crowned Crane. The magnitude of the effects of operation on Grey Crowned Crane habitat connectivity is considered high. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the operation phase of the Project, pre-mitigation. Following the application of mitigation measures, the impact significance is expected to remain major, because the magnitude of effects is likely to remain medium as the pressures on wetland integrity associated with human influx are expected to be difficult to mitigate, and the sensitivity of the receptor will remain the same (Table 22).

Impacts to Grey Crowned Crane abundance and distribution will be adverse. The geographical extent of impacts will be local, as impacts will be restricted to wetlands in the Buhuka Flats. Impact duration will be into the far future (that is, ~25 years) as the sensory disturbance and habitat degradation associated with the ongoing population influx to the Flats is likely to remain ongoing throughout the lifetime of the Production Facility. The magnitude of the effects of operation on Grey Crowned Crane abundance and distribution is thus considered high. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain major, because the magnitude of effects can at best





be reduced to moderate as the pressures on Grey Crowned Crane (wetland) habitat integrity associated with human influx are expected to be difficult to mitigate. The sensitivity of the receptor will remain the same (Table 22).

Impacts to the survival and reproduction of Grey Crowned Crane will be adverse. The geographical extent of impacts will be local because effects are restricted to wetlands on the Buhuka Flats. Impact duration will be into the far future (that is, ~25 years) as the sensory disturbance and habitat degradation associated with the ongoing population influx to the Flats is likely to remain ongoing throughout the lifetime of the Production Facility. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is considered probable that at least this percentage of individuals in the local population within the Buhuka Flats will be affected. The magnitude of the effects of construction on the survival and reproduction of the Grey Crowned Crane is therefore high. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain major, because the magnitude of effects can at best be reduced to moderate as the pressures on Grey Crowned Crane habitat integrity associated with human influx are expected to be difficult to mitigate. The sensitivity of the receptor will remain the same (Table 22).

**Table 22: Potential Production Facility operation phase impacts to Grey Crowned Crane**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quantity and quality	High – 4	High – 4	Major - 16	Medium - 3	High – 4	Major – 12
Habitat connectivity	Medium - 3	High – 4	Major – 12	Medium - 3	High – 4	Major – 12
Abundance and distribution	High – 4	High – 4	Major – 16	Medium - 3	High – 4	Major – 12
Survival and reproduction	High – 4	High – 4	Major - 16	Medium - 3	High – 4	Major – 12

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced (the upgraded R5 and P1 roads) impacts that the operation of the Production Facility could have on Grey Crowned Crane and its habitat in the CHAA.

The spatial extent of the wetland habitat in the CHAA was broadly mapped based on knowledge of the study area, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**10.2.2.3 What impact could the operation of the Project have on Nahan’s Francolin?**

Nahan’s Francolin is a Vulnerable and range-restricted species. In the CHAA, it is restricted to the Bugoma Central Forest Reserve. It triggers Tier 2 critical habitat under Criterion 2 (Section 6.3.2.2, Appendix G).

The potential induced and cumulative effects to this species due to the operation of the upgraded R5 and P1 roads are presented below.





### *Impact Indicators*

Indicators used to assess effects of the operation of the upgrade of the R5 and P1 roads on Nahan's Francolin were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to indirect disturbance and edge effects was estimated by applying a 200 m buffer to the road corridors. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance like light, noise, vibration, and edge effects.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the road corridor to qualitatively identify areas where habitat becomes fragmented.

Potential changes in abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration). These disturbances were considered in light of known or inferred effects to the survival and reproduction of buttonquail species for which data on these types of effects are available.

### *Impact Assessment*

#### ■ Habitat Quality and Quantity

The Nahan's Francolin's degree of vulnerability to disturbance is unknown. Birds are known to be sensitive to land use and habitat alteration (Lussier et al. 2006). The behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions, such as the availability of alternative foraging sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the population of this species in Bugoma CFR.

How tolerant the Nahan's Francolin may be to indirect disturbances, such as noise, light, vibration and edge effects, is not known. However, assuming it is sensitive to such disturbance because it is a shy, forest-dependent species (Sande et al. 2009a), then potential Nahan's Francolin habitat will be lost or reduced in quality as a result of edge effects, and possibly sensory disturbance.

The improved R5 and P1 roads could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of habitat alteration and deforestation, as well as increased bush meat hunting. These factors could combine to reduce the habitat quality and quantity for Nahan's Francolin.

Effects from loss of habitat are predicted to be well beyond the expected range of natural disturbance perturbations.

#### ■ Habitat Connectivity

The upgraded R5 and P1 roads will not directly remove suitable habitat for the Nahan's Francolin. However, may affect that habitat within the 200 m buffer through indirect impacts like edge effects and sensory disturbance. As such, the road may become an effective barrier to the movement of Nahan's Francolin if the levels of traffic on the road substantially increase over the life of the Project. Roads are recognised as creating sensory disturbances, which can elicit reduced use or complete avoidance of affected areas, thereby creating movement barriers (for example, see Kolowski and Alonso 2009, Gleeson and Gleeson 2012). However, given that the Hoima-to-Ikamiro Road already exists, if the Nahan's Francolin is sensitive to such effects, then it is reasonable to assume the existing road is already a barrier to a greater or lesser degree.

The improved R5 and P1 roads could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central



Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of habitat alteration and deforestation. These factors could reduce the habitat connectivity for Nahan's Francolin.

Effects from loss of habitat connectivity are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

Nahan's Francolin is reported to be relatively common in the Bugoma Central Forest Reserve (Plumptre et al. 2011). What effects the upgrade of the road, and the predicted increased traffic levels along the road during the operation of the Project, could have on the abundance and distribution of this species is unknown.

With the operation of the upgraded road, and the increased traffic along the road, it is reasonable to assume that the local abundance and distribution of this species may be altered. Individuals may avoid the resultant sub-optimal habitat brought about by the upgraded road and increased traffic volumes. Additionally, the increased traffic on the road could lead to an increase in direct mortality of individuals.

The improved R5 and P1 roads could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration. These factors could reduce the abundance and distribution of Nahan's Francolin.

Effects from the upgraded road are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

The Nahan's Francolin's degree of vulnerability to direct disturbance, particularly during the breeding season, is not well understood. Generally, for birds, the behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions such as the availability of alternative breeding sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the local population of Nahan's Francolin.

How tolerant the Nahan's Francolin may be to indirect disturbances, such as noise, light, vibration and edge effects, during the breeding season, are also not completely understood. What is known is that disturbance of mature forest generally appears to reduce the home range of this species (Sande et al. 2009b), and hence its potential to find suitable mates. Assuming it is sensitive to sensory disturbance because it is a shy, forest-dependent species (Sande et al. 2009a), the operation of the upgraded roads, and the increased traffic levels on the road could affect the breeding success of those individuals living in close proximity to the road.

The improved R5 and P1 roads could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration. These factors could reduce the survival and reproduction of Nahan's Francolin.

These effects to the survival and reproductive success of Nahan's Francolin are predicted to be well beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Nahan's Francolin's sensitivity is high because it is range-restricted, and triggers Tier 2 critical habitat.

Impacts to the Nahan's Francolin's habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 and P1 road corridors in the CHAA. Impact duration will be long-term. The magnitude of the effects of operation on the





habitat quantity and quality of the Nahan’s Francolin is low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operational phase of the Project, pre-mitigation (Table 25).

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 23).

Impacts to the Nahan’s Francolin’s habitat connectivity will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 and P1 road corridors in the CHAA. Impact duration will be long-term. The magnitude of the effects of operation on the habitat connectivity of the Nahan’s Francolin is Low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operational phase of the Project, pre-mitigation.

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 23).

Impacts to the Nahan’s Francolin’s abundance and distribution could be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 and P1 road corridors in the CHAA. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed during the operation phase. The magnitude of the effects of operation on the abundance and distribution of the Nahan’s Francolin is Low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Project, pre-mitigation.

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 23).

Impacts to the survival and reproduction of Nahan’s Francolin could be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the R5 and P1 road corridors in the CHAA. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed during the operation phase. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve will not be affected. The magnitude of the effects of operation on the survival and reproduction of the Nahan’s Francolin is therefore low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operational phase of the Project, pre-mitigation (Table 23).

Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, and the sensitivity of the receptor will remain the same (Table 23).

**Table 23: Potential impacts in the operational phase to the Nahan’s Francolin**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8
Habitat connectivity	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8
Abundance and distribution	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8





Survival and reproduction	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8
---------------------------	---------	----------	--------------	---------	----------	--------------

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced (the upgraded R5 and P1 roads) impacts that the operation of the Project could have on the Nahan’s Francolin and its habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**10.2.2.4 What impact could the operation of the Project have on Eastern Chimpanzee?**

Eastern Chimpanzee is an Endangered species. A population occurs in the Bugoma Central Forest Reserve. This species triggers Tier 1 critical habitat.

The potential induced effects of the upgraded R5 and P1 roads on Eastern Chimpanzee are presented below.

**Impact Indicators**

Indicators used to assess effects of the upgraded R5 and P1 roads on Eastern Chimpanzee were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to indirect disturbance and edge effects during operation was estimated by applying a 500 m buffer to the road corridor. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance like noise, vibration and traffic. The buffer width was determined based known chimpanzee sensitivities to noise disturbance (Parren and Byler 2003, Rabanal et al. 2010). The amount of loss or degradation of habitats within the buffer was evaluated across a range of possibilities, including that habitats become completely unavailable to chimpanzees during the operation phase. Evaluating the potential for complete avoidance of the buffer area is a conservative approach, which addresses uncertainty about the attenuation distance of sensory disturbances for chimpanzees, even though the likelihood of strict avoidance throughout the entire buffer may be low.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat in relation to the road corridor to qualitatively identify areas where habitat becomes fragmented.

Potential changes in abundance and distribution were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration). These disturbances were considered in light of known or inferred effects to the survival and reproduction of other populations of chimpanzees for which data on these types of effects are available.

To assess effects to survival and reproduction as a result of in-migration and potential associated increases in poaching and disease spread, in-migration rates were predicted based on the predictions in the influx management plan. A literature review of the impact of contact with humans was also conducted

**Impact Assessment**

■ Habitat Quality and Quantity

Based on the data presented in Plumptre et al. (2010), the distribution of nesting sites appears to be widely distributed through the forest; the exception being within the vicinity of the Hoima-to-Ikamiro Road. Here the frequency of nests encountered by those workers was a lot lower than elsewhere in the forest. This suggests that the Eastern Chimpanzees within the Bugoma Central Forest Reserve are actively avoiding the road, preferring to sleep some distance away from the disturbances arising from the road. The observation supports





the findings of Parren and Byler (2003) that chimpanzees actively avoid environments where they will be disturbed at night.

The degree of vulnerability to disturbance experienced by chimpanzees is reasonably well known (for example, see Parren and Byler 2003, Rabanal et al. 2010, Thompson and Wrangham 2013). The chimpanzees living in and around the Bugoma Central Forest Reserve most likely experience sensory disturbances at present from human activities, given the high human populations living around the reserve. Indeed, groups are known to forage in the agricultural fields surrounding the forest, and hence, would more than likely be used to human noises and disturbances (McLennan 2008). As such, the potential sensory disturbances arising from the operation of the upgraded road are likely to be minimal. Nevertheless, the magnitude of noise may not be the most important determinant of chimpanzee response. Instead, chimpanzees may respond to 'new' noises or may associate particular noises with other occurrences (for example, machine noise may be associated with human presence, which chimpanzees may, in turn, associate with the presence of danger). Where humans pose a threat, chimpanzees generally avoid them (Hockings and Humle 2009, Parren and Byler 2003). Therefore, the degree of avoidance may depend on the behaviour of people, highlighting the importance of managing contractor activity.

The improved R5 and P1 roads could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration. These factors could reduce the survival and reproduction of Eastern Chimpanzee.

Effects from the loss of habitat are predicted to be well beyond the expected range of natural disturbance perturbations.

### ■ Habitat Connectivity

Although the operation of the upgraded R5 and P1 roads will not cause direct losses of suitable habitat for the Eastern Chimpanzees; it may affect suitable habitat within the 500 m buffer through indirect impacts like edge effects and sensory disturbance.

However, the chimpanzees within Bugoma Central Forest Reserve are assumed to be accustomed to human activity, and they may, more than likely, regularly cross roads within their range. Therefore, it is possible that they will not completely avoid the road corridor when the road is operational.

As such, the effect of the operation of the upgraded roads and associated traffic volumes within Bugoma CFR as a barrier to chimpanzee movements is predicted to be negligible – assuming that the mitigation recommendation of restricted access on the R5 intersecting the reserve has been applied.

Effects from loss of habitat connectivity are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

The Bugoma Central Forest Reserve supports one of the top four Eastern Chimpanzee populations in Uganda, with a population of between 450 and 850 individuals (Plumptre et al. 2010). Although the increased traffic volumes expected on these roads during operation has the potential to cause direct mortality of individuals, should such individuals cross the road, the operation of the upgraded roads is not expected to detrimentally affect the abundance of chimpanzees in the forest directly. What effect the loss of individuals from the population could have is unknown. It is doubtful though that the population could be reduced by 10% and hence, reach that critical population threshold due to road mortalities alone.

The long-term presence of the improved R5 and P1 roads could continue to facilitate influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration. These factors could reduce the abundance and distribution of Eastern Chimpanzee.



Effects from the upgraded road are predicted to be well beyond the expected range of natural disturbance perturbations, although not beyond the disturbance from the human-induced deforestation rate and habitat loss, and bush meat hunting in Bugoma Central Forest Reserve.

### ■ Survival and Reproduction

In-migration associated with upgrade of the R5 and P1 roads may adversely affect survival and reproduction of chimpanzees through poaching and disease transfer. Hunting and poaching is a recognised threat to chimpanzees in Bugoma Central Forest Reserve (Plumptre et al. 2003), and this pressure could increase as in-migration of people from other areas occurs. Mitigation and management of in-migration will be very important to minimise potential effects on chimpanzees and other fauna species.

Disease is one of the major threats to Eastern Chimpanzees (Oates et al. 2008); increased abundance of people and competition for land and food resources between humans and chimpanzees could lead to higher rates of disease spread from humans to chimpanzees. Chimpanzees are closely related to humans; therefore, many diseases are transferrable between chimpanzees and humans (Formenty et al. 2003; Isabirye-Basulta and Lwanga 2008). Either direct or indirect contact with humans can spread disease. For example, there is evidence to suggest that respiratory illnesses have been transferred directly to chimpanzees from humans as a result of researcher and tourist contacts, often leading to outbreaks and death (Formenty et al. 2003). Human defecation in forest undergrowth can indirectly lead to spread of intestinal diseases, such as *Clostridium perfringens*, which can be fatal to chimpanzees (Fujita 2011). As forest fragments decrease in size, risks of contact with, and transmission of, disease from humans increases (Isabirye-Basulta and Lwanga 2008). Factors that lead to increased crop raiding or sharing of water resources can also increase the risk of disease spread (Hockings and Hulme 2009). Communicable respiratory diseases are a significant concern in the LSA.

An increasing human population, facilitated by an improved road, could lead to an increase in the demand for agricultural resources, which, in turn could further fragment, change, and degrade the Bugoma Central Forest Reserve and surrounding habitat. With the upgrade of the road, and the potential for the Project to be attractive for people seeking opportunities, could see a substantial increase the current population around the Bugoma Central Forest Reserve.

Effects from the upgraded road are predicted to be well beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Eastern Chimpanzee's sensitivity is high because it is Endangered.

Impacts to the Eastern Chimpanzee's habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the Hoima-to-Ikamiro Road corridor in the CHAA. Impact duration will be long-term. The magnitude of the effects of construction on the habitat quantity and quality of the Eastern Chimpanzee is low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Project, pre-mitigation (Table 20). Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 24).

Impacts to the Eastern Chimpanzee's habitat connectivity will be neutral. The geographical extent of impacts will be local because effects are restricted to those areas immediately adjacent to the Hoima-to-Ikamiro Road corridor in the CHAA. Impact duration will be long-term. The magnitude of the effects of construction on the habitat connectivity of the Eastern Chimpanzee is negligible. Therefore, the magnitude and sensitivity combine to produce a minor overall impact level during the operational phase of the Project (Table 24).

Impacts to the Eastern Chimpanzee's abundance and distribution could be adverse. The geographical extent of impacts will be regional because, although effects are restricted to those areas immediately adjacent to the Hoima-to-Ikamiro Road corridor in the CHAA, there is the potential to extend beyond that area. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed during the operational phase. The magnitude of the effects of operation on the abundance and distribution of the Eastern



Chimpanzee is medium. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the operation phase of the Project, pre-mitigation (Table 24). Following the application of appropriate mitigation measures, the impact significance is expected to become moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 24).

Impacts to the survival and reproduction of the population of Eastern Chimpanzee in Bugoma Central Forest Reserve could be adverse. The geographical extent of impacts will be national because effects could extend well beyond the CHAA. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed during the operation phase. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve could be affected through direct mortality from bush meat hunting or other disturbance. Therefore, the magnitude of the effects of operations on the survival and reproduction of the Eastern Chimpanzee high. Therefore, the magnitude and sensitivity combine to produce a major overall impact level during the road upgrade and construction phase of the Project, pre-mitigation (Table 24).

Following the application of appropriate mitigation measures, the impact significance is expected to become moderate, primarily because the effectiveness of mitigation measures in the long-term are unknown and beyond the control of CNOOC, although the sensitivity of the receptor will remain the same (Table 24).

**Table 24: Potential impacts in the operation phase to the Eastern Chimpanzee**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate – 8	Low – 2	High – 4	Moderate – 8
Habitat connectivity	Negligible – 1	High – 4	Minor – 4	Negligible – 1	High – 4	Minor – 4
Abundance and distribution	Medium – 3	High – 4	Major – 12	Low – 2	High – 4	Moderate – 8
Survival and reproduction	High – 4	High – 4	Major – 12	Low – 2	High – 4	Moderate – 8

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced (the upgraded R5 and P1 roads) impacts that the operation of the Project could have on the Eastern Chimpanzees and their habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.0). The actual extent of possible habitat may have been over-estimated.

**11.0 IMPACT ASSESSMENT – FEEDER PIPELINE**

This section presents an assessment of the possible interactions of biodiversity valued components with the Feeder pipeline infrastructure and activities, and the resulting impacts during the construction and operation phases.

The biodiversity valued components for the Feeder pipeline impact assessment are listed in Table 5 below. They include all of the species and habitats that trigger critical habitat designation within the CHAA. In addition, ecosystems of concern that will be potentially affected by the Project, and Grey Crowned Crane, were also







included as valued components for impact assessment, for reasons outlined in the Table below. As mentioned in Section 10.1.2, potential impacts to other species of concern are assessed at the habitat level (ecosystems of concern).

**Table 25: Biodiversity Valued Components for Feeder Pipeline Impact Assessment**

Valued Component	Confirmed CH Trigger? (see Table 4)	Reasoning (see Table 4)
Near-shore aquatic habitats of Lake Albert	<ul style="list-style-type: none"> <li>■ Yes – Criterion 13</li> <li>■ Possibly Criterion 1 and Criterion 2 (G. candida)</li> </ul>	<ul style="list-style-type: none"> <li>■ The near-shore habitats are important fishing grounds that support 11 fishing villages on the Buhuka Flats and surrounds (see Chapter 18 Ecosystem Services Review)</li> <li>■ May support the CR and range-restricted species <i>Gabiella candida</i></li> </ul>
Wetlands	<ul style="list-style-type: none"> <li>■ No</li> </ul>	<ul style="list-style-type: none"> <li>■ Natural habitat – priority habitat according to IFC (2012)</li> <li>■ Supports Endangered Grey Crowned Crane</li> <li>■ Important in supply of ecosystem services to local communities (see Chapter 18 Ecosystem Services Review)</li> </ul>
Escarpment vegetation corridor	<ul style="list-style-type: none"> <li>■ No</li> </ul>	<ul style="list-style-type: none"> <li>■ Natural habitat – priority habitat according to IFC (2012)</li> <li>■ Forms part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor</li> <li>■ The location of caves and cavities along the escarpment that could be important for cavity-roosting bats</li> </ul>
Bugoma Central Forest Reserve	<ul style="list-style-type: none"> <li>■ Yes –</li> <li>■ Criterion 4</li> <li>■ Criterion 1</li> <li>■ Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>■ Triggers CH on the basis of being a highly threatened and unique ecosystem (Criterion 4)</li> <li>■ Triggers Criterion 1 Tier 1 CH on the basis of support of a population of Eastern Chimpanzee, that is recognised as being one for the four largest in the region; apart from being an Endangered species, chimpanzees are also recognised as key stone species and ecosystem engineers</li> <li>■ Triggers Criterion 2 Tier 2 CH on the basis of support of range-restricted Nahan’s Francolin</li> <li>■ Recognised area of old growth forest</li> <li>■ The forest is recognised for its unique biodiversity values, including biome restricted species</li> <li>■ Is an important ecosystem service supply area for local people who harvest timber, fibre, fuel wood and charcoal, and non-timber forest products from the forest</li> <li>■ Bugoma Central Forest Reserve is recognised as an Important Bird Area</li> </ul>





Valued Component	Confirmed CH Trigger? (see Table 4)	Reasoning (see Table 4)
		<ul style="list-style-type: none"> <li>Nationally recognised as a high conservation priority area (NEMA 2010)</li> </ul>
Mud Snail ( <i>Gabbiella candida</i> )	<ul style="list-style-type: none"> <li>Possibly Criterion 1 and Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>Could occur on near-shore aquatic habitats (Bugoma Lagoon, large bays, open sandy shores, shallow river-associated water)</li> <li>Has not been confirmed in LSA to date and is included on basis of precautionary principle</li> </ul>
Grey Crowned Crane	<ul style="list-style-type: none"> <li>No</li> </ul>	<ul style="list-style-type: none"> <li>Although Grey Crowned Crane is not present in numbers that would trigger CH designation, it is an Endangered species and has been confirmed present on the Buhuka Flats during baseline fieldwork in 2014 and 2017</li> <li>Any potential Project impacts on a globally-recognised and nationally-protected Endangered species are unacceptable and warrant addressing via the impact assessment process</li> </ul>
Nahan's Francolin ( <i>Ptilopachus nahan</i> )	<ul style="list-style-type: none"> <li>Yes – Criterion 2</li> </ul>	<ul style="list-style-type: none"> <li>Occurs in Bugoma Central Forest Reserve, possibly one of less than 10 DMUs globally (including DRC)</li> <li>Potential for CHAA to support &gt;10% of this species' known global population</li> </ul>
Eastern Chimpanzee ( <i>Pan troglodytes schweinfurthii</i> )	<ul style="list-style-type: none"> <li>Yes – Criterion 1</li> </ul>	<ul style="list-style-type: none"> <li>Occurs in Bugoma Central Forest Reserve</li> <li>Great apes are an iconic species of anthropological and evolutionary significance</li> <li>They generally immediately trigger CH designation (see GN 74 and footnotes in PS6, IFC 2012a and b)</li> </ul>

It should be noted that there is limited potential for decommissioning phase impacts for the Feeder pipeline, as typically pipelines are left in situ following decommissioning. In any case, the decommissioning activity will be the subject of a separate ESIA process, to be conducted towards the end of the operational phase.

### 11.1 Construction Phase Impacts

The construction of the Feeder pipeline will occur over 10-12 months, and is expected to present direct impacts to valued components including sensory disturbances, the potential for pollution and erosion and sedimentation, and direct mortality from vehicle movements and entrapment in open trenches. Indirect impacts on valued components are also anticipated as a result of population influx, due to expectations of work on the construction contract.

For the assessment of impacts during the construction phase, the key questions were divided into sub-questions that focused on individual valued components within the CHAA and LSA. In answering each question, the individual components of the pipeline were considered with regards to their potential to affect a valued component. These questions are presented below.





### 11.1.1 What impact could the construction of the Feeder Pipeline have on habitats and ecosystem integrity?

This section presents the assessment of impacts that the construction and decommissioning of the Feeder Pipeline could have on the habitat and ecosystem integrity within the CHAA and the LSA. These habitats either do, or could, support populations of species of concern. Therefore, the assessment of potential impacts to those species, and others, occurring in the CHAA has been assessed in this section through the determination of the impacts to potential habitat for those species.

The impacts of the Feeder Pipeline on critical habitat, as triggered by species of concern, are covered under the individual assessment of those species in Section 11.1.2. Other triggers of critical habitat are discussed, as relevant, in the appropriate sections.

#### 11.1.1.1 What impact could the construction of the Feeder Pipeline have on the near-shore aquatic habitats of Lake Albert?

##### *Impact Indicators*

The Feeder pipeline will not be constructed in the vicinity of the near-shore aquatic habitats of Lake Albert and thus is not expected to directly impact these habitats in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. However, changes in water quality as a result of the proposed discharge of hydrotest water to Lake Albert could affect the habitat and ecosystem integrity of the near-shore environment of Lake Albert.

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the near-shore environment of Lake Albert were changes in: regional representativeness; topography (geomorphology) and sediments; water quality; ecosystem composition; ecosystem configuration.

Potential losses of habitat due to direct disturbance and clearing associated with the Project was quantified by overlaying the current, baseline extent of the habitat with the Project footprint.

Additional, indirect impacts to habitat were estimated by applying a 1 km buffer to the Project footprint, forming the LSA. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by sensory disturbance, changes in water quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential succession changes in species composition that could occur. This was accomplished by examining available literature about the ecology of Lake Albert, and scientific literature about the impacts of human activities on aquatic environments.

##### *Impact Assessment*

###### ■ Representativeness

The CHAA supports 16.2 km of near-shore aquatic habitats, equating to about 810 ha (based on the near-shore habitat extending 0.50 km into the lake, see Section 7.1.1).

The Feeder pipeline will not be constructed in the vicinity of the near-shore aquatic habitats of Lake Albert and thus is not expected to directly impact these habitats in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. Therefore, no effects on regional representativeness are predicted.

###### ■ Topography (geomorphology) and sediment transport

The Feeder pipeline will not be constructed in the vicinity of the near-shore aquatic habitats of Lake Albert and thus is not expected to directly impact these habitats in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. Therefore, no pipeline-related impacts on the topography (geomorphology) and/or sediment transport in the near-shore habitats are predicted.

###### ■ Water Quality





The commissioning of the Feeder pipeline will involve hydrotesting, in which the vessels are filled with water and pressurised to verify their integrity. On occasions, chemical additives such as biocides, oxygen scavengers, dyes and/or corrosion inhibitors are added to the water, depending on the residence time, before it is discharged. This water is proposed to be discharged directly to Lake Albert. Without management, its release can present a severe risk to the aquatic environment, resulting in possible mortality and degradation of downstream ecosystems and species.

### ■ Ecosystem composition

All six of the main habitat types in Lake Albert, as identified for fishes by Wandera and Balirwa (2010) (that is, shallow river-associated waters, open sandy shores, lagoons, large bays, rocky escarpments, and, open-water habitats), occur within the near-shore areas of the CHAA (see Section 7.1.1, Figure 5). Similarly, the species guilds associated with the near-shore habitats of the CHAA, in particular fish, are well represented throughout those regions of the lake that have been investigated (for example, see Wandera and Balirwa 2010, Taabu-Munyaho et al. 2012). Consequently, at baseline, the composition of these ecosystems can be said to be in good condition and reflective of the aquatic diversity of Lake Albert. Similarly, these aquatic habitats have a well-developed structure, that is, well-defined aquatic plant layers associated with underwater features and substrates.

If not correctly treated, the discharge of hydrotest fluids directly to Lake Albert during pipeline construction is likely to result in high-intensity, temporary and localised alteration of the ecosystem composition of the aquatic communities; particularly those inhabiting the near-shore habitats in the vicinity of the discharge outfall.

The point source discharge, and the quantity of hydrotest fluid involved, presents a risk of localised toxicity, which could cause mortality of fish and aquatic invertebrates, algae and plants, thereby changing diversity and complexity of the aquatic habitats and their ability to support associated aquatic faunal communities. Impacts from the changes to ecosystem composition as a result of this contamination are predicted to be beyond the expected range of natural disturbance perturbations.

Since the near-shore habitats of Lake Albert within the CHAA potentially support species that trigger critical habitat including the Critically Endangered mud snail (*G. candida*); any negative impacts on the ecosystem composition of this habitat have the potential to be of major significance.

### ■ Ecosystem configuration

The Feeder pipeline will not be constructed in the vicinity of the near-shore aquatic habitats of Lake Albert and thus is not expected to directly impact these habitats in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. Therefore, no pipeline-related impacts on the connectivity, or ecosystem configuration of the near-shore habitats are predicted.

## Impact Classification

The near-shore aquatic habitat's sensitivity is high because these habitats potentially support populations of the Critically Endangered Mud Snail (*G. candida*), the Vulnerable African Soft-shelled Turtle (*T. triunguis*), and the range-restricted and Near Threatened Snail (*Bellamya rubicunda*). Near-shore aquatic habitat within the CHAA potentially constitutes Tier 1 Critical Habitat for the Mud Snail (*G. candida*) (Table 4). Impacts on this habitat are therefore classified on the basis of high sensitivity to potential effects of the proposed development.

### ■ Representativeness

Impacts to the representativeness of the habitat will be neutral, as the Feeder pipeline will not be constructed in the vicinity of the near-shore aquatic habitats of Lake Albert, and thus is not expected to directly impact these habitats in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. Therefore, no effects on regional representativeness are predicted.

### ■ Topography (geomorphology) and sediment transport

Impacts to the topography (geomorphology) and sediment transport in the near-shore aquatic habitats of Lake Albert will be neutral, as the Feeder pipeline will not be constructed in the vicinity of these habitats, and thus is not expected to directly impact them in terms of direct or indirect habitat loss as a result of disturbances and





clearing during construction. Therefore, no pipeline-related impacts on the topography (geomorphology) and/or sediment transport in the near-shore habitats are predicted

■ Ecosystem composition – water quality

Impacts to the water quality, and subsequently the ecosystem composition of near-shore habitats will be adverse. The geographical extent of impacts is expected to be localised, around the Lake Albert discharge point. The duration will be temporary (that is, over the course of several weeks), and will only occur once the construction of the full pipeline has been completed i.e. at the end of the construction phase.

The magnitude of the potential impacts of the direct discharge of hydrotest effluent from the pipeline on the water quality is medium. This is because, prior to mitigation, the potential for hydrotest discharge to alter the baseline water quality is substantial, resulting localised changes to ecosystem composition as a result of toxic effects on aquatic invertebrate, fish and plant communities (i.e. the diversity and complexity of the habitat) around the discharge point. As mentioned previously, the near-shore aquatic habitat’s sensitivity is high, therefore, the magnitude and sensitivity combine to produce a major overall impact level to water quality as a result of potential contamination with potentially toxic chemicals during the hydrotesting of the Feeder pipeline, prior to the application of the recommended mitigation (Table 7). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible.

Following the application of the mitigation measures (Section 12.2), the impact significance is expected to be reduced to minor, primarily because the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

■ Ecosystem configuration

Impacts to the ecosystem configuration of the near-shore aquatic habitats of Lake Albert will be neutral, as the Feeder pipeline will not be constructed in the vicinity of these habitats, and thus is not expected to directly impact them in terms of direct or indirect habitat loss as a result of disturbances and clearing during construction. Therefore, no pipeline-related impacts on the ecosystem configuration of the near-shore habitats are predicted.

**Table 26: Potential impacts in the construction phase to near-shore habitats**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Ecosystem composition - Water quality	Medium - 3	High - 4	Major – 12	Negligible - 1	High - 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios with the level of Lake Albert, which has, in the recent and not so recent past, varied quite dramatically (Talbot et al. 2006), it is conceivable that level of the lake may increase or decrease thereby altering near-shore habitats.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 7.1.1). The actual extent of possible habitat may have been over-estimated.





**11.1.1.2 What impact could the construction of the Feeder Pipeline have on the escarpment vegetation corridors?**

**Impact Indicators**

Indicators used to assess impacts of the Project on the habitat and ecosystem integrity of the vegetation corridors on the escarpment were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

Loss of habitat due to direct disturbance and clearing associated with the Feeder pipeline was quantified by overlaying the current, baseline extent of the escarpment vegetation communities with the Feeder pipeline footprint.

Additional, indirect affects to habitat were estimated by applying a 0.50 km buffer to the footprint of the Feeder pipeline. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, changes in water quantity and quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur, and the disruption of known corridors. This was accomplished by examining available literature about the ecology of the vegetation communities on the escarpment, and scientific literature about the impacts of human activities on corridors.

**Impact Assessment**

■ Representativeness

The CHAA supports approximately 2443 ha of escarpment corridor vegetation communities (see Section 6.3.1.1.2), which are bounded on the east by agriculturally modified landscapes, and the Buhuka Flats on the west. As mentioned, these vegetation communities form part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2003) (see Section 7.1.2). Therefore, their continuous extent represents an important landscape feature in the CHAA.

The proposed Feeder pipeline will traverse approximately 0.8 km through escarpment vegetation communities before it enters agriculturally-modified habitats on the plateau above the escarpment. The total area of escarpment habitat that will be lost and disturbed as a result of the construction of the Feeder pipeline is 4.1 ha. All these vegetation communities are widely represented on the escarpment, and the CHAA.

The total loss of escarpment vegetation communities within the CHAA due to the construction of the Feeder pipeline is outlined in Table 27).

**Table 27: Permanent loss of escarpment corridor vegetation due to construction of Feeder pipeline**

Vegetation Type	Total area in the CHAA (ha)	Total area in Escarpment vegetation corridor (ha)	Area of escarpment vegetation corridor affected by the pipeline (ha)	% loss of vegetation corridor within CHAA
Cultivation and settlement	31860.9	0.1	-	-
Dense bushland	1097.6	337.7	1.6	0.5%
Dense wooded grassland	613.2	589.0	1.7	0.3%
Open grassland	568.5	31.5	-	-
Open wooded bushland	523.0	214.0	0.8	0.4%
Open wooded grassland	1900.9	552.2	-	-





Vegetation Type	Total area in the CHAA (ha)	Total area in Escarpment vegetation corridor (ha)	Area of escarpment vegetation corridor affected by the pipeline (ha)	% loss of vegetation corridor within CHAA
Riverine woodland	74.8	69.8	-	-

The loss of this quantity of vegetation in relation to the amount in the CHAA is not substantial. Nevertheless, the loss of this vegetation does open up a previously contiguous tract of vegetation with a linear corridor that introduces edge effects and the concomitant aspects associated with those, as discussed below.

Impacts from the changes to representativeness are predicted to be beyond the expected range of natural disturbance perturbations.

■ Ecosystem composition

The construction of the Feeder pipeline may result in loss of biodiversity at both local and regional scales due to restricting movement between populations, increased mortality, habitat fragmentation and edge effects, invasion by exotic species, or increased human access to wildlife habitats, all of which are expected to increase local extinction rates or decrease local recolonisation rates (Findlay and Bourdages 2000). Indeed, the construction of a linear corridor (such as the Feeder pipeline) through an area of relatively intact vegetation, like that on the escarpment, creates edge effects that could, in the long-term, alter the composition of the ecosystem through which the pipeline traverses. Vegetation clearing creates edges or boundaries where habitat (for example, riverine woodland) meets a disturbance (for example, the pipeline corridor). Edges associated with disturbance are different than transition areas, or ecotones, amongst vegetation communities, because disturbance edges tend to be abrupt with a high degree of contrast between two areas (for example, pipeline corridor and open wooded grassland). Edges open up habitat in areas where it was previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges often result in changes in species composition along the edge, with the edges typically becoming dominated by pioneer species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Often, these microclimates are favoured by weed and/or invasive plant species. However, certainly for fauna species, and depending on the species, edges can have either a positive or negative impact on habitat quality and quantity (Prevedello et al. 2013, Wellicome et al. 2014). Given the length of the Feeder pipeline (approximately 0.8 km) traversing the escarpment vegetation communities, this equates to approximately 1.6 km of edges in the escarpment vegetation corridor, which would have otherwise not existed.

What long-term impacts that these edge impacts could have on the vegetation communities' composition are unknown; in particular their resilience. The escarpment vegetation communities are already under pressure from harvesting of fuel wood and charcoal manufacture, and livestock grazing. Cattle have been shown to substantially increase the edge impacts in savannah habitats (Porensky et al. 2013). Vegetation clearing for the Feeder pipeline could facilitate the introduction and spread of invasive plant species throughout the escarpment vegetation communities, as well as create a nick point for erosion and subsequent degradation of adjacent vegetation. These changes could substantially alter the habitat structure and composition, which, in turn, could affect its utilisation by the current species guild. To some extent this may already be occurring. For example, very few medium sized mammals were recorded in the CHAA, and those that were recorded tended to be thicket and dense bushland specialists, such as bushbuck and duiker. The low populations and diversity of these species could also be a reflection of increased pressure for bush meat from the local human population, which has increased markedly over the last ten years (AECOM 2012). It is expected that the existing trends in this regard will increase markedly once people become aware that project construction is imminent.

The 800 m section of the Feeder pipeline that will intercept the escarpment vegetation corridor will be routed straight up a 76% (40°) slope. The trench for the pipeline will probably need to be drilled and blasted to achieve





the minimum 0.8m depth of cover for rocky ground. During construction, the trenches present a risk of injury and mortality to fauna, in particular smaller mammal and reptile species, which could affect ecosystem composition, albeit in a temporary and localised manner.

If properly managed during construction, including rolling rehabilitation of the buried pipeline once construction is complete, impacts on the ecosystem composition could be minor, nevertheless the potential construction impacts are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Ecosystem configuration

The wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Iwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor is recognised as being important for threatened species in the face of climate change adaptation (Ayebare et al. 2013), and as part of a much broader set of corridors running the length of the Albertine Rift. This corridor in the southern portion of the eastern shores of Lake Albert, in the vicinity of the Project, is very narrow compared to its extent elsewhere, and is recognised as being important for savannah species (Plumptre et al. 2010).

The construction of the pipeline requires a 30 m wide construction right-of-way and will involve digging/blasting of a trench for the pipeline, which will then be buried. The construction activity and concomitant environmental disturbances (visual, noise, lighting, physical barrier) are likely to temporarily reduce its value as a wildlife movement corridor.

Indications are that populations of highly mobile wildlife within the area, which may depend on the corridor, are not substantial, and potentially severely depleted (see Section 6.1). In particular, those species that could utilise this corridor, such as large ungulates, predators, primates and birds, are very rare in the area, apart from birds. Nevertheless, during the construction phase, the trenches for the pipeline could present a barrier to movement for fauna, in particular smaller mammal and reptile species. However, the Feeder pipeline is not expected to be a barrier for birds; for example, White-backed Vultures were seen moving south down the escarpment corridor during the surveys (see Section 6.1.4). What impact the Feeder pipeline could have on the movement of those terrestrial individuals and species that may utilise this corridor is unknown. The corridor is recognised as an important climate change refugium for a range of threatened species, which may become increasingly important for those species in the future (Ayebare et al. 2013).

Direct impacts from the changes to ecosystem configuration on the escarpment as a result of the Feeder pipeline are, therefore, predicted to be beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The vegetation community of the escarpment's sensitivity is medium because these habitats are already under stress from livestock grazing and harvesting of fuel wood and non-timber forest products. However, these communities do form part of a wider, regionally-important wildlife corridor.

Impacts to the representativeness of the habitat will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA, and will only constitute a loss of 4.1 ha, in a total area of escarpment vegetation in the CHAA of 2233.8 ha (Table 27). This amounts to 0.18% of the vegetation in the escarpment corridor in the CHAA. The impact is expected to be medium to long term; while rehabilitation of the affected pipeline corridor will occur immediately after construction is completed, the steep and rocky slopes make it unlikely that habitat recovery by plants other than pioneer species will occur in the short-term. Whether there is relatively complete habitat recovery will depend largely on the effectiveness of the rehabilitation strategy. If normal construction management methods are used (without a high degree of specialist rehabilitation intervention), full recovery is unlikely. Nevertheless, the small area affected is not expected to materially impact on the representativeness of the escarpment vegetation corridor, either within the CHAA or at regional scale, and the magnitude of the impacts of the Feeder pipeline on representativeness of the vegetation communities of the escarpment is rated as low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to representativeness during the construction phase of the Project, pre-mitigation (Table 28). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to low, providing that the recommended mitigation measures (rehabilitation of the







soil and vegetation cover on top of the buried pipeline) are implemented, with a high degree of management intervention necessary both during and after construction.

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. The impact is expected to be medium to long term; while rehabilitation of the affected pipeline corridor will occur immediately after construction is completed, the steep and rocky slopes make it unlikely that habitat recovery by plants other than pioneer species will occur in the short-term. There is also potential for changes to the vegetation community composition to occur through the introduction or spread of invasive plant species, often themselves pioneer-type species. The magnitude of the impact on ecosystem composition is medium because, prior to any mitigation, the potential for changes to ecosystem composition potentially brought about by edge effects is possible. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction phase of the Project, pre-mitigation (Table 28). Following the application of appropriate mitigation measures (rehabilitation of the soil and vegetation), vegetation recovery to primary levels of succession would be expected in the short term, with further recovery in the medium and long-term (dependent on a high degree of management and specialist intervention both during and after construction); therefore the impact significance is expected to be reduced to minor, as the magnitude will become negligible, and the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be regional because impacts are restricted to the CHAA. Impact duration will be medium-term (that is ~25 years) because impacts are expected to continue through the Project operations. The magnitude of the impact on ecosystem configuration is medium because, prior to any mitigation, the potential for changes to ecosystem configuration is possible, especially interference with wildlife movement corridors. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to ecosystem composition during the construction phase of the Project, pre-mitigation (Table 28). Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, because the burial and rehabilitation of the pipeline will restore habitat connectivity for fauna using the escarpment corridor.

**Table 28: Potential impacts in the construction phase to the vegetation communities of the escarpment**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low - 2	Medium – 3	Moderate – 6	Negligible – 1	Medium – 3	Minor - 3
Ecosystem composition	Medium – 3	Medium – 3	Moderate – 9	Negligible – 1	Medium – 3	Minor - 3
Ecosystem configuration	Negligible – 1	Medium – 3	Minor - 3	Negligible – 1	Medium – 3	Minor - 3

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential impacts of the Feeder pipeline to the vegetation communities of the escarpment, and the wildlife corridors of which they form part. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain. Indications are that they will increase in importance (Ayebare et al. 2013), provided human pressures do not overwhelm them.





The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 7.1.2). The actual extent of possible habitat may have been over-estimated.

**11.1.1.3 What impact could the construction of the Feeder Pipeline have on wetlands and riparian habitat in the CHAA?**

**Impact Indicators**

Indicators used to assess impacts of the Feeder Pipeline on the habitat and ecosystem integrity of the wetlands of the CHAA were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

Loss of habitat due to direct disturbance and clearing associated with the Feeder Pipeline was quantified by overlaying the current, baseline extent of the vegetation communities with the Feeder Pipeline footprint, plus a 30m construction right of way.

Additional, indirect impacts to habitat were estimated by applying a 0.50 km buffer to the Feeder Pipeline footprint. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, changes in water quantity and quality (drivers of ecosystem processes and functions), and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur, and the disruption of known corridors. This was accomplished by examining available scientific literature about the ecology of wetlands (permanent and seasonally flooded grasslands).

**Impact Assessment**

■ Representativeness

The CHAA supports approximately 1157.9 ha of wetlands, of which 85.3 ha are classified as seasonal. In addition, 840 ha of riparian vegetation communities associated with drainage lines and riparian systems occur within the CHAA. Table 29 quantifies the impact of the pipeline on the CHAA wetlands and riparian habitats, based on the area of physical disturbance during construction.

**Table 29: Area of wetland and riparian habitat impacted along the 30m wide pipeline construction right of way**

Habitat	Area in the CHAA (ha)	Area in the 30m-wide pipeline corridor (ha)
Cultivation and Settlement	31860.9	108.9
Dense Bushland	1097.6	1.6
Dense Wooded Grassland	613.2	1.7
Open Bushland and Shrubland	2896.3	14.9
Open Grassland	568.5	3.1
Wooded Grassland	184.3	3.9
Open Wooded Grassland	1900.9	0.8
Riverine Woodland and Bushland (riparian habitat along drainage lines)	74.8	0.0
Riverine Bushland (riparian habitat along drainage lines)	640.3	1.6
Seasonal Wetland	85.3	0.6
Permanent Wetland	83.8	0.0
Wetland	1072.6	0.4





Approximately 2.6 ha of wetlands, seasonal wetlands and riparian habitat along drainage lines will be directly affected, which is 0.13% of the total area of similar wetland and riparian habitat in the CHAA. Further indirect impacts could result from edge effects and channel incision, affecting downstream wetland function. The loss of this quantity of wetlands in relation to their representation in the wider CHAA is not substantial; however, because this loss is brought about by the construction of a linear feature, there is a potential for the downstream wetland habitat to be affected if proper management controls are not implemented.

Impacts from the changes to representation of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Wetland and Riparian Habitat Structure and Ecosystem Composition

The construction of linear infrastructure, such as pipelines, through wetland communities is known to have detrimental impacts on the functioning, processes and species composition of these communities (Roise et al. 2004). Pipeline construction may result in significant loss of biodiversity at both local and regional scales due to restricted movement between populations, increased mortality, habitat fragmentation and edge effects, invasion by exotic species, or increased human access to wildlife habitats, all of which are expected to increase local extinction rates or decrease local recolonisation rates (Findlay and Bourdages 2000). Consequently, it could be expected that the construction of the pipeline through wetlands and riparian habitats, could alter the ecosystem processes and functions driving these wetlands, especially downstream.

Buried pipelines crossing rivers and wetlands will not (of themselves) obstruct surface water flow. The risk to wetland function is mainly due to the disruption of wetland vegetation and soils by heavy machinery, particularly when tracked vehicles are used that have greater impact on soil structure and the soil profile is overturned due to careless construction management. Disruption of flows and ecosystem composition may also occur if fill material is imported into the wetland to provide stability for excavators and pipe layers, and is not completely removed and replaced with the natural soils after construction.

Alteration of wetland and riparian vegetation community composition could occur if flow paths for water, both surface and sub-surface, are not maintained. If these flow paths are not maintained, there is a potential that wetland vegetation community composition downstream of the obstruction could become altered – typically via flow concentration and channel incision in downstream wetlands that were previously supplied by diffuse, dispersed flows. The risk of incised drainage and associated loss of wetland function due to concentration of water flows is high. The vegetation within the seasonally flooded grasslands and riparian habitats is adapted to seasonal inundation, and, therefore, is dependent upon that cycle of wet and dry for survival. Additional, associated impacts that could occur in tandem with channel incision and wetland desiccation include exotic species invasion.

As part of standard construction methods, culverted wetland/drainage line crossings have been installed as part of the in-field road and airstrip construction. Nevertheless, even with such measures in place, there is still a potential for changes to the structure and ecosystem composition of wetland and riparian habitats to occur.

The wetland and riparian vegetation communities on the escarpment are already under pressure from livestock grazing, and harvesting of fibre for house construction. It is possible that these communities may change in the long-term as grazing pressure increases, and the human population of the region increases. These changes could alter the habitat structure and composition, which, in turn, could affect the utilisation of these habitats by the currently resident species guilds.

Impacts on wetland and riparian fauna will depend on the changes in vegetation communities as a result of construction; in addition, the noise and sensory disturbances created by the construction equipment could alter the behaviour of species frequenting the wetlands during construction/decommissioning. For example, wading birds and Grey Crowned Cranes are unlikely to tolerate construction nuisance within 500 m and could avoid these areas entirely during the construction period. If construction occurs during the breeding season of the Grey Crowned Crane (that is, the dry season (Archibald et al. 2013)), which could occur in wetlands on the escarpment, then disturbance from the construction activities could cause nest abandonment (Strasser et al. 2013).



Impacts to ecosystem composition of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Water Quality and Ecosystem Composition

Impacts to ecosystem composition of the wetlands in the CHAA are predicted to be beyond the expected range of natural disturbance perturbations.

#### *Erosion and Sedimentation*

The construction of the Feeder pipeline could cause increased erosion and sediment-laden run-off to report to the wetlands and riparian habitats being crossed by the pipeline. With the construction of the pipeline, and the consequent exposure of areas of soil, there is a potential that, without adequate erosion and sediment control measures in place, sediment loads within the watercourses draining the pipeline footprint could increase. Hence, there is a potential for increased sediment loads in the wetland and riparian habitats. The watercourses and associated wetlands draining the LSA support dense emergent vegetation (see Section 6.1.1). Such vegetation can form an impactful filter for most sediment (IECA 2008); therefore, it can be expected that sediment loads reporting to downstream wetland and riparian habitats could be minimal. Nevertheless, if sediment loads are substantial, there is a potential for that sediment to smother wetland vegetation and interfere with aquatic invertebrates. If this occurs, it could detrimentally affect the wetland and riparian processes and functions, which, in turn, could alter wetland and riparian community composition, albeit on a localised scale.

#### *Hydrocarbon and Chemical Spills*

Accidental spillage of fuels and chemicals during the construction of the Feeder pipeline are probable, where most construction activities will take place over a period of 10-12 months. Certain invertebrate species (for example, aquatic snails (Araujo et al. 2012), mayflies (Savić et al. 2011)) and juvenile fish (for example, Agamy 2013) are highly sensitive to chemical pollutants, particularly hydrocarbons; therefore, accidental spillages during construction could detrimentally affect aquatic species as well as the wetland and riparian habitat processes and functions, which, in turn, could alter ecosystem composition, albeit on a localised scale.

#### *Sewage Disposal*

Sewage at the personnel camp will be treated by a package sewage treatment plant, designed to comply with the Ugandan requirements for effluent disposal and the IFC guidelines for domestic wastewater. Treated effluent will be discharged into a soakaway. The personnel construction camp is situated more than 500 m from the nearest drainage line/riparian habitat, and since the discharge will be to a soakaway, potential impacts are discussed under vol 4a Groundwater.

#### *Increase in Population*

Indirect impacts on wetlands and riparian habitats as a result of water quality impacts could occur as a result of the migration of people into the CHAA in search of work. Existing sanitary conditions in the CHAA are poor, with all of the streams being contaminated with faecal waste from both animals and humans (Chapter 07 & Vol 4c Social IA). Increasing population pressures will exacerbate these conditions. Increased grazing pressure and erosion from denuded areas around expanding settlements will increase erosion and sedimentation in the wetlands.

### ■ Ecosystem configuration

Linear infrastructure such as pipelines can present significant barriers to, or can alter behaviours of, a range of wetland and riparian wildlife, from: amphibians (for example, Pontoppidan et al. 2013); to turtles (for example, Langen et al. 2012). The construction of the pipeline through the wetlands and riparian habitats of the CHAA could present a barrier to movement for those species inhabiting them, during those project phases. However, the pipeline will be buried, therefore once construction is complete, they are not expected to be major barriers.



The pipeline will cut currently contiguous wetlands in the LSA. If not managed correctly during the construction process, that is, the installation of appropriate drainage connections, the pipeline could cause permanent barriers between the two newly separated wetland habitats. The process of clearing the wetlands for the construction of pipeline will create edge impacts, and result in the fragmentation of the wetland habitats. As already discussed, vegetation clearing creates edges or boundaries where habitat meets a disturbance. These edges open up habitat in areas where it was previously continuous, and this generally changes the abiotic conditions (for example, temperature, light, and moisture regimes) (Porensky and Young 2013). Edges also often result in changes in species composition along the edge, with the edges typically becoming dominated by pioneer and weedy species adapted to the particular microclimate experienced on the edge (Porensky and Young 2013). Fragmentation of wetland habitat is known to reduce the viability of many species and the wetland as a whole, with the viability of the particular fragment dependent on its size, proximity and, hence, connectivity to other wetland habitats (Uzarski et al. 2009). These wetland communities are already under pressure from livestock grazing, and harvesting of fibre for house construction.

Impacts to ecosystem configuration of the wetlands and riparian habitats in the CHAA are therefore predicted to be beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The sensitivity of wetlands and riparian habitats is high, because permanent and seasonal wetlands are important breeding and foraging habitat for Grey Crowned Crane, a species of concern; and wetlands are susceptible to degradation as a result of changes in water flow patterns brought about by infrastructural developments. The habitats are also already under stress from livestock grazing and harvesting of fibre. This high sensitivity is weighed against the magnitude of each of the impacted indicators as described in the paragraphs below, in order to derive the overall impact level for each indicator.

#### ■ Representativeness

Impacts to the representativeness of the habitat will be adverse. The geographical extent of impacts will be local because impacts are restricted to the Feeder pipeline construction right of way on the escarpment, and will only constitute loss of approximately 0.13% of wetland and riparian habitat in relation to the CHAA. The magnitude of the impacts of the Feeder pipeline on representativeness is therefore considered to be low. In the context of high wetland sensitivity, this results in impacts of moderate significance.

Following the application of appropriate mitigation measures, the impact significance is expected to be reduced to minor, as rehabilitated wetlands re-establish following completion of construction activities.

#### ■ Wetland Structure and Ecosystem Composition

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be medium-term, because impacts are expected last as long as it takes the wetland rehabilitation to become established post-construction. The magnitude of the impact on ecosystem composition is low because, although changes to wetland structure and ecosystem composition in wetland areas crossed by the Feeder pipeline are likely, which could result in edge effects, changed flow regimes, and erosion and sedimentation of affected wetlands, the extent of the area affected is relatively small. Therefore, the magnitude and high sensitivity of the wetlands combine to produce a moderate overall impact level to ecosystem composition during construction of the Feeder pipeline, pre-mitigation (Table 30).

Following the application of site-specific mitigation measures (Section 12.2), including appropriate construction management methods, the impact significance is expected to be reduced to minor, because, as the magnitude will become negligible, the sensitivity of the receptor will remain the same.

#### ■ Water Quality and Ecosystem Composition

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be for the duration of construction, which is expected to take approximately 3 years. The magnitude of the different project-specific impacts on ecosystem composition varies, as discussed under each of the subheadings below.





### *Erosion and Sedimentation*

Clearing of vegetation will result in increased sediment loads in stormwater flows. These impacts will be of short duration, most likely occurring as a result of significant rain events, and will be most frequent in the early stages of construction when bulk earthworks are in progress and large areas of exposed earth are available. Impacts will be local (mainly in the catchment of wetlands that will be crossed by the pipeline). They will generally be reversible, and, consequently, have a low magnitude in the context of the Feeder pipeline construction. Wetland and riparian plant species are generally tolerant of occasional increases in sediment load in stormwater flows, and can serve as an effective sediment filter. In cases of severe and/or ongoing sediment loading, detrimental impacts on wetland and riparian vegetation and macro-invertebrates could be likely, which could detrimentally affect wetland processes and functions and, in turn, wetland composition at a localised scale. The high wetland and riparian habitat sensitivity, in combination with the low magnitude of predicted effects, will result in a moderate overall impact level to ecosystem composition during construction of the Project, pre-mitigation (Table 30).

Following the application of appropriate mitigation measures, the impact significance is expected to reduce to minor.

### *Hydrocarbon and Chemical Spills*

The baseline concentrations of hydrocarbons and other industrial pollutants in the wetlands and riparian habitats of the LSA were below levels that cause harm in the aquatic environment. Some invertebrate species (such as aquatic snails, described in Araujo et al. 2012), mayflies (Savić et al. 2011) and juvenile fish (Agamy 2013) are particularly sensitive to these pollutants.

Equipment working in the wetlands could contribute small quantities of oil into the aquatic environment as a result of leakages or spills. Hydrocarbon spills will only occur if insufficient care is taken during construction to prevent them, and will have a low probability of occurrence. In the absence of daily monitoring and management of site activities by competent personnel, an impact of low magnitude could occur, resulting in an overall impact of moderate significance (Table 30).

Following the application of appropriate mitigation measures, the impact magnitude is expected to reduce to negligible, which in combination with the high sensitivity of the wetland habitat, results in an overall impact of minor significance.

### *Increase in Population*

In the absence of project interventions, the impact of increased population density (and associated increases in grazing livestock) on the Buhuka Flats is expected to be long-term, and to have a material effect on water quality in the wetlands across the Flats through reductions in sanitary water quality, exacerbation of wetland erosion, increased harvest of plant species used for traditional home construction, increased fire frequency and increased grazing pressure. The effects will be irreversible, and of medium magnitude. In the context of the high wetland sensitivity, the predicted impact significance is major, prior to mitigation (Table 30).

Following the application of appropriate mitigation measures, including appropriate construction management methods, the impact significance associated with population increase is expected to remain moderate, as although the magnitude will become low, the sensitivity of the receptor will remain the same.

### ■ **Ecosystem Configuration**

Impacts to the ecosystem configuration will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be short-term and reversible, subject to the avoidance of permanent structural damage at the crossing points and the length of time that the rehabilitated crossing points (wetlands and riparian bank vegetation) takes to successfully re-establish. Taking into consideration the very localised geographic extent of construction through the wetlands and small drainage lines and the intensive cultivation in all of the surrounding areas, as well as the fact that the pipeline will be buried, the magnitude of impact is considered to be low.



Combined with the high sensitivity of wetland and aquatic environments, the overall impact on wetland and drainage line configuration, without mitigation, is expected to be of moderate significance. This suggests that project-specific measures will be necessary to minimize the impacts of construction equipment operating in the wetlands and drainage lines.

Following the application of project-specific mitigation measures (Section 11.2), the impact significance is expected to be reduced to minor (Table 30).

**Impact Significance Rating**

**Table 30: Potential impacts of construction of the Feeder pipeline on the wetlands of the CHAA**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low - 2	High - 4	Moderate - 8	Negligible - 1	High - 4	Minor - 4
Ecosystem composition – wetland structure	Low - 2	High - 4	Moderate - 8	Negligible - 1	High - 4	Minor - 4
Water quality						
Erosion and sedimentation	Low - 2	High - 4	Moderate - 8	Negligible - 1	High - 4	Minor - 4
Hydrocarbon and chemical spills	Low - 2	High - 4	Moderate - 8	Negligible - 1	High - 4	Minor - 4
Population increase	Medium - 3	High - 4	Major - 12	Low - 2	High - 4	Moderate - 8
Ecosystem configuration	Low - 2	High - 4	Moderate - 8	Negligible - 1	High - 4	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Feeder pipeline impacts to the wetlands of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**11.1.1.4 What impact could the construction of the Feeder Pipeline have on Bugoma CFR?**

**Impact Indicators**

Bugoma Central Forest reserve (Bugoma CFR) will not be directly affected by pipeline construction, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary. The indirect impacts of population influx, due to expectations of work on the construction contract, are considered since this could impact on the forest reserve.

A formal impact classification based on indicators was developed for induced and cumulative impacts to the Bugoma Central Forest Reserve; the impacts are discussed, and their significance assessed through a **reasoned narrative**. An overall impact significance classification was then developed. This was





accomplished by examining available literature about the ecology of the Bugoma Central Forest Reserve (BCFR), and scientific literature regarding the effects of migration and human population pressure on forests in Africa.

The impacts were assessed in light of the guidance provided by IFC (2013), and in consideration of other known projects being developed in the wider area. In particular, the development of the oil processing facility at Kabaale, the oil developments around the Kaiso-Tonya area (AECOM 2012), the Hoima-Mputa-Fort Portal-Nkenda power line, and the potential for regional population increases in the wider area.

### **Impact Analysis**

Bugoma CFR is identified as a valued component for this impact assessment, certainly in terms of biodiversity (see Section 6.3.1.1.4). As identified in that section, apart from being one of the last stands of tropical semi-deciduous forests in the region, it also supports known populations of the Endangered Eastern Chimpanzee and range-restricted Nahan's Francolin (Plumptre et al. 2011), potential non-breeding habitat for the Endangered Madagascar Pond Heron (see Section 6.3.3.1), as well as elephants and a host of other threatened and irreplaceable species.

Although Bugoma CFR will not be directly affected by pipeline construction, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary, the indirect effects of project-induced migration to the pipeline construction region due to expectations of work on the construction contract have the potential to affect the ecological integrity of the habitats within the CFR, as well as the species of conservation concern that occur within it.

Bugoma CFR is under severe pressure from the human population surrounding it. In 2013, more than 1,500 people evicted illegally after settling within the forest (Mugerwa 2013), and recent (2016) reports from the area suggest that land-grab activity has resulted in the conversion of an estimated 8000 ha for sugar cane plantation development. The land cover study (see Figure 4) clearly shows that the areas surrounding the forest have largely been transformed for agricultural and subsistence purposes, and the transformation is now encroaching within the boundary of the CFR. This trend of encroachment of protected areas is not unique. Protected areas are known to be particularly vulnerable to changes in human demographics and deforestation; and Wittemyer et al. (2008) identified that human population growth and encroachment around protected areas is significantly higher than the average population growth in rural areas. This difference was largely due to the immigration of people into these areas because of the perceived increased availability of opportunities, natural resources and potential jobs (Wittemyer et al. 2008). This is supported by research conducted in the forests of the Albertine Graben, and the wider CHAA. For example, in a study of Budongo Central Forest Reserve, Zommers and MacDonald (2012), identified that of the local communities that hunted bush meat in the forest, nearly 73% were immigrants to the area. Furthermore, these workers identified that the households of immigrants were also more likely to be involved with deforestation.

The results of the recently-updated (Nov 2017) social baseline data gathering fieldwork indicated that immigration to the CHAA has escalated significantly since the opening of the escarpment road, and commencement of preliminary construction works at the Kingfisher Field Development Area.

An Influx Management Strategy and Framework Plan was developed to manage the potential influx of people into the LSA. However, this plan can only focus on those measures over which CNOOC has control, and to support the range of government and donor projects in Uganda aimed at socio-economic development and environmental conservation. How this translates to the potential influx of people towards the Feeder pipeline route and construction camp is unknown.

### **Impact Classification**

Indirect impacts on Bugoma CFR due to population influx to the region during construction of the Feeder pipeline will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR, in the CHAA. Impact duration will be short-term (that is, limited to the construction phase of the Feeder pipeline, that is, 10-12 months). The magnitude of the effects of population influx associated with pipeline construction on the Bugoma Central Forest Reserve is expected to be low, given the short duration of the construction period, and the distance of the construction activities from the Bugoma CFR boundary.





The sensitivity of the Bugoma Central Forest Reserve is high because it is a threatened ecosystem that is already under pressure. Therefore, the intensity and sensitivity combine to produce a moderate overall impact level during the construction phase of the pipeline, prior to the implementation of strict Project-specific mitigation measures (**Table 31**).

The mitigation hierarchy is an important process that has been used to minimise impacts to Bugoma CFR. The focus for the continued use of the mitigation hierarchy during the Feeder pipeline construction phase will be continued development and implementation of mitigation measures through monitoring and adaptive management (see Section 12.0).

With the implementation of such mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude of impact could become negligible, and the sensitivity of the receptor will remain the same (**Table 31**).

**Table 31: Potential impacts in the construction phase to the Bugoma Central Forest Reserve**

	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat and ecosystem integrity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4

As noted above, the focus for reducing impacts to the Bugoma Central Forest Reserve is mitigation to lessen various types of disturbance that may occur (Section 12.0). Optimising mitigation will reduce residual impacts on the CFR to minor, and reduce the need for offsetting. However, possible offsetting options to ultimately achieve ‘net gain’ for the natural habitats of the forest have been identified and are discussed in Section 13.0.





### 11.1.2 What impact could the construction of the Feeder Pipeline have on species of concern?

This section presents the assessment of impacts that the construction of the Feeder Pipeline could have on those species of concern that trigger critical habitat, as identified in Section 8.0; that is, the Mud Snail (*G. candida*), Nahan's Francolin, and Eastern Chimpanzee; as well as Grey Crowned Crane which was also included as a valued component (see Table 25). Potential impacts to other species of concern are assessed at the habitat level (see Section 11.1.1).

Only Mud Snail and Grey Crowned Crane are potentially directly affected by construction of the Feeder pipeline. Other species, including the Eastern Chimpanzee and Nahan's Francolin, occur in the Bugoma Central Forest reserve (BCFR) but are not directly affected by pipeline construction, which at its nearest point will be at least 1,8 km from the BCFR boundary. The indirect impacts of population influx, due to expectations of work on the construction contract, are considered since this could impact on the forest reserve and the species of concern that it supports.

#### 11.1.2.1 What impact could the construction of the Feeder Pipeline have on the Mud Snail?

The Mud Snail (*Gabbiella candida*) is a Critically Endangered and range restricted species. Currently, the only known populations occur around Butiaba (see Section 9.1.1.1), which is on the eastern shore of Lake Albert approximately 90 km north of the LSA. Although this species was not confirmed within the CHAA during the course of baseline studies, there is a potential that this species could occur in the near-shore habitats of the CHAA, based on its known habitat preferences, and those of other Mud Snail species (*Gabbiella* spp.), which have previously been recorded in both the same locality as this species at Butiaba, as well as in the LSA. Hence, a precautionary approach has been adopted, and *G. candida* is assumed to occur in the near-shore habitats of the CHAA.

#### Impact Indicators

No habitat loss due to direct disturbance and clearing of habitat is expected during construction of the Feeder pipeline, as these habitats lie outside that portion of the LSA. It is unlikely that construction of the Feeder pipeline could alter the habitat connectivity of the near-shore habitats in the CHAA. No structures are being put in place that will alter the natural connectivity of the aquatic habitats of the lake. It is expected that the connectivity amongst the aquatic habitats will remain the same during construction as they were during baseline.

Predicted construction phase impacts of the Feeder pipeline on the Mud Snail were therefore limited to the proposed discharge of hydrotest water to Lake Albert and subsequent effects on habitat quality.

The assessment of changes to habitat quantity and quality was focussed on the prediction of changes to water quality in the water column from construction-related activities.

#### Impact Assessment

##### ■ Habitat Quality and Quantity

Very little information is available for this species. Information about the genus *Gabbiella* indicates that this group is generally found in lakes, rivers and, less commonly, in small water bodies, and rarely in habitats that dry out (Kristensen and Stensgaard 2010). Two sister species (*G. humerosa* and *G. walleri*) were recorded in the LSA (see Section 3.3.2.4, Appendix D). These two species have also been recorded in the same locality and habitat as *G. candida* (GBIF, 2014). Therefore, the habitat preferences of those two species were used as a model to predict the potential impacts of changes to habitat quantity and quality for this species.

The two sister Mud Snail species recorded from the CHAA were collected from bottom substrates in the open sandy shore habitats in the LSA. These habitats are characterised by a gently sloping lake bed extending from the shore line to deeper water. The substrate is typically comprised of sand and finer sediments (Wandera and Balirwa 2010). This habitat constitutes approximately linear (lake shore) 10.5 km of the CHAA; however it should be noted that the near-shore habitat is located well beyond the Feeder pipeline construction right-of-way.





Despite the large buffering capacity of Lake Albert, the proposed direct discharge of the Feeder pipeline hydrotest water to Lake Albert could ultimately report to the near-shore habitats of the CHAA, with subsequent effects on the Mud Snail.

Aquatic snails are highly sensitive to chemical pollutants, particularly hydrocarbons (Araujo et al. 2012). Currently, the concentrations of hydrocarbons and other pollutants in the lake waters of the near-shore habitats are below levels that could cause harm to the environment; this is supported by the healthy aquatic communities that were observed in the lake’s near-shore habitats at baseline (see Section 6.2). As such, the direct discharge of the hydrotest fluid to Lake Albert could have a detrimental impact on the near-shore habitats of the LSA through the introduction of toxic compounds and pollutants, and subsequent effects on the Mud Snail.

Impacts on habitat quality and quantity for *G. candida* arising from discharge of the Feeder pipeline hydrotest fluid following construction are predicted to be beyond the expected range of natural disturbance perturbations. Potential loss of critical habitat for the Mud Snail requires the consideration of offsets to meet IFC requirements.

**Impact Classification**

The Mud Snail’s sensitivity is high because this species is Critically Endangered, and potentially triggers a Tier 1 critical habitat designation.

Impacts to the Mud Snail’s habitat quantity and quality will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Although the impact duration will be short-term, and should be reversible with time as the Lake waters dilute and disperse the contaminants, the magnitude of the impacts of a potentially-toxic discharge to *G. candida*’s habitat could be high. Therefore, the magnitude and sensitivity combine to produce a major overall impact on habitat quality and quantity as a result of contamination, prior to the implementation of the recommended mitigation (Table 32).

The strict application of the recommended mitigation measures (Section 12.2) will reduce the risk, and thus the likely magnitude, of potential habitat contamination for the Mud Snail, reducing the overall impact significance to moderate.

**Table 32: Potential impacts in the construction phase to the Mud Snail**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	High – 4	High – 4	Major - 16	Negligible - 1	High - 4	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts to the near-shore habitats of the CHAA, and hence potential local populations of Mud Snail. However, there is some uncertainty with regard to this species actually occurring in the CHAA. As such, the above assessment has been undertaken based on a precautionary approach.

The spatial extent of the near-shore habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**11.1.2.2 What impact could the construction of the Feeder Pipeline have on Grey Crowned Crane?**

The Grey Crowned Crane is an Endangered species. Up to 14 individuals were regularly recorded on the Buhuka flats during baseline surveys, and a pair of Grey Crowned Crane was observed on the Flats near Well





Pad 1 during social baseline survey work conducted in November 2017. The local population of Grey Crowned Crane is regarded as highly sensitive, and the wetlands of the CHAA are considered an important conservation unit for the species.

The construction of the Feeder Pipeline could directly impact Grey Crowned Crane breeding and foraging habitat, given that construction impacts on wetlands above the escarpment are expected (Section 10.1.1.2).

The potential direct, indirect and induced impacts of the construction of the Feeder Pipeline on Grey Crowned Crane are presented below.

### **Impact Indicators**

Indicators used to assess effects of the construction of the Feeder Pipeline on Grey Crowned Crane were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Habitat loss due to direct disturbance and clearing of habitat was assessed by calculating the loss of suitable habitat from the CHAA as a result of the construction of the Feeder Pipeline, consisting of a 30m construction right of way. Changes to habitat quality due to indirect disturbance were estimated by applying a 200m buffer to the pipeline footprint. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality caused by indirect disturbance arising from light, noise, vibration, and edge effects.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat for Grey Crowned Crane in relation to the Feeder Pipeline to qualitatively identify areas where habitat becomes fragmented.

Potential changes in abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances (that is, vehicle traffic, light, noise, vibration) and site clearing activities. These disturbances were considered in light of known or inferred effects to the survival and reproduction of Grey Crowned Crane, where data on these types of effects are available.

### **Impact Assessment**

#### ■ Habitat Quality and Quantity

Grey Crowned Crane is distributed across eastern and southern Africa. Populations in many areas including Uganda have experienced rapid declines during the past 45 years (BirdLife International, 2018) for reasons including habitat loss to farming, human presence causing disturbance, collection of chicks for domestication, disruption of breeding activity, loss of roosting sites (large trees located remotely from areas frequented by humans) and dry-season fires in wetland habitats (Olupot, 2014). Its habitat preferences are diverse, including wetlands with tall emergent vegetation, open riverine woodland, shallowly flooded plains and temporary pools with adjacent grasslands, open savannas, croplands, pastures, fallow fields and irrigated areas (Archibald et al, 2018). It prefers short to medium height open grasslands adjacent to wetlands for foraging, which is consistent with the seasonally flooded grassland wetlands in the Buhuka Flats. For breeding, it prefers marshes with water 1 m deep and emergent vegetation 1 m above the water (Archibald et al, 2018); habitat which corresponds to the permanent wetlands of the Buhuka Flats. They are also often found foraging in agricultural land wherever available, in close proximity to the preferred mixture of wetlands and open grassland or savanna (Morrison, 2015).

Approximately 2.6 ha of wetlands, seasonal wetlands and riparian habitat along drainage lines will be directly affected by the construction of the pipeline, which is 0.13% of the total area of similar wetland and riparian habitat in the CHAA. In the context of available foraging and breeding habitat in the CHAA for Grey Crowned Crane, the direct losses are not substantial.

Although Grey Crowned Crane can tolerate a degree of anthropogenic disturbance in foraging habitat (e.g. in agricultural lands), in Ugandan wetlands it has been observed to be intolerant of human proximity within 100-200m (Olupot, 2014), flying away on approach; a factor which also affects breeding success as breeding birds flush from nests on approach, causing increased rates of predation, reduced time at the nest (either incubating or feeding), and ultimately nest abandonment. How tolerant foraging/roosting/breeding Grey Crowned Crane





may be to indirect disturbances, such as noise, light, vibration and edge effects, is not known. The application of a 200m buffer around the 120 m length of the Feeder Pipeline that intercepts wetlands indicates that approximately 2.4 ha of wetland habitat will be reduced in quality as a result of sensory disturbance.

This equates to a total potential habitat loss in the CHAA from direct losses from vegetation clearing, and indirect losses from sensory disturbances and edge effects of up to 5 ha (0.43%).

Effects from loss of habitat are predicted to be beyond the expected range of natural disturbance perturbations (for example, fire), although not beyond the human-induced rate of degradation via cattle grazing pressure etc.

### ■ Habitat Connectivity

The impact of the Feeder pipeline construction as a barrier to the movement of Grey Crowned Crane is likely to be adverse. The construction activity will create sensory disturbances in the short-term, which can elicit reduced use or complete avoidance of affected areas, thereby creating temporary movement barriers (for example, see Kolowski and Alonso 2009, Gleeson and Gleeson 2012). Construction of linear infrastructure (pipelines and service roads) through wetlands has the potential to create temporary barriers to movement as a result of the associated disturbance due to human presence in the area.

Impacts on habitat connectivity are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

Up to 14 individuals of Grey Crowned Crane were frequently observed on the Flats during the baseline fieldwork conducted in 2014, and a pair were observed in the same area during social baseline fieldwork conducted in November 2017. Grey Crowned Crane has potential to nest in wetland habitats and forage in the wetland/grassland/cultivated lands mosaic that occurs throughout the CHAA.

For the purposes of this assessment, it is assumed that Grey Crowned Crane is present in suitable habitat (permanent and seasonal wetlands) in the CHAA. The construction of the Feeder Pipeline is likely to generate increased levels of sensory disturbance in the locality, with resultant effects on habitat quality and the distribution of the species. Although significant losses of preferred seasonal grassland-type foraging habitat to the Feeder Pipeline construction right-of-way could affect the abundance and distribution of Grey Crowned Crane in the CHAA; the predicted potential habitat loss amounts to 5 ha or (0.43%) of similar habitat in the CHAA, which is miniscule in the context of the available foraging habitat for cranes, which includes wetlands and agricultural lands.

Effects on the abundance and distribution of Grey Crowned Crane are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

During the baseline fieldwork conducted in May 2014, the formation of breeding pairs of Grey Crowned Crane was indicated. Grey Crowned Crane nests are typically constructed within or on the edges of marshes with water of 1m depth and emergent vegetation 1m high (BirdLife International, 2018). Loss and disturbance of wetland habitat in the CHAA is expected to decrease the likelihood of Grey Crowned Crane selecting these areas for breeding.

Above the escarpment, migration into the area due to expectations about work on the construction projects could result in increased settlement in the CHAA. In the vicinity of the areas of permanent wetland, this could cause greater pressure on natural resources, including harvesting of reeds and an increased threat of discovery and removal of crane chicks for sale/domestication. Direct disturbance via human proximity within 100-200m causes breeding birds to flush from nests on approach (Olupot, 2014), which may result in increased rates of predation, reduced time at the nest, either incubating or feeding, and ultimately nest abandonment, affecting reproductive success. It is assumed for this assessment that indirect disturbances arising from mechanical noise, site lighting, vibration and edge effects during construction are also likely to affect the breeding success of Grey Crowned Crane in the CHAA.



Removal of large trees which have importance as night-time roosts and day-time shelter from the midday sun has been indicated in localised declines in Grey Crowned Crane in Uganda (Olupot, 2014); the loss of such trees during vegetation clearance for pipeline construction is therefore expected to have implications for the survival of flocks and individuals in the area.

These effects to the survival and reproductive success of Grey Crowned Crane are predicted to be well beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Grey Crowned Crane sensitivity is high because it is Endangered.

Impacts to Grey Crowned Crane habitat quantity and quality will be adverse. The geographical extent of impacts will be local because effects are restricted to those areas within 200 m of the Production Facility infrastructure footprint, with approximately 0.36 ha (0.5%) of potential habitat being cleared, and 2.4 ha (3.47 %) of potential habitat indirectly affected. Impact duration will be short-term (that is, limited to the construction phase of the Project, approx. 10-12 months). Changes to the habitat quality and quantity from sensory disturbances arising from construction activities are expected to be reversible after completion of the works. The magnitude of the effects of construction on Grey Crowned Crane habitat quantity and quality is considered low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, because the magnitude of effects could be reduced to negligible, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to Grey Crowned Crane habitat connectivity will be adverse. The geographical extent of impacts will be local because effects are restricted to the 30m construction right-of-way within which wetlands will be crossed by the Feeder pipeline. Impact duration will be short-term, as the pipeline will be buried and vegetation rehabilitated, and sensory disturbances arising from construction activities are expected to dissipate within a short time after completion of the construction phase. The magnitude of the effects of construction on Grey Crowned Crane habitat connectivity is considered to be low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the construction phase of the Feeder Pipeline, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, because the magnitude of effects could be reduced to negligible, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to Grey Crowned Crane abundance and distribution will be adverse. The geographical extent of impacts will be local because effects are restricted to wetlands within 200m of the proposed Feeder pipeline construction right of way. Impact duration will be short-term (that is, limited to the construction phase of 10-12 months) because physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities. There is a possibility that individuals could be killed or severely disturbed during the construction phase (e.g. chicks on nests during clearance works); however, in the context of overall species abundance, these losses or disturbances are expected to be reversible after completion of the works. The magnitude of the effects of construction on Grey Crowned Crane abundance and distribution is thus considered low. Therefore, the magnitude and sensitivity combine to produce an impact of moderate significance during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor, because the magnitude of effects could be reduced to negligible, and the sensitivity of the receptor will remain the same (Table 14).

Impacts to the survival and reproduction of Grey Crowned Crane will be adverse. The geographical extent of impacts will be local because effects are restricted to wetlands within 200m of the proposed Feeder pipeline construction right of way. Although physical and sensory disturbances arising from construction activities are expected to dissipate within a short time after cessation of activities, the effects of in-migration on Grey Crowned Crane are expected to be permanent, once households/settlements are established. The likelihood of impacts is uncertain (it is not known whether there are any breeding pairs of the crane in the permanent wetlands of the CHAA above the escarpment and the extent of migration into the area cannot be determined with certainty). A threshold of 10% for this species' survival and reproduction in the CHAA is reasonable, and



it is considered unlikely that this number of individuals in the CHAA will be affected. The magnitude of the effects of construction on the survival and reproduction of the Grey Crowned Crane is considered to be medium. The magnitude and sensitivity combine to produce a major overall impact level during the construction phase of the Project, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to decrease to moderate, because the magnitude of impact will be reduced to low, and the sensitivity of the receptor will remain the same (Table 33).

**Table 33: Potential construction phase impacts to Grey Crowned Crane**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate - 8	Negligible - 1	High – 4	Minor – 4
Habitat connectivity	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor – 4
Abundance and distribution	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor – 4
Survival and reproduction – direct construction effects e.g. mortality	Low – 2	High – 4	Moderate - 8	Negligible - 1	High – 4	Minor – 4
Survival and reproduction – population influx	Medium	High – 4	Major - 12	Low – 2	High – 4	Moderate - 8

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts on the Grey Crowned Crane and its habitat in the CHAA.

The spatial extent of the wetland habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.1). The actual extent of possible habitat may have been over-estimated.

**11.1.2.3 What impact could the construction of the Feeder Pipeline have on Nahan’s Francolin?**

Nahan’s Francolin is a Vulnerable and range-restricted species. It has a very restricted distribution, being found only in north-east DRC and western and south-central Uganda; in particular, the forests of Budongo, Bugoma and Mabira (McGowan and de Juana 1994). In the CHAA, it is restricted to the Bugoma Central Forest Reserve. It triggers Tier 2 critical habitat under Criterion 2 (Section 9.0, Appendix G).

Tier 2 habitats are considered to be sensitive, and, therefore, if a project is located in such a habitat, the IFC considers that compliance with the provisions of paragraph 17 of PS 6 (IFC 2012a) would be difficult. In summary, a project will not be developed in Tier 2 habitat unless: no other viable alternatives exist; and, the project does not lead to measurable and irreversible adverse impacts to the valued component that triggered critical habitat; and, the project does not lead to a net reduction in the global and/or national/regional population of the triggering species (such as Nahan’s Francolin) over a reasonable period of time; and, a robust,





appropriately designed, and long-term biodiversity monitoring and evaluation programme is part of the project's Environmental and Social Management System (ESMS). A Biodiversity Action Plan (BAP) will then be developed to achieve net gain for the affected species.

It is expected that the construction of the Feeder Pipeline will not have direct impacts to Bugoma CFR; its nearest point will be at least 1,8 km from the pipeline, and therefore no direct impacts on Nahan's Francolin are predicted. However, the indirect impacts of population influx, due to expectations of work on the construction contract, are considered since this could impact on the forest reserve and subsequently Nahan's Francolin.

### **Impact Indicators**

As the pipeline will be at least 1.8 km from the Bugoma CFR boundary, no effects on habitat connectivity are predicted. Therefore, the indicators used to assess effects of the population influx associated with the construction of the Feeder pipeline on Nahan's Francolin were: habitat quantity and quality, abundance and distribution, and survival and reproduction.

Potential changes in habitat quantity and quality, abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances arising from increased human presence in the CHAA. These disturbances were considered in light of known or inferred effects to the survival and reproduction of buttonquail species for which data on these types of effects are available.

### **Impact Assessment**

#### ■ Habitat Quality and Quantity

This species is confined to dense, mature, moist, sometimes swampy medium-altitude forest below 1,500 m (McGowan and de Juana 1994, BirdLife International 2014i); and is reasonably common in Budongo Central Forest Reserve (Plumptre et al. 2010, 2011). Suitable habitat was predicted to cover an area of ~35,201 ha (352 km<sup>2</sup>) in the CHAA, principally in the Bugoma Central Forest Reserve. No direct losses or impacts on the quantity of habitat suitable for Nahan's Francolin are predicted as a result of the construction of the Feeder pipeline. However, indirect effects on habitat quality and quantity may arise as a result of human-induced disturbance associated with population influx to the region,

The Nahan's Francolin's degree of vulnerability to disturbance is unknown. Birds are known to be sensitive to land use and habitat alteration (Lussier et al. 2006). The behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions, such as the availability of alternative foraging sites (Madsen 1998). A reduction in habitat quality and quantity may have conservation implications and consequences for the population of this species.

The construction phase of the Feeder pipeline, as a component of the Project, could result in an influx of people into the CHAA seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of habitat alteration and deforestation. Nahan's Francolin is a shy, forest-dependent species (Sande et al. 2009a) and thus is also probably sensitive to disturbance arising from deforestation/human presence in the forest; individuals may avoid the resultant sub-optimal habitat. These factors could combine to reduce the habitat quality and quantity for Nahan's Francolin.

Effects from reductions in habitat quality are predicted to be beyond the expected range of natural disturbance perturbations, although not beyond the baseline rate of direct loss as a result of the human induced deforestation rate.

#### ■ Abundance and Distribution

Although the pipeline will be at least 1.8 km from the Bugoma CFR boundary, the indirect impacts of population influx to the CHAA, due to expectations of work or opportunities generated during construction of the pipeline, could ultimately affect the abundance and distribution of Nahan's Francolin in the CHAA.





The construction of the pipeline, as a component of the overall Project, could result in an influx of people into the area seeking jobs and/or opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma CFR and within the CFR itself will exacerbate existing pressures on the forest. These pressures are likely to manifest in increased rates of bush meat hunting and habitat alteration, which could reduce the abundance and distribution of Nahan's Francolin in Bugoma CFR. Although this species within the forest is reported to be relatively common (*viz.*, Plumptre et al. 2011), the distribution of individuals within this habitat is unknown. For the purposes of this assessment, it is assumed that the current distribution of this species is evenly spread within suitable habitat in the Bugoma Central Forest Reserve.

Effects from the population influx to the CHAA associated with the construction of the pipeline are predicted to be beyond the expected range of natural disturbance perturbations, although not beyond the baseline disturbance levels from the human-induced deforestation rate.

### ■ Survival and Reproduction

The Nahan's Francolin's degree of vulnerability to direct disturbance, particularly during the breeding season, is not well understood. Nahan's Francolin is reliant upon large trees, with appropriate buttress formation, for breeding sites (Sande et al. 2009a). No direct losses or impacts on breeding habitat suitable for Nahan's Francolin are predicted as a result of the construction of the Feeder pipeline. However, indirect effects on habitat quality and quantity may arise as a result of human-induced disturbance associated with population influx to the region.

How tolerant the Nahan's Francolin may be to indirect disturbances during the breeding season, are not completely understood. What is known is that disturbance of mature forest generally appears to reduce the home range of this species (Sande et al. 2009b), and hence its potential to find suitable mates. Generally, for birds, the behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions such as the availability of alternative breeding sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the local population of Nahan's Francolin.

The construction of the pipeline, as a component of the overall Project, could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma CFR and within the CFR itself will exacerbate existing pressures on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration, which could reduce the survival and reproductive success of Nahan's Francolin. Assuming it is sensitive to sensory disturbance because it is a shy, forest-dependent species (Sande et al. 2009a), the disturbance generated by increased human presence in Bugoma CFR could further affect the breeding success of those individuals living in close proximity to the boundaries of Bugoma CFR in particular.

These effects to the survival and reproductive success of Nahan's Francolin are predicted to be well beyond the expected range of natural disturbance perturbations, although not beyond the current, baseline human-induced deforestation rate.

### **Impact Classification**

The Nahan's Francolin's sensitivity is high because it is range-restricted and triggers Tier 2 critical habitat.

Impacts to the Nahan's Francolin's habitat quantity and quality will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. The magnitude of the impact of population influx associated with the pipeline construction on the habitat quantity and quality of the Nahan's Francolin is expected to be low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the construction phase of the Feeder pipeline, pre-mitigation (**Table 34**). Following the application of appropriate mitigation measures, the impact significance is expected to decrease to minor,





primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (**Table 34**).

Impacts to the Nahan’s Francolin’s abundance and distribution could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. There is a possibility that as a result of influx during the construction phase, the distribution and abundance of Nahan’s Francolin in Bugoma CFR could be affected. The magnitude of the effects of population influx associated with pipeline construction on the abundance and distribution of the Nahan’s Francolin is Low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level, pre-mitigation (**Table 34**). Following the application of site-specific mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (**Table 34**).

Impacts to the survival and reproduction of Nahan’s Francolin could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. No direct effects on breeding habitat or direct species mortalities are anticipated as a result of pipeline construction; however indirect effects arising from sensory disturbance and poaching associated with population influx to Bugoma CFR could result in the injury, mortality and reduced breeding success of some individuals. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve will not be affected by the indirect impacts of pipeline construction. The magnitude of the effects of construction on the survival and reproduction of the Nahan’s Francolin is therefore low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the pipeline construction phase of the Project, pre-mitigation (**Table 34**).

Following the application of site-specific mitigation measures, the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same (**Table 34**).

**Table 34: Potential impacts in the construction phase to the Nahan’s Francolin**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Abundance and distribution	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4
Survival and reproduction	Low – 2	High – 4	Moderate – 8	Negligible – 1	High – 4	Minor – 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced impacts that the operation of the Feeder pipeline could have on the Nahan’s Francolin and its habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.1.1). The actual extent of possible habitat may have been over-estimated.





### 11.1.2.4 *What impact could the construction of the Feeder Pipeline have on Eastern Chimpanzee?*

The Eastern Chimpanzee is an Endangered species. The population of Eastern Chimpanzees in the Bugoma CFR is recognised as being one of the four largest in the region (Plumptre et al. 2010). Hence, the forest is recognised as an important chimpanzee conservation unit by the IUCN (Plumptre et al. 2010).

This species triggers Tier 1 critical habitat (Section 6.3.2.2, Appendix G). Tier 1 habitats are considered to be very sensitive, and, therefore, if a project is located in such a habitat, the IFC considers it unlikely that the client will be able to comply with the provision of PS 6, in particular paragraphs 17, 18 and 19 (IFC 2012a, as presented in Appendix C). In summary, a project will not be developed in Tier 1 habitat unless: no other viable alternatives exist; and, the project does not lead to measurable and irreversible adverse impacts to the valued component that triggered critical habitat; and, the project does not lead to a net reduction in the global and/or national/regional population of the triggering species (i.e. Eastern Chimpanzee) over a reasonable period of time; and, a robust, appropriately designed, and long-term biodiversity monitoring and evaluation programme is part of the project's Environmental and Social Management System (ESMS). A Biodiversity Action Plan (BAP) will be developed to achieve net gain for the affected species.

Bugoma Central Forest reserve (Bugoma CFR) will not be directly affected by pipeline construction, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary. The indirect impacts of population influx, due to expectations of work on the construction contract, are considered since this could impact on the forest reserve, and the population of Eastern Chimpanzee that lives there.

#### **Impact Indicators**

As the pipeline will be at least 1.8 km from the Bugoma CFR boundary, no effects on habitat connectivity during operation are predicted. Therefore, the indicators used to assess effects of the population influx associated with the operation of the Feeder pipeline on Eastern Chimpanzee were: habitat quantity and quality, abundance and distribution, and survival and reproduction.

Potential changes in habitat quantity and quality, and abundance and distribution, were assessed qualitatively by considering changes in disturbances arising from increased human presence in the CHAA. These disturbances were considered in light of known or inferred effects to the survival and reproduction of other populations of chimpanzees for which data on these types of effects are available.

To assess effects to survival and reproduction as a result of in-migration and potential associated increases in poaching and disease spread, in-migration rates were assessed based on the data presented in the social impact assessment (Vol 4c). A literature review of the impact of contact with humans was also conducted.

#### **Impact Assessment**

##### ■ Habitat Quality and Quantity

The entire Bugoma Central Forest Reserve is considered suitable habitat for Eastern Chimpanzees, as well as the areas beyond the boundaries of the forest reserve (see McLennan 2008). However, for the intents of this impact assessment, the habitat within and immediately surrounding the Bugoma Central Forest Reserve was only considered.

The potential for the construction of the Feeder pipeline (as a component of the overall Project) to be attractive for people seeking opportunities could result in a substantial increase the current population around the Bugoma Central Forest Reserve. An increasing human population could lead to an increase in the demand for agricultural and timber resources, resulting in increased rates of habitat alteration and deforestation, which, in turn could further fragment, change, and degrade Bugoma CFR and surrounding habitat, resulting in reduced habitat quantity and quality for Eastern Chimpanzee.

The degree of vulnerability to disturbance experienced by chimpanzees is reasonably well known (for example, see Parren and Byler 2003, Rabanal et al. 2010, Thompson and Wrangham 2013). The chimpanzees living in and around the Bugoma Central Forest Reserve most likely experience sensory disturbances at present from human activities, given the high human populations living around the reserve. Indeed, groups are known to forage in the agricultural fields surrounding the forest, and hence, would more than likely be used to human





noises and disturbances (McLennan 2008). As such, the potential sensory disturbances arising from population influx associated with the construction of the pipeline are likely to be minimal. Nevertheless, the magnitude of noise may not be the most important determinant of chimpanzee response. Instead, chimpanzees may respond to 'new' noises or may associate particular noises with other occurrences (for example, machine noise may be associated with human presence, which chimpanzees may, in turn, associate with the presence of danger). Where humans pose a threat, chimpanzees generally avoid them (Hockings and Humle 2009, Parren and Byler 2003). Therefore, the degree of avoidance may depend on the behaviour of people, resulting in varying levels of indirect effects on the quality of available habitat for Eastern Chimpanzee, and highlighting the importance of managing contractor activity during construction.

Effects from the reduced habitat quantity and quality during construction are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

The Bugoma Central Forest Reserve supports one of the top four Eastern Chimpanzee populations in Uganda, with a population of between 450 and 850 individuals (Plumptre et al. 2010). Given its distance from the forest, the construction of the Feeder pipeline is not expected to detrimentally affect the abundance or distribution of chimpanzees in the forest directly.

However, the construction of the Feeder pipeline could influence influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population both within and in the area surrounding Bugoma CFR, will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting, and reduced habitat availability due to human disturbance, which could affect the abundance and distribution of Eastern Chimpanzee within Bugoma CFR.

Effects are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

In-migration of people seeking opportunities associated with the construction of the Feeder pipeline may adversely affect survival and reproduction of chimpanzees through poaching and disease transfer. Hunting and poaching is a recognised threat to chimpanzees in Bugoma Central Forest Reserve (Plumptre et al. 2003), and this pressure could increase as in-migration of people from other areas occurs. Mitigation and management of in-migration will be very important to minimise potential effects on chimpanzees and other fauna species.

Disease is one of the major threats to Eastern Chimpanzees (Oates et al. 2008); increased abundance of people and competition for land and food resources between humans and chimpanzees could lead to higher rates of disease spread from humans to chimpanzees. Chimpanzees are closely related to humans; therefore, many diseases are transferrable between chimpanzees and humans (Formenty et al. 2003; Isabirye-Basulta and Lwanga 2008). Either direct or indirect contact with humans can spread disease. For example, there is evidence to suggest that respiratory illnesses have been transferred directly to chimpanzees from humans as a result of researcher and tourist contacts, often leading to outbreaks and death (Formenty et al. 2003). Human defecation in forest undergrowth can indirectly lead to spread of intestinal diseases, such as *Clostridium perfringens*, which can be fatal to chimpanzees (Fujita 2011). As forest fragments decrease in size, risks of contact with, and transmission of, disease from humans increases (Isabirye-Basulta and Lwanga 2008). Factors that lead to increased crop raiding or sharing of water resources can also increase the risk of disease spread (Hockings and Hulme 2009). As identified in the social impact assessment (Vol 4c), communicable respiratory diseases are a significant concern in the LSA; as such there is a risk of loss of Eastern Chimpanzee individuals due to transmission of diseases as a result of construction-phase population influx.

What effect the loss of individuals from the population could have is unknown. It is doubtful that the population could be reduced by 10% and hence, reach that critical population threshold due to population influx associated with the pipeline alone, nevertheless the potential impacts are predicted to be beyond the expected range of natural disturbance perturbations.



**Impact Classification**

The Eastern Chimpanzee’s sensitivity is high because it is Endangered.

Impacts to the Eastern Chimpanzee’s habitat quantity and quality will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. The amount of influx that could be specifically attributed to the construction of the Feeder pipeline is expected to be low, therefore the magnitude of the effects of construction on the habitat quantity and quality of the of the Eastern Chimpanzee is expected to be low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Feeder pipeline, to the application of Project-specific mitigation measures (Table 41). Following the application of the recommended mitigation measures (Section 11.0), the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same.

Impacts to the Eastern Chimpanzee’s abundance and distribution could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. Given its distance from the forest, the operation of the Feeder pipeline is not expected to affect the abundance and distribution of Eastern Chimpanzee in Bugoma CFR, however the Project-associated population influx has the potential to generate impacts. The amount of influx that could be specifically attributed to the construction of the Feeder pipeline is expected to be low, therefore the magnitude of the effects of construction on the abundance and distribution of the Eastern Chimpanzee in Bugoma CFR is low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Feeder pipeline, to the application of Project-specific mitigation measures (Table 41). Following the application of the recommended mitigation measures (Section 11.0), the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same.

Impacts to the survival and reproduction of the population of Eastern Chimpanzee in Bugoma Central Forest Reserve could be adverse. The geographical extent of impacts will be national because effects could extend well beyond the CHAA. Impact duration will be long-term (that is, extending far beyond the pipeline construction period of 10-12 months), as once settlements created by migrants to the area become established, they may be difficult to reverse. There is a possibility that individuals could be killed or severely disturbed as a result of Project-related population influx during the operation phase. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve could be affected through direct mortality from bush meat hunting or other disturbance. However, the amount and intensity of influx that could be specifically attributed to the construction of the Feeder pipeline is expected to be low, therefore the magnitude of the effects of construction on the survival and reproduction of the Eastern Chimpanzee is expected to be low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level during the operation phase of the Feeder pipeline, prior to the application of Project-specific mitigation measures (Table 41). Following the application of the recommended mitigation measures (Section 11.0), the impact significance is expected to decrease to minor, primarily because the magnitude could become negligible, although the sensitivity of the receptor will remain the same.

**Table 35: Potential impacts in the construction phase to the Eastern Chimpanzee**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low - 2	High – 4	Moderate - 8	Negligible - 1	High – 4	Minor - 4





Abundance and distribution	Low - 2	High – 4	Moderate - 8	Negligible - 1	High – 4	Minor - 4
Survival and reproduction	Low - 2	High – 4	Moderate - 8	Negligible - 1	High – 4	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced impacts that the construction of the Feeder pipeline could have on the Eastern Chimpanzees and their habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**11.2 Operation Phase Impacts**

For the assessment of impacts during the operations phase, the key questions were divided into sub-questions that focused on individual valued components within the CHAA and LSA. In answering each question, the individual components of the Project were considered with regards to their potential to affect a valued component. These questions are presented below.

**11.2.1 What impact could the operation of the Feeder Pipeline have on habitats and ecosystem integrity?**

This section presents the assessment of impacts that the operation of the Project could have on the habitat and ecosystem integrity within the CHAA and the LSA. These habitats either do, or could, support populations of species of concern. Therefore, the assessment of potential impacts to those species, and others, occurring in the CHAA has been assessed in this section through the determination of the impacts to potential habitat for those species.

The impacts of the Project on critical habitat, as triggered by species of concern, are covered under the individual assessment of those species in Section 10.2.2. Other triggers of critical habitat are discussed as relevant in the appropriate sections.

As discussed, construction-phase rehabilitation measures should return the site to a grassed, herbaceous vegetation cover similar to open grassland-type habitats that occur in the escarpment vegetation corridor at baseline. Ongoing recovery of vegetation cover from the initial construction-phase habitat losses, and associated edge effects, risk of alien invasive vegetation spread are considered in the construction phase impact assessment (Section 11.1.1). Operation phase impacts of the Feeder pipeline on habitats and ecosystem integrity are therefore limited to those caused by new components and activities, including:

- Maintenance activities particularly clearance of woody vegetation with roots that could damage the pipeline beneath;
- The presence of the heated pipeline approximately 1.0 m beneath the surface; and
- The effect of ongoing in-migration into the CHAA in response to the expectation of job opportunities.

**11.2.1.1 What impact could the operation of the Feeder Pipeline have on the escarpment vegetation corridors?**

**Impact Indicators**

Indicators used to assess impacts of the Feeder Pipeline on the habitat and ecosystem integrity of the vegetation corridors on the escarpment were changes in: regional representativeness; ecosystem composition, and ecosystem configuration.





Literature was reviewed to understand the long-term direct impacts of pipelines through natural habitats. Indirect effects to habitat were estimated by applying a 0.50 km buffer to the pipeline route. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, changes in water quantity and quality, and air emissions and dust.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur over the life of the Project, and the disruption of known corridors. This was accomplished by examining available literature about the ecology of the vegetation communities on the escarpment, and scientific literature about the impacts of human activities on corridors in the long-term.

### **Impact Assessment**

#### ■ Representativeness

The CHAA supports approximately 2443 ha of escarpment corridor vegetation communities (see Section 6.3.1.1.2), which are bounded on the east by agriculturally modified landscapes, and the Buhuka Flats on the west. As mentioned, these vegetation communities form part of a contiguous vegetation corridor that is part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor (Plumptre et al. 2003) (see Section 6.3.1.2.1). Therefore, their continuous extent represents an important landscape feature in the CHAA.

Indirect effects on the representativeness of the escarpment vegetation communities during operation are expected as a result of the ongoing effects of population influx to the region. The presence of a relatively-easy to navigate, grassed linear feature through the escarpment vegetation corridor is likely to facilitate access to other areas of the escarpment vegetation communities for timber harvest and livestock grazing on accessible slopes, resulting in habitat loss and degradation in the vicinity, affecting the representativeness of this section of the escarpment vegetation corridor in the landscape.

Impacts from the changes to representativeness are predicted to be less than those experienced during the construction phase because temporary construction areas will be rehabilitated. Nevertheless, the potential for renewed reduction in habitat extent and condition during operation as a result of population influx in the area is predicted to be beyond the expected range of natural disturbance perturbations.

#### ■ Ecosystem composition

The construction phase rehabilitation measures should return the corridor to a vegetation cover similar to baseline vegetation communities that are suitable for establishment over the underground pipeline (i.e. communities that lack large trees, such as open grassland). Ongoing maintenance will be conducted throughout the operational lifetime of the pipeline to ensure that shrubs and trees, whose roots could potentially damage or penetrate the buried pipeline, do not establish on the surface directly above the pipeline, and is to have a 10m wide permanent right of way along the buried pipeline route, meaning that the climax vegetation community achievable for this area will be open grassland.

The pipeline will be buried to 1.0 m depth and it will be heated to 82°C. The heated pipeline could warm the soil overhead, which has the potential to influence the dominant type of flora species in the grass/herbaceous layer on the surface. For example, drought-tolerant or xerophytic species may dominate, over species that require seasonally-cold periods for dormancy, or species that require more moisture, affecting the ecosystem composition of vegetation communities and associated faunal guilds (e.g. invertebrates).

Ongoing migration into the CHAA are expected to increase existing pressures on escarpment vegetation communities, through harvest of woody resources for timber and fuel, overgrazing, and hunting fauna for bushmeat. Overgrazing pressures could result in shifts in vegetation community composition to more hardy species, whilst hunting pressure would further reduce the remaining mammalian fauna on the escarpment within the CHAA. Birds and other faunal community composition may also change due to hunting and a deterioration in cover.

Impacts from the long-term changes to ecosystem composition are predicted to be less than those experienced during the construction phase because temporary construction areas will be rehabilitated, but still beyond the expected range of natural disturbance perturbations.





### ■ Ecosystem configuration

The wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor is recognised as being important for threatened species in the face of climate change adaptation (Ayebare et al. 2013), and as part of a much broader set of corridors running the length of the Albertine Rift. This corridor in the southern portion of the eastern shores of Lake Albert, in the vicinity of the Project, is very narrow compared to its extent elsewhere, and is recognised as being important for savannah species (Plumptre et al. 2010). The corridor is recognised as an important climate change refugium for a range of threatened species, which may become increasingly important for those species in the future (Ayebare et al. 2013), that is, within the life time of the Project.

The Feeder pipeline will be buried and rehabilitated following construction, and rehabilitation measures should return the site to a vegetation cover similar to baseline vegetation communities that are suitable for establishment over the underground pipeline. During operation, the pipeline is therefore not expected to be a significant barrier for species that could use the escarpment vegetation as a wildlife corridor, such as large ungulates, predators, primates (all of which are very rare in the area in any case), and birds.

Therefore, no significant changes to ecosystem configuration of the escarpment vegetation communities as a result of the long-term operation of the pipeline are predicted.

### **Impact Classification**

The vegetation community of the escarpment's sensitivity is medium because, although these habitats are already under stress from livestock grazing and harvesting of fuel wood and non-timber forest products, they form part of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor.

Impacts to the representativeness of the habitat will be negative because the indirect effects of population influx to the region are likely to result in further loss and degradation of escarpment vegetation communities, in particular in the vicinity of the short-stretch of easily traversed, grassed pipeline route. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be medium-long term because the pipeline will remain in place throughout the project lifetime. The magnitude of the impacts of operation on representativeness of the vegetation communities of the escarpment is low, given the extent of the area in question. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level to representativeness during the operational phase of the Project, pre-mitigation (Table 36). Following the application of appropriate mitigation measures during operation, the impact significance could remain moderate, as the effects of population influx on the escarpment vegetation community extent and condition in the vicinity of the pipeline. may be difficult to mitigate.

Impacts to the ecosystem composition will be adverse in the long-term, because any re-establishing woody vegetation species will be cleared from the pipeline, and the vegetation community may shift to more heat/drought-tolerant species. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be medium-term (that is, ~25 years) because the pipeline will remain in place throughout the project lifetime. The magnitude of the impact on ecosystem composition is low, as although the woody species will be removed, and the vegetation community may see a shift to more heat tolerant species, the re-established grass and herbaceous vegetation layer will remain. Therefore, low magnitude and medium sensitivity combine to produce a moderate overall impact level to ecosystem composition during the operational phase of the Project, pre-mitigation (Table 36). Following the application of appropriate mitigation measures, the impact significance is expected to become minor, as the magnitude will become negligible, while the sensitivity of the receptor will remain the same.

Impacts to the ecosystem configuration will be neutral, as the buried and rehabilitated pipeline will not present a significant barrier to movement for faunal species using the escarpment vegetation corridor. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be medium-term (that is, ~25 years) because the pipeline will remain in place throughout the project lifetime. The magnitude of the impact on ecosystem configuration is therefore negligible, and combines with medium sensitivity to produce a minor overall impact level to ecosystem composition during the operational phase of the Project (Table 36), and no further mitigation measures are required.





**Table 36: Potential impacts in the operation phase of the Feeder Pipeline to the vegetation communities of the escarpment**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low – 2	Medium – 3	Moderate - 6	Low – 2	Medium – 3	Moderate - 6
Ecosystem composition	Low – 2	Medium – 3	Moderate - 6	Negligible – 1	Medium – 3	Minor - 4
Ecosystem configuration	Negligible – 1	Medium – 3	Minor - 4	Negligible – 1	Medium – 3	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Feeder Pipeline-related impacts to the vegetation communities of the escarpment, and the wildlife corridors of which they form part. However, there is some uncertainty with regards to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain. Indications are that they will increase in importance (Ayebare et al. 2013), provided human pressures do not overwhelm them.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**11.2.1.2 What impact could the operation of the Feeder Pipeline have on wetlands and riparian habitat in the CHAA?**

**Impact Indicators**

Indicators used to assess impacts of the Feeder Pipeline on the habitat and ecosystem integrity of the wetland and riparian habitats in the CHAA were changes in: regional representativeness; ecosystem composition; and ecosystem configuration.

Literature was reviewed to understand the long-term direct impacts of pipelines through natural habitats. Indirect affects to habitat were estimated by applying a 200 m buffer to the pipeline route. Specifically, the buffer was selected to account for changes in habitat quantity and/or quality that could be caused by edge effects, fragmentation, sensory disturbance, and changes in water quantity and quality.

Changes in habitat composition and ecosystem configuration were assessed by identifying potential changes in species composition that could occur over the life of the Project, and the disruption of wetland and riparian systems. This was accomplished by examining available literature about wetland ecology, and scientific literature about the impacts of human activities on wetland systems in the long-term.

**Impact Assessment**

■ Representativeness

The CHAA supports approximately 1157.9 ha of wetlands, and 840 ha of riparian vegetation communities associated with drainage lines and riparian systems, of which, approximately 2.6 ha will have been directly affected by construction activity.

During operation, no additional direct effects on the extent and/or condition of wetland and riparian habitats are predicted. However, indirect effects on the representativeness of the wetland and riparian habitats during operation are expected, as a result of the ongoing effects of population influx to the region. Knock-on effects





on the ecological integrity and resilience of the wetlands are predicted, as increased levels of livestock grazing, harvest of reeds and grasses for home construction, and conversion of wetland and riparian habitats for subsistence cropping are likely to occur.

The construction phases losses of habitat are expected to be partially recovered because the pipeline will be buried and rehabilitated following construction. Rehabilitation measures should return the affected areas to a vegetation cover similar to that that was there before the pipeline was constructed.

Impacts from the changes to representativeness are predicted to be less than those experienced during the construction phase because temporary construction areas will be rehabilitated. Nevertheless, the potential for renewed reduction in habitat extent and condition during operation as a result of population influx in the area is predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Ecosystem Composition

The presence of linear infrastructure, such as pipelines, through wetlands is known to have detrimental impacts on the functioning, processes and species composition of these communities (Roise et al. 2004). Buried pipelines passing through wetlands may result in losses of biodiversity at local and downstream scales largely due to changes in geomorphological properties and interruption in hydrological regimes, which could affect the maintenance of flow to wetland areas downstream of the crossing, or result in increased erosion potential around the buried pipeline. Alteration of wetland structure and composition could occur if flow paths for water, both surface and sub-surface, are not maintained. If these flow paths are not maintained throughout the operational lifetime of the project, there is a potential that the wetland community downstream of the obstruction could become altered – typically via flow concentration and channel incision in downstream wetlands that were previously supplied by diffuse, dispersed flows. The risk of incised drainage and associated loss of wetland function due to concentration of water flows may be relatively low, assuming that the construction mitigation measures were successfully applied.

The pipeline will be buried to 1.0 m depth and it will be heated to 82°C. The heated pipeline has the potential to warm the wetland and riparian habitat in its immediate vicinity, which could influence the moisture regime of some habitats (particularly seasonal wetlands) and associated flora and fauna community composition, albeit on a very localised basis.

Although the Feeder pipeline will have been buried and rehabilitated following construction, and rehabilitation measures should return the corridor to a vegetation cover similar to baseline vegetation communities; the likely intensification in use of the wetlands as the human population of the region increases could generate indirect impacts on wetlands. Existing sanitary conditions in the CHAA are poor, with all of the streams being contaminated with faecal waste from both animals and humans (Vol4c Social IA). The wetland communities in the CHAA are already under pressure from livestock grazing, and harvesting of fibre for house construction. It is likely that these communities will become degraded in the long-term as the human population of the region increases. Such changes would alter the habitat structure and vegetation community composition of the wetland and riparian habitats in the CHAA, which, in turn, could affect the utilisation of these habitats by the currently resident faunal species guilds (e.g. the Endangered Grey Crowned Crane).

Impacts from the changes to ecosystem composition of wetland and riparian habitats are predicted to be less than those experienced during the construction phase, because temporary construction areas will be rehabilitated. Nevertheless, the potential for renewed reduction in habitat extent and condition during operation as a result of population influx in the area is predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Ecosystem configuration

Linear infrastructure such as pipelines can present significant barriers to, or can alter behaviours of, a range of wetland wildlife, from: amphibians (for example, Pontoppidan et al. 2013); to turtles (for example, Langen et al. 2012).

Although the Feeder pipeline will cross currently contiguous wetlands and riparian habitats in the LSA, it will be buried and rehabilitated following construction, and rehabilitation measures should eventually return the



affected areas to a vegetation cover similar to baseline vegetation communities. During operation, the pipeline is not expected to represent a significant barrier to movement, or break in habitat connectivity, for wetland and riparian faunal species.

Therefore, no significant changes to ecosystem configuration of the wetland and riparian habitats as a result of the long-term operation of the pipeline are predicted.

### **Impact Classification**

The sensitivity of wetlands and riparian habitats is high, because permanent and seasonal wetlands are important breeding and foraging habitat for Grey Crowned Crane, a species of concern; and wetlands are susceptible to degradation as a result of changes in water flow patterns brought about by infrastructural developments. The habitats are also already under stress from livestock grazing and harvesting of fibre. This high sensitivity is weighed against the magnitude of each of the impacted indicators as described in the paragraphs below, in order to derive the overall impact level for each indicator.

#### ■ Representativeness

Impacts to the representativeness of the habitat will be adverse because the indirect effects of population influx to the region are likely to result in further loss and degradation of wetland and riparian habitat. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be medium-long term because the pipeline will remain in place throughout the project lifetime. The magnitude of the impacts of operation on representativeness of the wetland and riparian habitats is low, given the extent of the potentially-affected areas in question. In the context of high wetland sensitivity, this results in impacts of moderate significance.

Following the application of appropriate mitigation measures, the impact significance could remain moderate, as the effects of population influx on the extent and condition of wetland habitats in the vicinity of the pipeline. may be difficult to mitigate.

#### ■ Ecosystem Composition

Impacts to the ecosystem composition will be adverse. The geographical extent of impacts will be local because impacts are restricted to the LSA. Impact duration will be for the duration of operation (~25 years). The magnitude of the different project-specific impacts on ecosystem composition varies, as discussed under each of the subheadings below.

Impacts to the ecosystem composition arising from the heated pipeline could be adverse in the long term, as the wetland vegetation community in its direct vicinity may be affected. The geographical extent of impacts will be highly localised with impacts restricted to the direct vicinity of the buried pipeline. Impact duration will be medium-term, lasting throughout the operational lifetime of the project. The magnitude of the impact on ecosystem composition is negligible because, although localised changes to the floral and faunal community composition could occur, the extent of the potentially affected area is small, and the likelihood of any significant changes in the wetland or riparian community composition is considered very low. Therefore, the magnitude and high sensitivity of the wetlands combine to produce a minor overall impact level to ecosystem composition during operation of the Feeder pipeline, and no project-specific mitigation measures are required (Table 37).

The ongoing in-migration to the CHAA is likely to result in increased human use of wetlands in the CHAA. In the absence of project interventions, the impact of increased population density (and associated increases in grazing livestock) in the CHAA is expected to be long-term, and to have a material effect on both wetland structure and water quality in the wetlands across the CHAA through reductions in sanitary water quality, exacerbation of wetland erosion, increased harvest of plant species used for traditional home construction, increased fire frequency and increased grazing pressure. The impacts will be irreversible; and of medium magnitude. In the context of the high wetland sensitivity, the predicted impact significance on ecosystem composition is major, prior to mitigation. Following the application of Project-specific mitigation measures during operation, the impact significance could remain moderate, as the effects of population influx on the ecosystem composition of wetlands and riparian habitat in the vicinity of the pipeline may be difficult to mitigate.

#### ■ Ecosystem Configuration





Impacts to the ecosystem configuration will be neutral, as the buried and rehabilitated pipeline will not present a significant barrier to movement for faunal species using the affected wetland and riparian habitats. The magnitude of the impact on ecosystem configuration is therefore negligible, and combines with medium sensitivity to produce a minor overall impact level to ecosystem composition during the operational phase of the Project (Table 37), and no further mitigation measures are required.

**Impact Significance Rating**

**Table 37: Potential impacts of operation of the Feeder pipeline on the wetlands of the CHAA**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Representativeness	Low - 2	High – 4	Moderate – 8	Low - 2	High – 4	Moderate – 8
Ecosystem composition – (heated pipeline)	Negligible – 1	High – 4	Minor - 4	Negligible – 1	High – 4	Minor - 4
Ecosystem composition – (population influx)	Medium - 3	High – 4	Major - 12	Low – 2	High – 4	Moderate – 8
Ecosystem configuration	Negligible – 1	High – 4	Minor - 4	Negligible – 1	High – 4	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential operational impacts that the Feeder pipeline may have on the wetlands of the CHAA. However, there is some uncertainty in regard to the irreplaceability and vulnerability of those habitats in the CHAA. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future is uncertain.

The spatial extent of habitats in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1.1). The actual extent of possible habitat may have been over-estimated.

**11.2.1.3 What impact could the operation of the Feeder Pipeline have on Bugoma CFR?**

**Impact Indicators**

Bugoma Central Forest reserve (Bugoma CFR) will not be directly affected by pipeline operation, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary. The indirect impacts of population influx to the CHAA, due to expectations of work or opportunities generated by the Project, are considered since this could impact on the forest reserve.

An analysis of the potential impacts of such influx on the habitat and ecosystem integrity of Bugoma CFR was accomplished by examining available literature about the ecology of Bugoma CFR, and scientific literature regarding the effects of migration and human population pressure on forests in Africa. The impacts are discussed, and their significance assessed through a **reasoned narrative**.

**Impact Analysis**

Bugoma CFR is identified as a valued component for this impact assessment, certainly in terms of biodiversity (see Table 25). As identified in that section, apart from being one of the last stands of tropical semi-deciduous forests in the region, it also supports known populations of the Endangered Eastern Chimpanzee and range-





restricted Nahan's Francolin (Plumptre et al. 2011), potential non-breeding habitat for the Endangered Madagascar Pond Heron (see Section 6.3.3.1), as well as elephants and a host of other threatened and irreplaceable species.

Although Bugoma CFR will not be directly affected by pipeline operation, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary, the indirect effects of project-induced migration to the CHAA due to expectations of work on the Project or induced opportunities, have the potential to affect the ecological integrity of the habitats within the CFR, as well as the species of conservation concern that occur within it.

Bugoma CFR is under severe pressure from the human population surrounding it. In 2013, more than 1,500 people evicted illegally after settling within the forest (Mugerwa 2013), and recent (2016) reports from the area suggest that land-grab activity has resulted in the conversion of an estimated 8000 ha for sugar cane plantation development. The land cover study (see Figure 4) clearly shows that the areas surrounding the forest have largely been transformed for agricultural and subsistence purposes, and the transformation is now encroaching within the boundary of the CFR. This trend of encroachment of protected areas is not unique. Protected areas are known to be particularly vulnerable to changes in human demographics and deforestation; and Wittemyer et al. (2008) identified that human population growth and encroachment around protected areas is significantly higher than the average population growth in rural areas. This difference was largely due to the immigration of people into these areas because of the perceived increased availability of opportunities, natural resources and potential jobs (Wittemyer et al. 2008). This is supported by research conducted in the forests of the Albertine Graben, and the wider CHAA. For example, in a study of Budongo Central Forest Reserve, Zommers and MacDonald (2012), identified that of the local communities that hunted bush meat in the forest, nearly 73% were immigrants to the area. Furthermore, these workers identified that the households of immigrants were also more likely to be involved with deforestation.

The results of the recently-updated (Nov 2017) social baseline data gathering fieldwork indicated that in-migration to the CHAA has escalated significantly since the opening of the escarpment road, and commencement of preliminary construction works at the Kingfisher Field Development Area; this pattern of in-migration will probably continue throughout the operational lifetime of the Project.

An Influx Management Strategy and Framework Plan was developed to manage the potential influx of people into the LSA. However, this plan can only focus on those measures over which CNOOC has control, and to support the range of government and donor projects in Uganda aimed at socio-economic development and environmental conservation. How this translates to the potential influx of people to the CHAA throughout the operational lifetime of the Project is unknown.

### **Impact Classification**

Indirect impacts on Bugoma CFR due to population influx to the region during the operation of the feeder pipeline will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR, in the CHAA. Impact duration will be long-term, lasting for at least as long as the Project is in operation, and may be irreversible.

However, the magnitude of the impacts of population influx specifically associated with the operation of the pipeline on Bugoma CFR is expected to be negligible, given that the pipeline will be buried, rehabilitated and will be generally remote from the main Production Facility where in-migration is expected to be concentrated, and taking into consideration the distance of the rehabilitated pipeline from the Bugoma CFR boundary.

The sensitivity of the Bugoma Central Forest Reserve is high because it is a threatened ecosystem that is already under pressure. Therefore, the intensity and sensitivity combine to produce a minor overall impact level during the operation phase of the pipeline (Table 31).

The mitigation hierarchy is an important process that has been used to minimise impacts to Bugoma CFR. The focus for the continued application of the mitigation hierarchy to Bugoma CFR during the feeder pipeline operation phase will consist of monitoring and adaptive management, as well as implementation of the Influx Management Strategy and Framework Plan for the Project (see Section 12.0).



With the implementation of such mitigation measures, the impact significance is expected to remain minor for the operational lifetime of the Project. (Table 31).

**Table 38: Potential impacts in the construction phase to the Bugoma Central Forest Reserve**

	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat and ecosystem integrity	Negligible – 1	High – 4	Minor – 4	Negligible – 1	High – 4	Minor – 4

As noted above, the focus for reducing impacts to the Bugoma Central Forest Reserve is mitigation to avoid and minimise various types of disturbance that may occur (Section 12.0). Optimising mitigation will reduce residual impacts on the CFR to minor, and reduce the need for offsetting. However, possible offsetting options to ultimately achieve ‘net gain’ for the natural habitats of the forest have been identified and are discussed in Section 12.0.

**11.2.2 What impact could the operation of the Feeder Pipeline have on species of concern?**

This section presents the assessment of impacts that the operation of the feeder Pipeline could have only on those relevant species of concern identified as biodiversity valued components; that is, Grey Crowned Crane, Nahan’s Francolin and Eastern Chimpanzee (see Table 25). Since the operation of the Feeder pipeline will not affect the near-shore habitats of Lake Albert, potential effects on the Mud Snail *G. candida* are not considered. Potential impacts to other species of concern are assessed at the habitat level (see Section 11.2.1).

**11.2.2.1 What impact could the operation of the Feeder Pipeline have on Grey Crowned Crane?**

Grey Crowned Crane is an Endangered species. In the CHAA, it is associated with permanent and seasonal wetlands habitats.

Given that operational impacts on wetlands in the CHAA are expected (Section 10.2.1.2), the operation of the feeder Pipeline could indirectly impact Grey Crowned Crane breeding and foraging habitat.

The potential direct, indirect and induced impacts to this species due to the operation of the Production Facility are presented below.

**Impact Indicators**

Indicators used to assess effects of the operation of the feeder Pipeline on Grey Crowned Crane were: habitat quantity and quality, habitat connectivity, abundance and distribution, and survival and reproduction.

Potential changes in habitat quantity and quality, abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances arising from increased human presence in the CHAA. These disturbances were considered in light of known or inferred effects to the survival and reproduction of Grey Crowned Crane for which data on these types of effects are available.

Changes in habitat connectivity were assessed by identifying potential barriers to movement and loss of corridors. This was accomplished by visually examining the spatial distribution of critical habitat for Grey Crowned Crane (permanent and seasonal wetlands) in relation to the feeder Pipeline to qualitatively identify any areas where habitat remains/may become fragmented during operation.





### *Impact Assessment*

#### ■ Habitat Quality and Quantity

Construction-phase losses of wetland habitat (permanent wetland) to the 30m-wide Feeder Pipeline construction right of way, consisting of 0.36 ha (0.5%) of 69 ha available habitat in the CHAA, will be mitigated through rehabilitation of the vegetation following construction; the ongoing reduction in habitat quantity and quality for Grey Crowned Crane during the time it takes for vegetation to re-establish is addressed in the construction phase impact assessment (Section 10.1.2.2).

The operational phase of the feeder pipeline, as a component of the Project, could result in an influx of people into the area seeking employment/other opportunities. It can reasonably be expected that an increased human population in the CHAA will put pressure on the wetland habitats preferred by Grey Crowned Crane. These pressures could manifest in increased rates of habitat alteration and deforestation, reducing habitat quality and quantity for Grey Crowned Crane.

Although Grey Crowned Crane can tolerate a low degree of anthropogenic disturbance in foraging habitat (e.g. subsistence and commercial farming practises), in Ugandan wetlands it has been observed to be intolerant of human proximity within 100-200m (Olupot, 2014), flying away on approach; a factor which also affects breeding success as breeding birds flush from nests on approach, causing increased rates of predation, reduced time at the nest (either incubating or feeding), and ultimately nest abandonment. Increased human presence in the CHAA is likely to further reduce habitat quality for Grey Crowned Crane in the CHAA.

Effects from reductions in habitat quality and quantity during operation are predicted to be beyond the expected range of natural disturbance perturbations.

#### ■ Habitat Connectivity

The impact of the feeder pipeline operation as a barrier to the movement of Grey Crowned Crane during operation is likely to be neutral. The pipeline will be buried and the overlying vegetation will have been rehabilitated. Disturbance levels due to human presence in the area are expected to be minimal, compared to construction phase.

Impacts on habitat connectivity are predicted to be within the expected range of natural disturbance perturbations during operation.

#### ■ Abundance and Distribution

The impact of the Feeder pipeline operation on the abundance and distribution of Grey Crowned Crane during operation is likely to be adverse. Although the pipeline will be buried and the overlying vegetation will be rehabilitated, restoring the extent of available foraging habitat, increased human presence in the CHAA throughout the operational lifetime of the Feeder pipeline is likely to result in settlement patterns that focus on wetland and riparian areas, which provide livestock grazing and crop production opportunities, and sources of freshwater. This is likely to affect the abundance and distribution of Grey Crowned Crane within the CHAA.

Impacts on species abundance and distribution are predicted to be beyond the expected range of natural disturbance perturbations during operation.

#### ■ Survival and Reproduction

The impact of the Feeder pipeline operation on the abundance and distribution of Grey Crowned Crane during operation is likely to be adverse. As mentioned, increased human presence in the CHAA throughout the operational lifetime of the Feeder pipeline is likely to result in settlement patterns that focus on wetland and riparian areas. This further reduces the likelihood that Grey Crowned Crane would use the affected wetlands for breeding during operation, and increases the likelihood of chick capture/nest abandonment if they did continue to use the area for breeding.

Impacts on species survival and reproduction are predicted to be beyond the expected range of natural disturbance perturbations during operation.



**Impact Classification**

The Grey Crowned Crane sensitivity is high because it is Endangered, and triggers Tier 2 critical habitat.

Impacts to Grey Crowned Crane habitat quantity and quality will be adverse. The geographical extent of impacts will be regional because population influx is expected to affect wetland habitat across the CHAA which will be reduced in quality as a result of habitat degradation and sensory disturbances, associated with in-migration. Impact duration will be into the far future (~25 years). Nevertheless, the magnitude of the effects of operation of the pipeline on Grey Crowned Crane habitat quantity and quality across the CHAA is considered low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as the pressures on Grey Crowned Crane associated with increased human presence across the CHAA are expected to be difficult to mitigate, affecting habitat quantity and quality (Table 39).

Impacts to Grey Crowned Crane habitat connectivity will be neutral. The geographical extent of impacts will be local because effects will be restricted to the 120 m of rehabilitated wetland habitat that will be crossed by the pipeline. The magnitude of the effects of operation of the pipeline on Grey Crowned Crane habitat connectivity is therefore considered negligible. The magnitude and sensitivity combine to produce a minor overall impact significance, and no further mitigation measures are considered necessary (Table 39).

The impact of the Feeder pipeline operation on the abundance and distribution of Grey Crowned Crane in the CHAA during operation is likely to be adverse. The geographical extent of impacts will be regional because population influx is expected to affect wetland habitat across the CHAA. Impact duration will be into the far future (~25 years). The magnitude of the effects of population influx during the operation of the pipeline on Grey Crowned Crane abundance and distribution is considered low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as the pressures on Grey Crowned Crane habitat suitability associated with increased human presence in the CHAA are expected to be difficult to mitigate, affecting the local distribution of the species (Table 39).

Impacts to the survival and reproduction of Grey Crowned Crane will be adverse. The geographical extent of impacts will be regional because population influx is expected to affect wetland habitat across the CHAA. Impact duration will be into the far future (~25 years). The magnitude of the effects of in-migration during operation of the pipeline on Grey Crowned Crane survival and reproduction is considered low. Therefore, the magnitude and sensitivity combine to produce a moderate overall impact level, pre-mitigation. Following the application of appropriate mitigation measures, the impact significance is expected to remain moderate, as the pressures on Grey Crowned Crane habitat suitability for breeding associated with human presence in the CHAA are expected to be difficult to mitigate, and the sensitivity of the receptor will remain the same (Table 39).

**Table 39: Potential operation phase impacts to Grey Crowned Crane**

Indicator of potential impact	Pre-mitigation			Post-mitigation (pre-offsets)		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Low – 2	High – 4	Moderate - 8	Low – 2	High – 4	Moderate - 8
Habitat connectivity	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Abundance and distribution	Low – 2	High – 4	Moderate - 8	Low – 2	High – 4	Moderate - 8
Survival and reproduction	Low – 2	High – 4	Moderate - 8	Low – 2	High – 4	Moderate - 8







### **Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential Project impacts on the Grey Crowned Crane and its habitat in the CHAA.

The spatial extent of wetland habitats in the CHAA was broadly mapped based on knowledge of the study area, baseline studies, literature and reports (see Section 6.1.1). The actual extent of possible habitat may have been over-estimated.

### **11.2.2.2 What impact could the operation of the Feeder Pipeline have on Nahan's Francolin?**

Nahan's Francolin is a Vulnerable and range-restricted species. In the CHAA, it is restricted to the Bugoma Central Forest Reserve. It triggers Tier 2 critical habitat under Criterion 2 (Section 9.2, Appendix G).

Bugoma Central Forest reserve (Bugoma CFR) will not be directly affected by pipeline operation, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary. The indirect impacts of population influx to the CHAA, due to expectations of work or opportunities generated by the Project, are considered since this could impact on the forest reserve, and subsequently Nahan's Francolin.

### **Impact Indicators**

As the pipeline will be at least 1.8 km from the Bugoma CFR boundary, no effects on habitat connectivity are predicted. Therefore, the indicators used to assess effects of the population influx associated with the operation of the Feeder pipeline on Nahan's Francolin were: habitat quantity and quality, abundance and distribution, and survival and reproduction.

Potential changes in habitat quantity and quality, abundance and distribution, and survival and reproduction were assessed qualitatively by considering changes in disturbances arising from increased human presence in the CHAA. These disturbances were considered in light of known or inferred effects to the survival and reproduction of buttonquail species for which data on these types of effects are available.

### **Impact Assessment**

#### ■ Habitat Quality and Quantity

The Nahan's Francolin's degree of vulnerability to disturbance is unknown. Birds are known to be sensitive to land use and habitat alteration (Lussier et al. 2006). The behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions, such as the availability of alternative foraging sites (Madsen 1998). A reduction in habitat quality and quantity may have conservation implications and consequences for the population of this species.

The operational phase of the Feeder pipeline, as a component of the Project, could result in an influx of people into the CHAA seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma Central Forest Reserve will put pressure on the forest. These pressures could manifest in increased rates of habitat alteration and deforestation. Nahan's Francolin is a shy, forest-dependent species (Sande et al. 2009a) and thus is also probably sensitive to disturbance arising from deforestation/human presence in the forest. These factors could combine to reduce the habitat quality and quantity for Nahan's Francolin.

Effects from loss of habitat are predicted to be beyond the expected range of natural disturbance perturbations.

#### ■ Abundance and Distribution

Nahan's Francolin is reported to be relatively common in the Bugoma Central Forest Reserve (Plumptre et al. 2011). Although the pipeline will be at least 1.8 km from the Bugoma CFR boundary, the indirect impacts of population influx to the CHAA, due to expectations of work or opportunities generated by the Project, could ultimately affect the abundance and distribution of Nahan's Francolin in the CHAA.





The operation of the pipeline, as a component of the overall Project, could result in an influx of people into the area seeking jobs and/or opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma CFR and within the CFR itself will exacerbate existing pressures on the forest. These pressures are likely to manifest in increased rates of bush meat hunting and habitat alteration. These factors could reduce the abundance and distribution of Nahan's Francolin.

Effects from the operation of the pipeline are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

The Nahan's Francolin's degree of vulnerability to direct disturbance, particularly during the breeding season, is not well understood. Generally, for birds, the behavioural response of species to disturbance will depend on species-specific tolerance levels, disturbance type and frequency, group size for socially-foraging animals, and local conditions such as the availability of alternative breeding sites (Madsen 1998). Many studies have reported a reduction in breeding success attributable to human disturbance (for a review, see Hill et al. 1992). Mechanisms include: increased rates of predation, nest abandonment and reduced time at the nest, either incubating or feeding. Clearly, a reduction in breeding output may have conservation implications and consequences for the local population of Nahan's Francolin.

The operation of the pipeline, as a component of the overall Project, could result in an influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding the Bugoma CFR and within the CFR itself will exacerbate existing pressures on the forest. These pressures could manifest in increased rates of bush meat hunting and habitat alteration, which could reduce the survival and reproductive success of Nahan's Francolin.

These effects to the survival and reproductive success of Nahan's Francolin are predicted to be beyond the expected range of natural disturbance perturbations.

### **Impact Classification**

The Nahan's Francolin's sensitivity is high because it is range-restricted, and triggers Tier 2 critical habitat.

Impacts to the Nahan's Francolin's habitat quantity and quality will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term. The magnitude of the impact of population influx associated with the operation of the pipeline, on the habitat quantity and quality of the Nahan's Francolin is expected to be negligible (see Section 10.2.1.3). Therefore, the intensity and sensitivity combine to produce a minor overall impact level during the operational phase of the Project, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 23).

Impacts to the Nahan's Francolin's abundance and distribution could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term. The magnitude of the impact of population influx associated with the operation of the pipeline on the abundance and distribution of the Nahan's Francolin is expected to be negligible (see Section 10.2.1.3). Therefore, the intensity and sensitivity combine to produce a minor overall impact level during the operational phase of the Feeder pipeline, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 23).

Impacts to the survival and reproduction of Nahan's Francolin could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed as a result of population influx to the region during the operation phase. A threshold of 10% for this species' survival and reproduction in the CHAA is reasonable, and it is not expected that this number of individuals in the local population within Bugoma CFR will be affected. The magnitude of the effects of operation of the Feeder pipeline on the survival and reproduction of the Nahan's Francolin is therefore negligible. Therefore, the intensity and sensitivity combine to produce a minor overall impact level during the operational phase of the Feeder pipeline, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 23).



**Table 40: Potential impacts in the operational phase to the Nahan’s Francolin**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude	Sensitivity of the Receptor	Significance	Magnitude	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Abundance and distribution	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Survival and reproduction	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4

**Prediction Confidence**

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced impacts that the operation of the Feeder pipeline could have on the Nahan’s Francolin and its habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.3.1.1). The actual extent of possible habitat may have been over-estimated.

**11.2.2.3 What impact could the operation of the Feeder Pipeline have on Eastern Chimpanzee?**

Eastern Chimpanzee is an Endangered species. A population occurs in the Bugoma Central Forest Reserve. This species triggers Tier 1 critical habitat.

Bugoma Central Forest reserve (Bugoma CFR) will not be directly affected by pipeline operation, which at its nearest point will be at least 1,8 km from the Bugoma CFR boundary. The indirect impacts of population influx to the CHAA, due to expectations of work or opportunities generated by the Project, are considered since this could impact on the forest reserve, and subsequently Eastern Chimpanzee.

**Impact Indicators**

As the pipeline will be at least 1.8 km from the Bugoma CFR boundary, no effects on habitat connectivity during operation are predicted. Therefore, the indicators used to assess effects of the population influx associated with the operation of the Feeder pipeline on Eastern Chimpanzee were: habitat quantity and quality, abundance and distribution, and survival and reproduction.

Potential changes in habitat quantity and quality, and abundance and distribution, were assessed qualitatively by considering changes in disturbances arising from increased human presence in the CHAA. These disturbances were considered in light of known or inferred effects to the survival and reproduction of other populations of chimpanzees for which data on these types of effects are available.

To assess effects to survival and reproduction as a result of in-migration and potential associated increases in poaching and disease spread, in-migration rates were assessed based on the data presented in the social impact assessment (Vol 4c). A literature review of the impact of contact with humans was also conducted.

**Impact Assessment**

- Habitat Quality and Quantity

The potential for the Feeder pipeline (as a component of the overall Project) to be attractive for people seeking opportunities could result in a substantial increase the current population around the Bugoma Central Forest Reserve. An increasing human population could lead to an increase in the demand for agricultural and timber





resources, resulting in increased rates of habitat alteration and deforestation, which, in turn could further fragment, change, and degrade Bugoma CFR and surrounding habitat.

The degree of vulnerability to disturbance experienced by chimpanzees is reasonably well known (for example, see Parren and Byler 2003, Rabanal et al. 2010, Thompson and Wrangham 2013). The chimpanzees living in and around the Bugoma Central Forest Reserve most likely experience sensory disturbances at present from human activities, given the high human populations living around the reserve. Indeed, groups are known to forage in the agricultural fields surrounding the forest, and hence, would more than likely be used to human noises and disturbances (McLennan 2008). As such, the potential sensory disturbances arising from population influx associated with the operation of the pipeline are likely to be minimal. Nevertheless, the magnitude of noise may not be the most important determinant of chimpanzee response. Instead, chimpanzees may respond to 'new' noises or may associate particular noises with other occurrences (for example, machine noise may be associated with human presence, which chimpanzees may, in turn, associate with the presence of danger). Where humans pose a threat, chimpanzees generally avoid them (Hockings and Humle 2009, Parren and Byler 2003). Therefore, the degree of avoidance may depend on the behaviour of people, resulting in varying levels of indirect effects on the quality of available habitat for Eastern Chimpanzee.

Effects from the reduced habitat quantity and quality during operation are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Abundance and Distribution

The Bugoma Central Forest Reserve supports one of the top four Eastern Chimpanzee populations in Uganda, with a population of between 450 and 850 individuals (Plumptre et al. 2010). Given its distance from the forest, the operation of the Feeder pipeline is not expected to detrimentally affect the abundance or distribution of chimpanzees in the forest directly.

The long-term presence of the Feeder pipeline as a component of the overall Project could continue to influence influx of people into the area seeking opportunities. It can reasonably be expected that an increased human population in the area surrounding Bugoma CFR will put pressure on the forest. These pressures could manifest in increased rates of bush meat hunting, and reduced habitat availability, which could affect the abundance and distribution of Eastern Chimpanzee within Bugoma CFR.

Effects are predicted to be beyond the expected range of natural disturbance perturbations.

### ■ Survival and Reproduction

In-migration of people seeking opportunities associated with the Project and Feeder pipeline may adversely affect survival and reproduction of chimpanzees through poaching and disease transfer. Hunting and poaching is a recognised threat to chimpanzees in Bugoma Central Forest Reserve (Plumptre et al. 2003), and this pressure could increase as in-migration of people from other areas occurs. Mitigation and management of in-migration will be very important to minimise potential effects on chimpanzees and other fauna species.

Disease is one of the major threats to Eastern Chimpanzees (Oates et al. 2008); increased abundance of people and competition for land and food resources between humans and chimpanzees could lead to higher rates of disease spread from humans to chimpanzees. Chimpanzees are closely related to humans; therefore, many diseases are transferrable between chimpanzees and humans (Formenty et al. 2003; Isabirye-Basulta and Lwanga 2008). Either direct or indirect contact with humans can spread disease. For example, there is evidence to suggest that respiratory illnesses have been transferred directly to chimpanzees from humans as a result of researcher and tourist contacts, often leading to outbreaks and death (Formenty et al. 2003). Human defecation in forest undergrowth can indirectly lead to spread of intestinal diseases, such as *Clostridium perfringens*, which can be fatal to chimpanzees (Fujita 2011). As forest fragments decrease in size, risks of contact with, and transmission of, disease from humans increases (Isabirye-Basulta and Lwanga 2008). Factors that lead to increased crop raiding or sharing of water resources can also increase the risk of disease spread (Hockings and Hulme 2009). As identified in the social impact assessment (Vol 4c), communicable respiratory diseases are a significant concern in the LSA.



What effect the loss of individuals from the population could have is unknown. It is doubtful that the population could be reduced by 10% and hence, reach that critical population threshold due to population influx associated with the pipeline alone, nevertheless the potential impacts are predicted to be beyond the expected range of natural disturbance perturbations.

**Impact Classification**

The Eastern Chimpanzee’s sensitivity is high because it is Endangered.

Impacts to the Eastern Chimpanzee’s habitat quantity and quality will be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term. The amount of influx that could be specifically attributed to the operation of the Feeder pipeline is expected to be negligible, therefore magnitude of the impact of population influx associated with the operation of the pipeline, on the habitat quantity and quality of the of the Eastern Chimpanzee is expected to be negligible.

Therefore, the magnitude and sensitivity combine to produce a minor overall impact level during the operation phase of the Feeder pipeline, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 41).

Impacts to the Eastern Chimpanzee’s abundance and distribution could be adverse. The geographical extent of impacts will be regional because effects are restricted to Bugoma CFR. Impact duration will be long-term. Given its distance from the forest, the operation of the Feeder pipeline is not expected to affect the abundance and distribution of Eastern Chimpanzee in Bugoma CFR, however the Project-associated population influx has the potential to generate impacts. The amount of influx that could be specifically attributed to the operation of the Feeder pipeline is expected to be negligible, therefore the magnitude of the effects of operation on the abundance and distribution of the Eastern Chimpanzee is negligible. Therefore, the magnitude and sensitivity combine to produce a minor overall impact level during the operation phase of the Feeder pipeline, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 41).

Impacts to the survival and reproduction of the population of Eastern Chimpanzee in Bugoma Central Forest Reserve could be adverse. The geographical extent of impacts will be national because effects could extend well beyond the CHAA. Impact duration will be long-term. There is a possibility that individuals could be killed or severely disturbed as a result of Project-related population influx during the operation phase. A threshold of 10% for this species’ survival and reproduction in the CHAA is reasonable, and it is expected that this number of individuals in the local population within Bugoma Central Forest Reserve could be affected through direct mortality from bush meat hunting or other disturbance. However, the amount of influx that could be specifically attributed to the operation of the Feeder pipeline is expected to be negligible, therefore, the magnitude of the effects of operation on the survival and reproduction of the Eastern Chimpanzee is expected to be negligible. Therefore, the magnitude and sensitivity combine to produce a minor overall impact level during the operation phase of the Feeder pipeline, and no further project-specific mitigation measures for the operation phase are considered necessary (Table 41).

**Table 41: Potential impacts in the operation phase to the Eastern Chimpanzee**

Indicator of potential impact	Pre-mitigation			Post-mitigation		
	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance	Magnitude (the expected size of the impact)	Sensitivity of the Receptor	Significance
Habitat quality and quantity	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Abundance and distribution	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4
Survival and reproduction	Negligible - 1	High – 4	Minor - 4	Negligible - 1	High – 4	Minor - 4





### Prediction Confidence

Given the information available, there is a reasonable understanding of the potential direct, indirect and induced impacts that the operation of the Feeder pipeline could have on the Eastern Chimpanzees and their habitat in the CHAA.

The spatial extent of the dense forest habitat in the CHAA was broadly mapped based on knowledge of the site, baseline studies, literature and reports (see Section 6.1.1). The actual extent of possible habitat may have been over-estimated.

## 12.0 RECOMMENDATIONS FOR MITIGATION AND MONITORING

In applying mitigation to the impact assessment process, it was assumed that the proposed Project design followed the mitigation hierarchy; that is: avoidance; minimisation; rehabilitation; offsets and additional conservation actions (after BBOP 2012, and CNOOC's in-house biodiversity management and aquatic and terrestrial habitat management specifications), and that the Project components and activities assessed here have taken into consideration the application of the mitigation hierarchy. The general principals of the mitigation hierarchy incorporated into the Project design are outlined in Section 12.1.

Nevertheless, additional, impact-specific mitigation measures have been developed for each impacted valued component, in order to minimise residual impacts as much as possible, and ensure that these specific measures are carried through to the Project ESMP for ongoing monitoring and management. These recommended mitigation measures for specific construction and operation-related Project impacts and the schedule for their completion is provided in Section 12.2.

### 12.1 Mitigation Hierarchy

The following sections describe, at a high level, the mitigation measures that CNOOC will use. These mitigation measures follow the mitigation hierarchy (sensu, BBOP 2012), as embodied in CNOOC's *Biodiversity Management Specification* (CNOOC 2014a) and *Aquatic and Terrestrial Habitat Management Specification* (CNOOC 2014b).

#### 12.1.1 Avoidance

Avoidance measures include:

- Locate Project infrastructure in already disturbed and affected habitats, where feasible.
- Locate primary roads (and other linear facilities) away from sensitive areas, where feasible.
- Reuse existing or remnant road networks.
- Limit areas of surface disturbance.

#### 12.1.2 Minimisation

Minimisation measures are measures taken to reduce the duration, magnitude and/or extent of impacts to valued components that cannot be completely avoided. Minimisation measures include:

- Footprint and Infrastructure
  - Limit width of roads and size of Project component areas.
  - Plan an access and road network to minimise total length.
- Physical Hazards
  - Restrict vehicle speeds on roads.
  - Use of buses to transport workers.
  - Restrict construction traffic to daylight hours to reduce the risk of vehicle collisions.



- Install under road crossing structures (for example, culverts) suitable for amphibians and small reptiles, along the access road to reduce road mortalities and improve habitat connectivity.
- Do not leave pipeline trenches open overnight.
- Employ appropriate identification techniques for guy wires and other tall structure like drill rigs.
- People Management
  - Preferentially hire from the local communities to minimise regional human population growth and the associated increase in human encroachment into valued component habitat and direct mortality from illegal hunting.
  - No hunting or collection of fish, flora and fauna permitted for staff and/or contractors.
  - Identify no-go areas for Project personnel; for example, the lake shore and the Bugoma Lagoon.
  - Diseases and pests are controlled to a large extent by proper cleaning, disinfecting, and/or sterilising. To this end, for example, CNOOC will implement widely accepted protocols aimed at minimising the risk of transmitting amphibian chytrid fungus disease in/around the Project site on footwear, vehicles, field equipment, or while handling amphibians in the wild.
  - Prevent or minimise the potential for community exposure to water-borne, water-based, water-related, vector-borne disease, and other communicable diseases that could result from Project activities. Explore opportunities during the project life cycle to improve environmental conditions that could help reduce their incidence.
  - Mandatory environmental training for all workers and contractors that highlights conservation issues and species-specific sensitivities.
- Noise
  - Noise mitigation developed for human receptors will benefit valued components receptors.
  - Limit hours of operation to avoid nocturnal valued component activity periods (that is, nocturnal species).
  - Minimise noise-related disturbances (for example, vehicular traffic) within 200 m of identified sensitive receptors. Some mitigations with special relevance to sensitive valued components include:
    - selection of equipment with lower sound power levels
    - installing enclosures or acoustic barriers without gaps, and with continuous minimum surface density of 10 kg.m<sup>-2</sup> to minimise transmission of sound through the barrier
    - installing vibration isolation devices for mechanical equipment or ensuring that foundations for equipment are designed to minimise transmission of vibration
    - re-locating noise sources to less sensitive areas to take advantage of distance and shielding
    - installation of barriers at facility boundaries such as vegetation curtains or overburden storage areas/soil berms.
  - Minimise higher frequency noises.
  - Stage any progressive noise disturbances (for example, construction activities) to begin at the centre of disturbance (that is, start construction farthest away from known sensitive receptors like population centres) and move progressively towards the edge of habitats, allowing populations to adjust to increasingly closer noise.
- Light



- Do not use bare light bulbs or any lighting pointing upward or outwards, particularly at the periphery of the project footprint or out into the lake.
  - Use the minimum number and brightness of lights required for safety.
  - Use narrow spectrum bulbs to minimise the range of species affected by lighting (for example, longer wave length red or yellow bulbs rather than “natural” or white light).
  - Light only high-risk stretches of roads such as crossings and merges.
  - Implement a Light Management Plan, where lighting that is not needed at a given time is turned off.
  - Retention of at least 10 m of unlit habitat (for example, on either side of key bat commuting/foraging areas) to prevent loss of habitat connectivity.
- Other
- Develop and adhere to airborne pollutant critical load benchmarks for terrestrial and/or aquatic system impacts for the Project.
  - Use dust control methods, such as covers, water suppression, or increased moisture content for open materials storage piles.
  - Control garbage through incineration and covering of landfill to avoid attracting fauna.
  - Conduct site clearing in a pattern to avoid creating habitat 'islands' within the footprint.
  - Conduct site clearing so that progression of clearing moves animals towards intact habitat.

### 12.1.3 Reclamation

- Reclamation of all disturbed areas with indigenous species after decommissioning to approximate pre-existing vegetation.
- Reclamation of those areas in the Project's construction footprint that are no longer needed for the operational phase will minimise the amount of disturbed habitat. Areas no longer required for the Project should be reclaimed by the contractor before close of construction phase. For example, reclamation of existing disturbances within the CHAA that will not be needed for the project.
- Access roads used for construction that will not be needed for Project operation will be closed and reclaimed to natural vegetation when they are no longer needed.

### 12.1.4 Offsetting

In line with the IFC's PS6, offsetting has been considered as an option to achieve no net loss and, preferably, net gain, when residual impacts are identified for valued components that trigger critical habitat and/or natural habitat designations, and where reclamation following the Project's decommissioning are expected not to meet the no net loss philosophy for a valued component. As required, possible offsetting strategies for each affected valued component are discussed in Section 13.0. Actual offsetting strategies should be developed on a landscape-scale, in liaison with other partners and as part of a Biodiversity Action Plan.

## 12.2 Site-Specific Mitigation Measures and Schedule for Completion

### 12.2.1 Pre-construction Phase

Prior to the commencement of construction activities, a number of additional studies should be conducted to gain a more thorough understanding of certain biodiversity valued components. These studies will also better inform those impacts and impacts that have a high degree of uncertainty. These include the following:

- h) Confirm that the Critically Endangered Mud Snail (*Gabbiella candida*), and the range restricted and Near Threatened Snail (*Bellamya rubicunda*) do not occur in CHAA, prior to commencement of construction of the new water intake and pumping station.







- i) The nesting times of Grey Crowned Crane on the Buhuka Flats.
- j) The relative importance of the Buhuka Flats for migratory bird species.
- k) The precise boundaries and locations of the wetlands along the pipeline route.
- l) The location of caves and cavities along the escarpment that could be important for cavity-roosting bats.

### 12.2.2 Construction and Decommissioning Phase

The mitigation measures are presented below and summarised on Table 42. Possible offset mechanisms are summarised in Table 50.

#### 12.2.2.1 Construction and Decommissioning Phase Impacts on Ecosystems of Concern

##### Near-shore habitats of Lake Albert

The following impact mitigation is recommended to minimise the risks of spillages and discharges affecting the near-shore habitats of Lake Albert:

- Make provision for the designs of well pads to be checked by pollution control experts to ensure that the risks of spillage/overflow associated with drilling pollution management systems is minimised.
- Establish a pollution management system, to be fully defined in the Contractor's contractual commitments, covering personnel, training, lines of responsibility, immediate action requirements, on-site spill kits, and all other factors necessary to ensure there is a provision for effective preventative and corrective action during all stages of construction and drilling.
- All machinery and vehicles must be certified as being free of weed propagules, prior to travelling to the construction site. Issue clearance certificates for each piece of machinery and equipment.
- Develop a culture of zero tolerance for pollution during the construction phase of the project.
- Provide a high level of competent environmental oversight during drilling of wells and construction of the CPF.
- Provide for thorough induction training of all construction personnel regarding pollution management, and ongoing refresher training throughout the construction/drilling contracts.
- Provide specific training to staff responsible for the oversight of pollution control systems
- Ensure structured, daily, monitoring of pollution control systems on the well pads and at the CPF to minimise the risk of inadvertent spills and to respond quickly and effectively to any spills that occur. Emphasis must be on preventative measures.
- If biocides are used in hydrotest water, then the discharge should be retained until it is confirmed to pass biotoxicity testing standards; if it fails to achieve these it should be treated appropriately to achieve required standards.
- Develop specific biological monitoring performance indicators as a part of the Construction and Drilling EMPs.

The following impact mitigation is recommended to minimise the risks of the construction of the new water intake station affecting sediment drift patterns on the near-shore habitats of Lake Albert:

- Construction of the water intake station on wooden or concrete piles, rather than extending a rock foundation-type structure (similar to the existing jetty) from the lake shore to the intake point, to minimise effects on lakeshore currents and long-shore drift of sediments.



- Sediment build-up on the wave-ward side of the water-intake station structure should be removed and redistributed to any eroded areas down-drift of the intake structure, as well as at the existing jetty structure.

### Escarpment Vegetation Corridors

The following mitigation and monitoring is proposed:

- Pre-construction surveys for species of concern associated with the escarpment vegetation corridors, in particular, bats and potential bat roosts, should be completed.
- Construction activities that generate noise and vibrations, particularly blasting activity that may be necessary to excavate trenches for the pipeline, should only occur during designated times.
- The use of high-frequency noise emitters (e.g. vehicle reverse signals) should be minimised.
- Rehabilitation of vegetation communities following completion of construction work and burial of pipeline.
- Develop plans, with Government, to reduce the risk of in-migration and subsequent increases in natural resource harvesting pressure into the area during the construction phase (further detail of the actions required are included in Section Vol 4c & Chapter 7 Social Impacts)
- Develop specific biological monitoring performance indicators as a part of the Construction and Drilling EMPs.

### Wetlands

The following impact mitigation and monitoring is recommended:

- Undertake pre-clearance surveys for wetland species of concern within and near the project footprint, such as nesting and foraging sites of the Grey Crowned Crane. Implement measures to ensure that the risk to these species is minimised.
- Minimise wetland vegetation cleared for the Kamansinig flowline crossing to the smallest footprint.
- Demarcate the construction right of way across the Kamansinig wetland to prevent inadvertent damage outside of this footprint.
- Preferably undertake the construction of the flowline crossing across the Kamansinig wetland in the dry season.
- Prohibit access to personnel outside of the defined project work sites and access roads. Train personnel to understand the sensitivity of the local environment in induction and ongoing tool box talks.
- Specifically prohibit project personnel from access to the Bugoma Lagoon which is resource of exceptionally high ecological and cultural value. The Bugoma Lagoon is a part of the Kamansinig wetland system, all of which is regarded as sensitive
- Ensure that erosion protection measures are in place during construction to minimise runoff from disturbed areas into the rivers and wetlands
- Develop a detailed method statement for the flowline wetland crossing of the Kamansinig River to well Pad 3; defining the requirements to contain construction equipment within the construction footprint, to minimise compaction of wetland soils, to reinstate any clay layers and replace soils in the correct order and to return the wetland to the same profile that existed before construction.
- Ensure that all vehicles and machinery are in sound mechanical order, do not have any oil leaks and are fitted with appropriate mufflers to minimise nuisance affecting wildlife.
- Ensure that any pumps, generators or other equipment containing oil used to manage water at the wetland crossing are located on impervious plastic sheeting or drip trays.





- Prohibit any refuelling of equipment within 100 m of a wetland.
- Prohibit the use of backfill intended to provide firm footing for vehicles in wetlands.
- Manage all hazardous products and wastes to minimise the risk of escaped outside of controlled areas (management according to measures recommended in Chapter 02 Waste Management)
- Adjust the final design of the canals channelling stormwater and treated sewage effluent from the CPF to remain outside of the seasonally wet areas associated with River 1, crossing the river channel just upstream of the road culvert.
- From the culvert onward, it may be necessary to canalise the flow to the lake. Use open cross section swales for this purpose (not concrete canalisation), reinforced if necessary and grassed. Finalise the canal design and the alignment of the stormwater drains with the assistance of a wetland ecologist.

### Bugoma CFR

The following impact mitigation and monitoring is recommended:

- De-list the R5 from the proposed oil road upgrades. CNOOC has confirmed that it does not need this road, either for construction or operational purposes. The Ugandan Government has been formally notified. CNOOC will use the P1 as the major haul road during the construction phase and, if upgraded in time, the R7.
- Limit vehicle speeds to 40 km/h along the P1 road in the section from Mpanga to Nsozi. Monitor vehicle speeds and fine drivers who do not comply with the speed limit.
- Prohibit transport of construction materials in the area of the forest at night.
- Widen the P1, where necessary, on the non-forest side of the road in order to minimise forest habitat loss.
- Ensure that all EPC contract transporters are fully aware of the risks to wildlife in the Bugoma Forest.
- Increase monitoring of population changes in the CHAA and, in particular, any incursions into the BCFR by settlement or people harvesting natural resources. A strategy for this initiative is discussed in further detail in Chapter 17, Cumulative Impacts.
- Implement the Influx Management Strategy and Framework Plan that has been developed along side the ESIA to manage the potential influx of people into the LSA. However, this plan can only focus on those measures over which CNOOC has control, and to support the range of government and donor projects in Uganda aimed at socio-economic development and environmental conservation. How this translates to the potential influx of people along an improved P1 road is unknown.

#### 12.2.2.2 Construction and Decommissioning Phase Impacts on Species of Concern

##### Mud Snail (*Gabbiella candida*)

The following impact mitigation and monitoring is recommended:

- Undertake final targeted, once off, specialist surveys for *G. candida* before construction starts at the sites where construction disturbance will occur in Lake Albert (jetty expansion and water intake). The surveys should ascertain the presence/absence of *G. candida* in the near-shore habitats of Lake Albert within the CHAA.
- Since this species triggers Tier 1 Critical Habitat, if it is found to be present, work in near-shore habitats should be postponed until appropriate solutions for the conservation and management of *G. candida* are devised by suitably experienced molluscan specialists, and approved by NEMA.



- If found to be present, the Client will need to demonstrate that the proposed construction activities will affect less than 10% of the known global population of the species – a comprehensive survey of habitats with potential to support the Mud Snail on the shores of Lake Albert will be required to support this demonstration. Thereafter, if less than 10% of the known population would be affected, a Species-Specific Action Plan as part of the overall Biodiversity Action Plan (BAP) must be developed to achieve net gain for the affected species.
- The construction phase mitigation measures for near-shore aquatic habitats (Section 12.2.2.1) should be implemented and strictly adhered to in order to minimise potential loss, fragmentation or degradation of the Mud Snail's habitat.

### Grey Crowned Crane

The following impact mitigation and monitoring is recommended:

- Implement the mitigation set out for wetlands under Section 12.2.2.1 to reduce further loss, fragmentation and degradation of habitats.
- Implement measures to minimise impacts on Grey Crowned Crane abundance and distribution, and reproduction and survival in the CHAA, and the Buhuka Flats in particular. Measures should include:
  - Restrict access by any CNOOC staff, subcontractors and members of the public from any identified areas of breeding habitat used by Grey Crowned Crane within 200 m of suitable nesting sites.
  - Erect/plant screens between construction activities and wetland habitats in order to reduce the likelihood of disturbance of Grey Crowned Crane via human presence, and minimise noise disturbance.
  - Prohibit CNOOC staff and construction subcontractors from entering areas beyond the construction rights of way.
  - Develop and disseminate community education programmes on Grey Crowned Crane habitat conservation, prevention of illegal trade in wild birds and chicks, and prevention of incidences of poisoning.
- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of Grey Crowned Crane in the CHAA (this recommendation is developed further in chapter 17, Cumulative Impacts).

### Nahan's Francolin

The following impact mitigation and monitoring is recommended:

- Implement the mitigation set out for Bugoma CFR under Section 12.2.2.1.
- Large, mature buttressed trees that constitute suitable nesting habitat for Nahan's Francolin should be avoided during vegetation clearance works for the P1 road upgrade.
- Implement measures to minimise impacts on Nahan's Francolin abundance and distribution, and reproduction and survival in the CHAA, particularly those arising from sensory disturbance caused by human presence and mechanical noise generated during construction activity associated with the P1 road upgrade activity. These should include restrictions in operating hours for heavy machinery, use of low-pitched reverse alerts, and restriction of access for road construction workers to areas beyond the road upgrade right of way.
- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of Nahan's Francolin in Bugoma Forest (this recommendation is developed further in Chapter 17, Cumulative Impacts).
- Support the government in enforcement of existing government forestry policies in Uganda.



### Eastern Chimpanzee

The following impact mitigation and monitoring is recommended:

- Implement the mitigation set out for Bugoma CFR under Section 12.2.2.1.
- Implement measures to minimise impacts on Eastern Chimpanzee abundance and distribution, and reproduction and survival in the CHAA, particularly those arising from sensory disturbance caused by human presence and mechanical noise generated during construction activity associated with the P1 road upgrade activity. These should include restrictions in operating hours for heavy machinery, use of low-pitched reverse alerts, and restriction of access for road construction workers to areas beyond the road upgrade right of way.
- Develop and disseminate community education programmes on Eastern Chimpanzee habitat conservation, and prevention of illegal trade in wild animals for live trade and bushmeat, in liaison with existing Eastern Chimpanzee conservation programmes (e.g. Jane Goodall Institute Uganda's environmental education programme).
- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of the Eastern Chimpanzee in Bugoma Forest (this recommendation is developed further in Section 17, Cumulative Impacts).
- Support the government in enforcement of existing government forestry policies in Uganda.



## BIODIVERSITY IMPACT ASSESSMENT

**Table 42: Mitigation and management measures for biodiversity, during the construction and decommissioning phases of the Project**

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
<p>Reduced abundance/distribution of species of concern</p> <p>Reduced survival/reproduction of species of concern</p>	<p>Grey Crowned Crane</p> <p>Escarpment vegetation corridor</p> <p>Wetlands</p>	<p>Pre-clearance surveys for species of concern and habitats of special significance within and near the Project footprint; e.g. nesting sites of species of concern (Grey Crowned Crane), bat roosting sites</p>	<ul style="list-style-type: none"> <li>■ Number of species located</li> <li>■ Locality and populations of invasive species</li> <li>■ Location of significant habitats, including nesting sites</li> <li>■ Locations of suitable relocation sites for individuals</li> <li>■ Number of individuals relocated</li> <li>■ Realignment of Project footprint to avoid sensitive habitats</li> </ul>	<p>Weekly, before any clearing activities</p>	<p>CNOOC and Contractors</p>	<ul style="list-style-type: none"> <li>■ Identification of species of concern</li> <li>■ Management and control of invasive species</li> <li>■ Impactive management of bat roosts</li> <li>■ Handling of animals</li> <li>■ Impactive translocation of threatened plants, and/or collection of reproductive material</li> </ul>
	<p>Mud Snail (<i>Gabbiella candida</i>)</p>	<p>Targeted surveys for Mud Snail (<i>Gabbiella candida</i>)</p>	<ul style="list-style-type: none"> <li>■ Number of survey plots and samples</li> </ul>	<p>Once off before construction activities</p>	<p>CNOOC</p>	<ul style="list-style-type: none"> <li>■ Specialist required to survey for, and identify, snails</li> </ul>
<p>Spread of invasive species</p>	<p>Near-shore habitats</p> <p>Escarpment vegetation corridor</p> <p>Wetlands</p> <p>Bugoma CFR</p>	<p>All machinery and vehicles entering the site should be certified clear of weed propagules</p>	<ul style="list-style-type: none"> <li>■ Inspection forms and clearance certificates for each vehicle and piece of machinery</li> </ul>	<p>As required before new machinery or vehicles arrive at site</p>	<p>CNOOC and Contractors</p>	<ul style="list-style-type: none"> <li>■ Impactive clean down of machinery and vehicles to remove weed seeds</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
			working on the site			
Sensory disturbance	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee	All vehicles and machinery are in sound mechanical order and fitted with appropriate mufflers	<ul style="list-style-type: none"> <li>Service records for each piece of machinery and vehicle</li> <li>Daily pre-start inspection logs for each vehicle and piece of machinery</li> </ul>	Daily	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Daily pre-start vehicle inspection</li> </ul>
Sensory disturbance	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR Grey Crowned Crane Nahan's Francolin	Noise management	<ul style="list-style-type: none"> <li>Noisy construction activities only occur during designated times</li> <li>High frequency noise emitters minimised</li> </ul>	Daily	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Noise management</li> </ul>
Clearing of vegetation	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR	Restrict access Minimise clearance footprint	<ul style="list-style-type: none"> <li>No go areas identified and marked on the ground</li> <li>Amount of vegetation cleared kept to absolute minimum</li> </ul>	Daily	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Information on erosion and sediment control</li> <li>Impactive fauna management during vegetation clearing</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
			<ul style="list-style-type: none"> <li>■ Clearing of vegetation to occur at the edges of contiguous vegetation patches first to allow disturbed fauna to move away</li> <li>■ Area of bare ground exposed at any one time</li> </ul>			
Levelling and grubbing of ground for placement of infrastructure	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR	Minimise extent of vegetation clearance Implement erosion sediment and dust control measures	<ul style="list-style-type: none"> <li>■ Area of bare ground exposed at any one time</li> <li>■ Erosion and sediment control measures in place and functional</li> <li>■ Dust fall-out beyond the construction footprint</li> <li>■ Appropriate storage and handling of topsoil to be used in rehabilitation works</li> </ul>	Daily, before works commence Immediately before impending rain event	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Impactive erosion and sediment control</li> <li>■ Topsoil handling and management</li> </ul>
Levelling and grubbing		Rehabilitation of disturbed areas	<ul style="list-style-type: none"> <li>■ Area of exposed soil</li> </ul>	Monthly	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Rehabilitation of disturbed ground</li> </ul>





## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
of ground for placement of infrastructure	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR	following completion of construction activities	<ul style="list-style-type: none"> <li>■ Success of rehabilitation measures</li> <li>■ Establishment of a cover crop</li> </ul>			
Accidental spillages	Near-shore habitats Wetlands	Management and clean-up of chemical spills	<ul style="list-style-type: none"> <li>■ Number of spills, including volume spilt</li> <li>■ No adverse impacts from spills</li> </ul>	Monthly	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Impactive spill response and clean-up</li> </ul>
Sensory disturbance Collision risk	Wetlands Bugoma CFR	Traffic management	<ul style="list-style-type: none"> <li>■ Speed limits of traffic within Project footprint</li> <li>■ No mortality of wildlife in the LSA</li> <li>■ No driving at night</li> </ul>	Monthly	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Safe driving</li> </ul>
Mortality/ injury risk	Escarpment vegetation corridor Wetlands	Trenching and pipeline laying – prevention of fauna injuries and mortality	<ul style="list-style-type: none"> <li>■ Trenches are not left open overnight</li> <li>■ If trenches are to be left open overnight, adequate ramps for the egress of trapped animals must be installed</li> <li>■ Adequate refuges must be installed at</li> </ul>	Every morning before work commences	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Safe handling and removal of animals trapped in trenches</li> </ul>





## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
			<p>regular intervals along the open trench (e.g., wet sacks)</p> <ul style="list-style-type: none"> <li>Number of animals relocated from the trench</li> </ul>			
<p>Vegetation clearance Earthworks Pipeline crossings</p>	Wetlands	Construction in wetlands and watercourses	<ul style="list-style-type: none"> <li>Maintenance of natural flow regimes</li> <li>Flow volumes the same downstream of excavation compared to upstream</li> <li>No sediment-laden water leaves construction area, use of coffer dams</li> <li>Assessment of potential acid sulphate soils before excavation commences</li> <li>If acid sulphate soils detected, adequate lime must be available on site to treat spoil from the</li> </ul>	Weekly, or as required before excavation in wetlands	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Management and treatment of acid sulphate soils</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
			excavation activities			
Reductions in Habitat quality	Near-shore habitats Escarpment vegetation corridor Wetlands	Control Night lighting	<ul style="list-style-type: none"> <li>■ All night lighting face in towards the Project footprint</li> <li>■ No lights face out towards the lake</li> <li>■ Lighting should be kept to a functional minimum in all areas</li> <li>■ Lamps should not emit light at angles greater than 70°</li> <li>■ Lights that emit a broad spectrum of light with a high UV component should be avoided</li> <li>■ Polarised light sources should not be used</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Impactive night lighting</li> </ul>
Reduction in habitat quality Reduction in survival/	Grey Crowned Crane	Putrescible and industrial waste	<ul style="list-style-type: none"> <li>■ All food wastes must be stored appropriately to discourage vermin</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Waste management</li> </ul>





## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
reproduction due to attraction of predatory/scavenging species			<ul style="list-style-type: none"> <li>Volume of waste removed, recycled, reused</li> </ul>			
Accidental spillages	Near-shore habitats Wetlands Grey Crowned Crane	Water quality and water management	<ul style="list-style-type: none"> <li>In situ water quality downstream of Project footprint (pH, EC, TDS, TSS, DO)</li> <li>Monthly water quality parameters in the lake, wetlands and watercourses (pH, EC, TDS, TSS, DO, metals, hydrocarbons)</li> <li>Volume of water extracted and treated on site</li> </ul>	Weekly and monthly	CNOOC	<ul style="list-style-type: none"> <li>Water quality monitoring</li> <li>Waste water treatment</li> </ul>
Reduction in habitat quality	Grey Crowned Crane	Air quality	<ul style="list-style-type: none"> <li>No noticeable odours at boundary of Project footprint</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>Odour management and detection</li> </ul>
Population influx	Near-shore habitats Escarpment vegetation corridor Wetlands	People management	<ul style="list-style-type: none"> <li>No personnel and/or contractors allowed beyond</li> </ul>	As required	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Inductions for all staff</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
	Mud Snail Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee		footprint of Project ■ Designated no-go areas, e.g., Lake Albert, wetlands, Bugoma Central Forest Reserve			



### 12.2.3 Operation Phase

The mitigation measures for the operation phase of the Project are presented in tabular format in Table 43. Possible offset mechanisms are summarised in Table 50.

#### 12.2.3.1 Operation Phase Impacts on Ecosystems of Concern

##### Near-shore habitats of Lake Albert

The following impact mitigation is recommended to minimise the risks of spillages affecting the near-shore habitats of Lake Albert:

- Establish a pollution management system, to be fully defined in CNOOC's EMP as well as in any sub-contractor's contractual commitments, covering personnel, training, lines of responsibility, immediate action requirements, on-site spill kits, and all other factors necessary to ensure there is a provision for effective preventative and corrective action during all stages of operation.
- All machinery and vehicles must be certified as being free of weed propagules, prior to travelling to the site. Issue clearance certificates for each piece of machinery and equipment.
- Develop a culture of zero tolerance for pollution during the operation phase of the project.
- Provide a high level of competent environmental oversight during drilling of wells and operation of the CPF.
- Provide for thorough induction training of all operation personnel regarding pollution management, and ongoing refresher training throughout the operational lifetime of the project.
- Provide specific training to staff responsible for the oversight of pollution control systems
- Ensure structured, daily, monitoring of pollution control systems on the well pads and at the CPF to minimise the risk of inadvertent spills and to respond quickly and effectively to any spills that occur. Emphasis must be on preventative measures.
- Develop specific biological monitoring performance indicators as a part of the Drilling EMPs.

##### Escarpment Vegetation Corridors

The following mitigation and monitoring is proposed:

- The use of high-frequency noise emitters (e.g. vehicle reverse signals) should be minimised.
- Ongoing monitoring and additional rehabilitation of vegetation communities over the buried pipeline as necessary, throughout the lifetime of the Project.
- Develop plans, with Government, to reduce the risk of in-migration and subsequent increases in natural resource harvesting pressure into the area during the operation phase (further detail of the actions required are included in Vol 4c Social Impacts)
- Develop specific biological monitoring performance indicators as a part of the Project Operation EMPs.

##### Wetlands

The following impact mitigation and monitoring is recommended:

- Prohibit access to personnel outside of the defined project work sites and access roads. Train personnel to understand the sensitivity of the local environment in induction and ongoing tool box talks.
- Specifically prohibit project personnel from access to the Bugoma Lagoon which is resource of exceptionally high ecological and cultural value. The Bugoma Lagoon is a part of the Kamansinig wetland system, all of which is regarded as sensitive



- Ensure that erosion protection measures are in place during operation to minimise runoff from disturbed areas into the rivers and wetlands.
- Ensure that all vehicles and machinery are in sound mechanical order, do not have any oil leaks and are fitted with appropriate mufflers to minimise nuisance affecting wildlife.
- Ensure that any pumps, generators or other equipment containing oil used to manage water at the wetland crossing are located on impervious plastic sheeting or drip trays.
- Prohibit any refuelling of equipment within 100 m of a wetland.
- Manage all hazardous products and wastes to minimise the risk of escaped outside of controlled areas (management according to measures recommended under Vol 1-3)
- Monitoring should remain ongoing at frequent intervals to ensure that rehabilitated vegetation continues to thrive and progress, and to confirm that the recommended mitigation measures to minimise wetland erosion and flow concentration remain satisfactorily functioning.

### Bugoma CFR

The following impact mitigation and monitoring is recommended:

- De-list the R5 from the proposed oil road upgrades. CNOOC has confirmed that it does not need this road, either for construction or operational purposes. The Ugandan Government has been formally notified. CNOOC will use the P1 as the major haul road during the construction phase and, if upgraded in time, the R7.
- Limit vehicle speeds to 40 km/h along the P1 road in the section from Mpanga to Nsozi. Monitor vehicle speeds and fine drivers who do not comply with the speed limit.
- Prohibit transport of Project-related materials in the area of the forest at night.
- Ensure that all EPC contract transporters are fully aware of the risks to wildlife in the Bugoma Forest.
- Increase monitoring of population changes in the CHAA and, in particular, any incursions into the Bugoma CFR by settlement or people harvesting natural resources. A strategy for this initiative is discussed in further detail in Chapter 17, Cumulative Impacts.
- Implement the Influx Management Strategy and Framework Plan that has been developed to manage the potential influx of people into the LSA. However, this plan can only focus on those measures over which CNOOC has control, and to support the range of government and donor projects in Uganda aimed at socio-economic development and environmental conservation.

#### 12.2.3.2 Operation Phase Impacts on Species of Concern

##### Mud Snail (*Gabbiella candida*)

The following impact mitigation and monitoring is recommended:

- If found to be present within the CHAA, the Client will need to demonstrate that the proposed operation of the Project will affect less than 10% of the known global population of the species –a comprehensive survey of habitats with potential to support the Mud Snail on the shores of Lake Albert will be required to support this demonstration. Thereafter, if less than 10% of the known population would be affected, a Species-Specific Action Plan as part of the overall Biodiversity Action Plan (BAP) must be developed to achieve net gain for the affected species.

##### Grey Crowned Crane

The following impact mitigation and monitoring is recommended:



- Implement the mitigation set out for wetlands under Section 11.2.3.1 to reduce further loss, fragmentation and degradation of habitats.
- Implement measures to minimise impacts on Grey Crowned Crane abundance and distribution, and reproduction and survival in the CHAA, and the Buhuka Flats in particular. Measures should include:
  - Restrict access by any CNOOC staff, subcontractors and members of the public from any identified areas of breeding habitat used by Grey Crowned Crane within 200 m of suitable nesting sites.
  - Erect/plant screens between operation activities and wetland habitats in order to reduce the likelihood of disturbance of Grey Crowned Crane via human presence, and minimise noise disturbance.
  - Prohibit CNOOC staff and construction subcontractors from entering areas beyond the operational footprint.
  - Develop and disseminate community education programmes on Grey Crowned Crane habitat conservation, prevention of illegal trade in wild birds and chicks, and prevention of incidences of poisoning.
- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of Grey Crowned Crane in the CHAA (this recommendation is developed further in Chapter 17, Cumulative Impacts).

### Nahan's Francolin

The following impact mitigation and monitoring is recommended:

- Implement the mitigation set out for Bugoma CFR under Section 12.2.3.1.
- Implement measures to minimise impacts on Nahan's Francolin abundance and distribution, and reproduction and survival in the CHAA, particularly those arising from sensory disturbance caused by traffic generated during operation, and effects arising from increased public access to the forest facilitate by the upgraded road. These should include restrictions in operating hours for heavy machinery, use of low-pitched reverse alerts, and restriction of access for Project employees and contractors.
- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of Nahan's Francolin in Bugoma Forest (this recommendation is developed further in Chapter 17, Cumulative Impacts).
- Encourage ecotourism projects run by community groups, and initiate community conservation awareness programmes.
- Support the government in enforcement of existing government forestry policies in Uganda.

### Eastern Chimpanzee

The following impact mitigation and monitoring is recommended:

- Implement the mitigation set out for Bugoma CFR under Section 11.2.2.1.
- Implement measures to minimise impacts on Eastern Chimpanzee abundance and distribution, and reproduction and survival in the CHAA, particularly those arising from sensory disturbance caused by human presence and mechanical noise generated by traffic and enhanced public access to the forest.
- Develop and disseminate community education programmes on Eastern Chimpanzee habitat conservation, and prevention of illegal trade in wild animals for live trade and bushmeat, in liaison with existing Eastern Chimpanzee conservation programmes (e.g. Jane Goodall Institute Uganda's environmental education programme).





- Develop and implement a long-term research and monitoring programme to improve understanding of the behaviour and status of the Eastern Chimpanzee in Bugoma Forest (this recommendation is developed further in Chapter 17, Cumulative Impacts).
- Support the government in enforcement of existing government forestry policies in Uganda.



## BIODIVERSITY IMPACT ASSESSMENT

**Table 43: Mitigation and management measures for biodiversity, during the operation phase of the Project**

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Reductions in Habitat quality and quantity Sensory disturbances Collision risk	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee	Ongoing monitoring of sensitive habitats and populations of species of concern	<ul style="list-style-type: none"> <li>■ Populations of species of concern</li> <li>■ Locality and populations of invasive species</li> <li>■ Condition and extent of significant habitats, including nesting sites</li> </ul>	Six-monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Identification of species of concern</li> <li>■ Management and control of invasive species</li> <li>■ Impactive management of bat roosts</li> <li>■ Impactive translocation of threatened plants, and/or collection of reproductive material</li> </ul>
Reductions in Habitat quality and quantity	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR	Ongoing monitoring of rehabilitated areas	<ul style="list-style-type: none"> <li>■ Area of exposed soil</li> <li>■ Success of rehabilitation measures</li> <li>■ Establishment of a cover crop</li> </ul>	Six-monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Monitoring and assessment of vegetation</li> </ul>
Introduction/spread of invasive species	Near-shore habitats Wetlands	Inspection of vehicles entering the site for weed propagules	<ul style="list-style-type: none"> <li>■ Inspection forms and clearance certificates for each vehicle and piece of machinery working on the site</li> </ul>	As required before new machinery or vehicles arrive at site	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Impactive clean down of machinery and vehicles to remove weed seeds</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Sensory disturbance	Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee	All vehicles are in sound mechanical order and fitted with appropriate mufflers	<ul style="list-style-type: none"> <li>Service records for each piece of machinery and vehicle</li> <li>Daily pre-start inspection logs for each vehicle and piece of machinery</li> </ul>	Daily	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Daily pre-start vehicle inspection</li> </ul>
Sensory disturbance	Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee	Noise management	<ul style="list-style-type: none"> <li>Noisy activities only occur during designated times</li> <li>High frequency noise emitters minimised</li> </ul>	Daily	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Noise management</li> </ul>
Accidental spillages	Near-shore habitats Wetlands	Management and clean-up of chemical spills	<ul style="list-style-type: none"> <li>Number of spills, including volume spilt</li> <li>No adverse impacts from spills</li> </ul>	Monthly	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Impactive spill response and clean-up</li> </ul>
Collision risk	Grey Crowned Crane Nahan's Francolin Eastern Chimpanzee	Traffic management	<ul style="list-style-type: none"> <li>Speed limits of traffic within Project footprint</li> <li>No mortality of wildlife in the LSA</li> <li>No driving at night</li> </ul>	Monthly	CNOOC and Contractors	<ul style="list-style-type: none"> <li>Safe driving</li> </ul>
Reductions in Habitat quality	Grey Crowned Crane Nahan's Francolin	Night lighting	<ul style="list-style-type: none"> <li>All night lighting face in towards the Project footprint</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>Impactive night lighting</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
	Eastern Chimpanzee		<ul style="list-style-type: none"> <li>■ No lights face out towards the lake</li> <li>■ Lighting should be kept to a functional minimum in all areas</li> <li>■ Lamps should not emit light at angles greater than 70°</li> <li>■ Lights that emit a broad spectrum of light with a high UV component should be avoided</li> <li>■ Polarised light sources should not be used</li> </ul>			
Reduction in habitat quality Reduction in survival/ reproduction due to attraction of predatory/ scavenging species	Grey Crowned Crane	Putrescible and industrial waste	<ul style="list-style-type: none"> <li>■ All food wastes must be stored appropriately to discourage vermin</li> <li>■ Volume of waste removed, recycled, reused</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Waste management</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Accidental spillages	Near-shore habitats Wetlands	Water quality and water management	<ul style="list-style-type: none"> <li>■ In situ water quality downstream of Project footprint (pH, EC, TDS, TSS, DO)</li> <li>■ Monthly water quality parameters in the lake, wetlands and watercourses (pH, EC, TDS, TSS, DO, metals, hydrocarbons)</li> <li>■ Volume of water extracted and treated on site</li> </ul>	Weekly and monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Water quality monitoring</li> <li>■ Waste water treatment</li> </ul>
Reduction in survival/ reproduction Reduced habitat quality	Grey Crowned Crane	Air quality	<ul style="list-style-type: none"> <li>■ No noticeable odours at boundary of Project footprint</li> </ul>	Monthly	CNOOC	<ul style="list-style-type: none"> <li>■ Odour management and detection</li> </ul>
Population influx	Near-shore habitats Escarpment vegetation corridor Wetlands Bugoma CFR Mud Snail Grey Crowned Crane Nahan's Francolin	People management	<ul style="list-style-type: none"> <li>■ No personnel and/or contractors allowed beyond footprint of Project</li> <li>■ Designated no-go areas, e.g., Lake Albert, wetlands, Bugoma Central</li> </ul>	As required	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Inductions for all staff</li> </ul>



## BIODIVERSITY IMPACT ASSESSMENT

Specific impact	Affected ecosystem/species of Concern	Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
	Eastern Chimpanzee		Forest Reserve, wetlands			





### 13.0 OFFSETTING STRATEGY

Possible offset options as part of the mitigation strategy for the Project are outlined on Table 50. Post-offset residual risks are discussed in the sections that follow. It is noted that a biodiversity action plan for the design and implementation of appropriate offsets to ensure no-net-loss of natural habitat, and net gain of critical habitat, must be developed in association with partnering organisations, including the other partners (TUOP and TOTAL) and BLAC. This is outside the current scope of the ESIA, but will be required to demonstrate compliance with Lender standards.

#### 13.1 Construction and Operation Phase Impacts on the Near-Shore Environment of Lake Albert

Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for natural habitat, such as the near-shore habitats of Lake Albert, is a target that CNOOC is working towards to meet IFC requirements, and partner expectations.

Offsetting is required to compensate for adverse residual impacts to natural habitat. A possible offsetting mechanism is supporting improved management of the near-shore aquatic habitats within the CHAA, which should curb degradation of these habitats. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting habitat.

One way to do this is to educate the local population about the potential impacts their activities could have on the near-shore aquatic habitats. For example: the safe handling and disposal of fuels and oils used for out-board motors; and the use of less destructive fishing methods in the near-shore habitats, such as seining, which are known to affect these habitats detrimentally (Wandera and Balirwa 2010).

The mitigation hierarchy is an important process that has been used to minimise impacts to the near-shore habitats. The focus for the continued use of the mitigation hierarchy during construction will be continued development and implementation of mitigation measures through monitoring and adaptive management during construction. Direct footprint impacts have largely been avoided and the level of realisation of impacts from disturbance is uncertain.

As noted above, the focus for reducing impacts to near-shore habitats is mitigation to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to these habitats have been identified above. The goal of these offsets would be to work towards no net loss of natural habitats within in the CHAA. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as possible at this time, which results in a high-risk level for the Project (Table 44).

**Table 44: Post-offset residual risk: construction phase impacts on near-shore environment of L. Albert**

Description of Offsets	Residual Impact	Risk Level for not being successful
Education of the local people about the potential impacts their activities could have on the near-shore aquatic habitats. For example: the safe handling and disposal of fuels and oils used for out-board motors; and the use of less destructive fishing methods in the near-shore habitats, such as seining.	positive	high risk

#### 13.2 Construction and Operation Phase Impacts on Wetlands in the CHAA

The construction and operation of the airstrip, flowlines and in-field roads will be see the direct and indirect loss of natural habitat on the Buhuka Flats; the construction of the Feeder pipeline will also result in direct and





indirect losses of natural wetland habitat. Under PS6, no net loss of natural habitat is required to meet the standards. Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for highly sensitive and unique biodiversity values is a target that CNOOC is working towards to meet IFC requirements, and partner expectations.

Offsetting is required to compensate for loss of natural habitat supported on the Buhuka Flats. A possible offsetting mechanism is the support of improved management of wetlands in the wider Albertine Graben; in particular, known breeding sites for Grey Crowned Cranes and Shoebills. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting the habitat.

One way to do this is to provide assistance to UWA, and other research organisations, with monitoring and conservation of the Grey Crowned Crane and Shoebill.

The mitigation hierarchy is an important process that has been used to minimise impacts to the wetlands of the CHAA. The focus for the continued use of the mitigation hierarchy during construction will be continued development and implementation of mitigation measures through monitoring and adaptive management during construction.

As noted above, the focus for reducing impacts to the wetlands of the CHAA to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, natural vegetation of permanent wetlands and seasonally flooded grasslands will still be lost. Therefore, offsetting options to compensate for that loss have been identified above. The goal of those offsets would be to work towards no net loss of the habitats in the CHAA. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as possible at this time, which results in a high-risk level for the Project (Table 45).

**Table 45: Post-offset residual risk: construction phase impacts on wetlands in the CHAA**

Description of Offsets	Residual Impact	Risk Level for not being successful
Assistance to UWA, and other research organisations, with monitoring and conservation of the Grey Crowned Crane and Shoebill in the local area and their habitats.	positive	high risk

### **13.3 Construction and Operation Phase Impacts on the Mud Snail *G. candida***

Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for highly sensitive and unique biodiversity values, such as the Mud Snail, is a target that CNOOC is working towards to meet IFC requirements, and partner expectations.

Offsetting is required to compensate for adverse residual impacts to Mud Snails. A possible offsetting mechanism is supporting improved management of the near-shore aquatic habitats within the CHAA, which should curb degradation of these habitats. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting habitat.

One way to do this is to educate the local population about the potential impacts their activities could have on the near-shore aquatic habitats. For example: the safe handling and disposal of fuels and oils used for out-board motors; and the use of less destructive fishing methods in the near-shore habitats, such as seining, which are known to affect these habitats detrimentally (Wandera and Balirwa 2010).

The mitigation hierarchy is an important process that has been used to minimise impacts to the Mud Snail. The focus for the continued use of the mitigation hierarchy during construction will be continued development and implementation of mitigation measures through monitoring and adaptive management during construction.







Direct footprint impacts have largely been avoided and the level of realisation of impacts from disturbance is uncertain.

As noted above, the focus for reducing impacts to Mud Snails is mitigation to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to Mud Snails have been identified above. The goal of these offsets would be to work towards no net loss and eventually net gain to Mud Snail habitat and populations in the CHAA. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as possible at this time, which results in a high-risk level for the Project (Table 46).

**Table 46: Post-offset residual risk: construction phase impacts on Mudsnaails in the CHAA**

Description of Offsets	Residual Impact	Risk Level for not being successful
Education of the local people about the potential impacts their activities could have on the near-sure aquatic habitats. For example: the safe handling and disposal of fuels and oils used for out-board motors; and the use of less destructive fishing methods in the near-shore habitats, such as seining	positive	high risk

### 13.4 Construction and Operation Phase Impacts on Bugoma CFR

Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for highly sensitive and unique biodiversity values, such as the Bugoma Central Forest Reserve, is a target that CNOOC is working towards to meet IFC requirements, and partner expectations. Indeed, because the Bugoma Central Forest Reserve is identified as critical habitat under criteria 1 and 4, a net positive impact will be the goal.

Offsetting is required to compensate for adverse residual impacts to Bugoma Central Forest Reserve from induced and cumulative effects (see Chapter 17 Cumulative Impact Assessment).

A possible offsetting mechanism is supporting improved management of the Bugoma Central Forest Reserve, and research on the threatened species that inhabit the forest. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting habitat.

One way to do this is to provide assistance to the NFA for management of the Bugoma Central Forest Reserve, and the UWA, and other research organisations involved with the monitoring and conservation of threatened species.

The mitigation hierarchy is an important process that has been used to minimise impacts to the Eastern Chimpanzee. The focus for the continued use of the mitigation hierarchy during the road upgrade and construction will be continued development and implementation of mitigation measures through monitoring and adaptive management; including suggested measures for the Ugandan Government to apply in the management of the upgrade of the Hoima-to-Ikamiro Road.

As noted above, the focus for reducing impacts to the Bugoma Central Forest Reserve is mitigation to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to the forest have been identified above. The goal of these offsets would be to work towards no net loss of this forest in the CHAA and net positive impact for the reserve. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as unlikely at this time due to increased human pressures around the forest, which results in a high-risk level for the Project (Table 47).





**Table 47: Post offset residual risk: Construction and operation phase impacts on Bugoma CFR**

Description of Offsets	Residual Impact	Risk Level for not being successful
Supporting improved management of the Bugoma Central Forest Reserve.	positive	high risk

### 13.5 Construction and Operation Phase Impacts on Nahan’s Francolin

Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for highly sensitive and unique biodiversity values, such as the Nahan’s Francolin, is a target that CNOOC is working towards to meet IFC requirements, and partner expectations.

Offsetting is required to compensate for adverse residual impacts to Nahan’s Francolin from induced and cumulative effects. Within Bugoma Central Forest Reserve, the population trend of this species appears to be decreasing, with the primary threats thought to be habitat loss through logging and clearance of forest for charcoal burning and agriculture (BirdLife International 2017).

A possible offsetting mechanism is supporting improved management of the Bugoma Central Forest Reserve, and research on the population of Nahan’s Francolin in that forest. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting habitat.

One way to do this is to provide assistance to UWA and other research organisations with monitoring and conservation of the Nahan’s Francolin, and the NFA for management of the Bugoma Central Forest Reserve.

The mitigation hierarchy is an important process that has been used to minimise impacts to the Nahan’s Francolin. The focus for the continued use of the mitigation hierarchy during construction will be continued development and implementation of mitigation measures through monitoring and adaptive management during construction; including suggested measures for the Ugandan Government to apply in the management of the upgrade of the P1 Road.

As noted above, the focus for reducing impacts to Nahan’s Francolin is mitigation to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to Nahan’s Francolin have been identified above. The goal of these offsets would be to work towards no net loss and eventually net gain to Nahan’s Francolin populations in the CHAA. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as possible at this time, which results in a moderate risk level for the Project (Table 48).

**Table 48: Post offset residual risk: construction and operation phase impacts on Nahan’s Francolin**

Description of Offsets	Residual Impact	Risk Level for not being successful
Supporting improved management of the Bugoma Central Forest Reserve, and research on the population of Nahan’s Francolin.	positive	moderate risk

### 13.6 Construction and Operation Phase Impacts on Chimpanzee

Successful application of offsets means that the overall impact level from the Project becomes negligible or positive. A positive overall result for highly sensitive and unique biodiversity values, such as the Eastern Chimpanzee, is a target that CNOOC is working towards to meet IFC requirements, and partner expectations.

Offsetting is required to compensate for adverse residual impacts to Eastern Chimpanzee from induced and cumulative effects. Within Bugoma Central Forest Reserve, the population trend of this species appears to be





decreasing, with the primary threats thought to be habitat loss through logging and clearance of forest for charcoal burning and agriculture, bush meat hunting, and killing of crop raiding individuals (Plumptre et al. 2003, 2010).

A possible offsetting mechanism is supporting improved management of the Bugoma Central Forest Reserve, and research on the Eastern Chimpanzee population in that forest. However, to be considered an offset, this improved management must work to reduce human disturbance that may be affecting habitat.

One way to do this is to provide assistance to UWA and other research organisations involved with the monitoring and conservation of the Eastern Chimpanzee, and the NFA for management of the Bugoma Central Forest Reserve.

The mitigation hierarchy is an important process that has been used to minimise impacts to the Eastern Chimpanzee. The focus for the continued use of the mitigation hierarchy during the road upgrade and construction will be continued development and implementation of mitigation measures through monitoring and adaptive management; including suggested measures for the Ugandan Government to apply in the management of the upgrade of the P1 Road.

As noted above, the focus for reducing impacts to Eastern Chimpanzees is mitigation to lessen various types of disturbance that may occur. Optimising mitigation will reduce the need for offsetting. However, offsetting options to compensate for impacts to Eastern Chimpanzees have been identified above. The goal of these offsets would be to work towards no net loss and eventually net gain to Eastern Chimpanzee populations in the CHAA. Although reasonable mechanisms exist, there is still uncertainty associated with offsetting because proposed actions require third-party participation beyond the control of the Project. Offset feasibility is classed as possible at this time, which results in a moderate risk level for the Project (Table 49).

**Table 49: Post offset residual risk: construction and operation phase impacts on Eastern Chimpanzee**

Description of Offsets	Residual Impact	Risk Level for not being successful
Supporting improved management of the Bugoma Central Forest Reserve, and research on the population of Nahan’s Francolin.	positive	moderate risk





## BIODIVERSITY IMPACT ASSESSMENT

**Table 50: Possible offset options as part of the mitigation strategy for the Project**

Offset required	Mechanism	Responsibility	Details
Natural habitat - near-shore aquatic Critical habitat – Mud Snail	<ul style="list-style-type: none"> <li>Improved management of the near-shore aquatic habitats within the CHAA</li> </ul>	CNOOC	<ul style="list-style-type: none"> <li>Education programme for the local population about the potential impacts their activities could have on the near-shore aquatic habitats</li> <li>The safe handling and disposal of fuels and oils used for out-board motors</li> <li>The use of less destructive fishing methods in the near-shore habitats, such as seining</li> </ul>
Natural habitat - vegetation of the escarpment	<ul style="list-style-type: none"> <li>Improved management of the wider Murchison Falls National Park-Budongo-Bugoma-Kagombe-Itwara Forest Reserves-Semliki/Toro Wildlife Reserve corridor</li> </ul>	CNOOC, TOTAL, Tullow	<ul style="list-style-type: none"> <li>Provision of assistance to UWA, and other research organisations, in forming and collaborating in a committee to devise long-term management goals and measures for the wildlife corridor</li> <li>Action plan to ensure that the management goals and measures devised by the committee are actioned, implemented and monitored on the ground</li> </ul>
Natural habitat - wetlands	<ul style="list-style-type: none"> <li>Improved management of wetlands in the wider Albertine Graben; in particular, known breeding sites for Grey Crowned Cranes and Shoebills</li> </ul>	CNOOC	<ul style="list-style-type: none"> <li>Provision of assistance to UWA, and other research organisations, with monitoring and conservation of the Grey Crowned Crane and Shoebill</li> </ul>
Natural habitat – Bugoma Central Forest Reserve Critical habitat – Nahan’s Francolin Critical habitat – Eastern Chimpanzee	<ul style="list-style-type: none"> <li>Improved management of the Bugoma Central Forest Reserve, and research on the threatened species that inhabit the forest</li> </ul>	CNOOC	<ul style="list-style-type: none"> <li>Assistance to the NFA for management of the Bugoma Central Forest Reserve, and the UWA, and other research organisations involved with the monitoring and conservation of threatened species.</li> </ul>



### 14.0 REFERENCES

- AECOM Limited (2012). Report on the Environmental Baseline in Exploration Area 2, Tullow Uganda Operations Pty.
- AECOM Limited (2013). Report on Biodiversity Conservation Opportunities in the Albertine Rift, Tullow Uganda Operations Pty.
- African Regional Workshop (Conservation & Sustainable Management of Trees, Zimbabwe, July 1996) 1998.
- Cordia millenii*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.
- Agamy E (2013). Sub-chronic exposure to crude oil, dispersed oil and dispersant induces histopathological alterations in the gills of the juvenile rabbit fish (*Siganus canaliculatus*). *Ecotoxicology & Environmental Safety*. 92:180-190
- Air Water Earth Ltd (2008a). Environmental and Social Impact Assessment for the KF2 well. Heritage Oil Plc.
- Air Water Earth Ltd (2008b). Environmental and Social Impact Assessment for the KF3 well. Heritage Oil Plc.
- Air Water Earth Ltd (2013a). Environmental and Social Impact Assessment for Camp and Yard upgrade. CNOOC.
- Air Water Earth Ltd (2013b). Airstrip rehabilitation Project Brief, CNOOC.
- Air Water Earth Ltd (2014a). Environmental and Social Impact Assessment for KF4 well, CNOOC.
- Air Water Earth Ltd (2014b). Jetty rehabilitation Project Brief, CNOOC.
- Air Water Earth Ltd (2014c). Escarpment Road ESIA, CNOOC.
- Akinyi E, Awaïss A, Azeroual A, Getahun A, Lalèyè P and Twongo T (2010a). *Alestes baremoze*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Akinyi E, Bouso T, Getahun A, Hanssens M, Lalèyè P, Moelants T and Twongo T (2010b). *Distichodus niloticus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Akinyi E, Azeroual A, Entsua-Mensah M, Getahun A, Lalèyè P and Twongo T (2010c). *Alestes dentex*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 19 December 2017.
- Alliance for Zero Extinction (2010). <http://www.zeroextinction.org/>; downloaded, 26 March 2014.
- American Public Health Association (1992). Standard Methods for the Examination of Water and Waste Water. 18th edition. ALPHA-AWWA-WEF, Washington.
- Araujo CVJ, Blasco J and Moreno-Garrido I (2012). Measuring the avoidance behaviour shown by the snail *Hydrobia ulvae* exposed to sediment with a known contamination gradient. *Ecotoxicology* (2012) 21:750–758.
- Archibald GW, Meine CD and Garcia EFJ (2013). Grey Crowned Crane (*Balearica regulorum*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA and de Juana E (eds). Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/53550> on 23 December 2017).
- Awaïss A, Azeroual A, Getahun A, Lalèyè P, Moelants T and Twongo T (2010). *Synodontis nigrita*. The IUCN





Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 19 December 2017.

Ayebare S, Ponce-Reyes R, Segan DB, Watson JEM, Possingham HP, Seimon A, and Plumtre AJ (2013) Identifying climate resilient corridors for conservation in the Albertine Rift. Unpublished Report by the Wildlife Conservation Society to MacArthur Foundation.

Azeroual A, Getahun A, Lalèyè P, Moelants T and Twongo T (2010a). *Brycinus macrolepidotus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.

Azeroual A, Bayona J, Entsua-Mensah M, Getahun A and Lalèyè P (2010b). *Citharinus latus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.

Azeroual A, Bills R, Cambray J, Getahun A, Hanssens M, Marshall B, Moelants T and Tweddle D (2010c). *Hydrocynus vittatus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.

Azeroual A, Entsua-Mensah M, Getahun A, Lalèyè P and Kazembe J (2010d). *Malapterurus electricus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.

Azeroual A, Bouso T, Getahun A, Lalèyè P, Moelants T and Twongo T (2010e). *Schilbe mystus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 19 December 2017.

Begon M, Harper JL and Townsend CR (1990). *Ecology: Individuals, Populations and Communities*. Blackwell Scientific Publications, Boston.

Bennun L and Njoroge P (eds) (1996). *Birds to Watch in East Africa: A Preliminary Red Data List*. Research Reports of the Centre for Biodiversity, National Museums of Kenya, Ornithology 23.

Bennun L, Njoroge P and Pomeroy D (2000). Birds to watch: a Red Data List for east Africa. *Ostrich* 71: 310-314.

Beresford AE, Buchanan GM, Donald PF, Butchart SHM, Fishpool LDC and Rondinini C (2011). Poor overlap between the distribution of Protected Areas and globally threatened birds in Africa. *Animal Conservation*, 14(2):99-107.

Berthinussen A and Altringham J (2012). The Effect of a Major Road on Bat Activity and Diversity. *Journal of Applied Ecology*. 49: 82-89.

Beveridge MCM (1996). *Cage Aquaculture*. Second Edition. Blackwell Science Ltd, Oxford, UK.

BirdLife International (1998). *Endemic Bird Areas of the World: Priorities for Biodiversity Conservation*, Cambridge.

BirdLife International (2017). IUCN Red List for birds. Downloaded from <http://www.birdlife.org> on 15 December 2017.

Blanc J (2008). *Loxodonta africana*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.

Board of Trustees of the Royal Botanic Gardens, Kew (2014). *Tamarindus indica* – species information. <http://www.kew.org/science-conservation/plants-fungi/tamarindus-indica-tamarind>. Accessed 11 December 2017.

Brehme CS, Tracey JA, McClenaghan LR, and Fisher RN (2013). Permeability of roads to movement of scrubland lizards and small mammals. *Conservation Biology*. 27(4):710-720



- Brooks JL (1957). The systematics of North American Daphnia. *Memoirs of the Connecticut Academy of Arts and Sciences*, 13:1-18.
- Bruce-White C and Shardlow M (2011). A Review of the Impact of Artificial Light on Invertebrates. Buglife – The Invertebrate Conservation Trust Bunn SE and Arlington AH (2002). Basic Principles and Ecological Consequences of Altered Flow Regimes for Aquatic Biodiversity. *Environmental Management* 30(4): 492-507.
- Business and Biodiversity Offsets Programme (2012). Standard on Biodiversity Offsets. BBOP, Washington, D.C.
- Buttynski TM, Kingdon J and Kalina J (2013). *Mammals of Africa Volume II: Primates*; Bloomsbury Publishing, London
- Byaruhanga A, Kasoma P and Pomeroy D (2001). *Important Bird Areas in Uganda*. East Africa Natural History Society, Kampala.
- Campbell LM, Wandera SB, Thacker RJ, Dixon DG and Hecky RE (2005). Trophic niche segregation in the Nilotic ichthyofauna of Lake Albert (Uganda, Africa). *Environmental Biology of Fishes*, 74:247–260.
- Carder N and Tindimubona L (2002). *Butterflies of Uganda: a field guide to butterflies and silk moths from the collection of the Uganda Society*. The Uganda Society, Cornell University.
- Carvalho P, Thomaz SM, Kobayashi JT and Bini LM (2013). Species richness increases the resilience of wetland plant communities in a tropical floodplain *Austral Ecology*. 38(5):592-598.
- Channing A and Howell KM (2006). *Amphibians of East Africa*, Cornell University Press.
- Chapman CA and Lambert JE (2000). Habitat alteration and the conservation of African primates: case study of Kibale National Park, Uganda. *American Journal of Primatology*. 50:169-185.
- Chapman CA, Bonnell TR, Gogarten JF, Lambert, JE, Omeja PA, Twinomugisha D, Wasserman MD and Rothman JM (2013). Are primates ecosystem engineers? *International Journal of Primatology* 34:1-14.
- Chen Y, Huang T and Shen Y (2013). Evaluation of ground vibration induced by rail systems. *Noise Control Engineering Journal* 61(2): 145-158.
- CITES (2017). The Convention on International Trade in Endangered Species of Wild Fauna and Flora Secretariat. <<https://www.cites.org/eng/disc/sec/index.php>>. Accessed 15 Dec 2017.
- Clausnitzer V (2002). The identification key of East African dragonflies. Draft version.
- Clausnitzer V (2010a). *Agriocnemis palaeforma*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 21 December 2017
- Clausnitzer V (2010b). *Tetrathemis corduliformis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 21 December 2017
- Claustinitzer V and Kityo R (2001). Altitudinal distribution of rodents (Muridae and Gliridae) on Mt. Elgon, Uganda. *Tropical Zoology*, 14:95-118.
- Clausnitzer V, Suhling F and Dijkstra K-DB (2010). *Mesocnemis singularis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.
- CNOOC Uganda Limited (2014a). Biodiversity Management Specification Rev A, QHSE CUL-QHSE-L3 (GE) - 057.
- CNOOC Uganda Limited (2014b). Natural Resource Management Specification Rev A, CUL-QHSE-L3 (GE)-051.





- Conservation International (2014). Eastern Afromontane Biodiversity Hotspot. <http://www.cepf.net/resources/hotspots/africa/Pages/Eastern-Afromontane.aspx>. Accessed, 7 December 2017.
- Convention on the Conservation of Migratory Species of Wild Animals (2014). [www.cms.int](http://www.cms.int). Accessed 18 December 2017.
- Convention on International Trade in Endangered Species of Wild Fauna and Flora Secretariat (2014). <http://www.cites.org/>. Accessed 12 December 2017.
- Cooley HS, Wielgus RB, Koehler GM, Robinson HS and Maletzke BT (2009). Does hunting regulate cougar populations? A test of the compensatory mortality hypothesis. *Ecology* 90(10):2913–2921
- Cotterill FPD (2013). *Tadarida ventralis* Giant Free-tailed Bat; pp 539-540. In: Happold M and Happold DCD (eds). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Crocodile Specialist Group (1996). *Crocodylus niloticus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 12 December 2017.
- Cumberlidge N (2008). *Potamonautes bipartitus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Davenport TRB (1996). *The Butterflies of Uganda - An Annotated Checklist*. Uganda Forest Department, Kampala, Uganda. pp 48.
- Davenport TRB (2003). *Endemic Butterflies of the Albertine Rift – an annotated checklist*. Wildlife Conservation Society, Mbeya, Tanzania.
- Dean WJR (2000). Alien birds in Southern Africa: what factors determine success? *South African Journal of Science* 96: 9-14.
- de Blois S, Domon G and Bouchard A (2002). Landscape issues in plant ecology. *Ecography* 25(2):244-256.
- Delince G (1992). *The ecology of the fish pond ecosystem with special reference to Africa*. Developments in Hydrobiology 72, Kluwer Academic Publishers.
- de Moor IJ, Day JA and de Moor FC (eds) (2003). *Freshwater Invertebrates of South Africa*. Vol. 8: Insect II. Prepared for the Water Research Commission, Pretoria.
- Dietz C, von Helversen O and Nill D (2009). *Bats of Britain, Europe and Northwest Africa*. A & C Black Publishers Ltd, London.
- Ehrlich PR, Breedlove DE, Brussard PE and Sharp MA (1972). Weather and the "regulation" of subalpine butterfly populations. *Ecology* 53:243-247.
- Emerton L and Muramira E (1999). *Uganda Biodiversity: Economic Assessment*. IUCN, Switzerland.
- Environmental Assessment Consult Ltd (2013). *Environmental and Social Impact Assessment for 2D Seismic Survey*, CNOOC.
- Environmental Assessment Consult Ltd (2014). *Environmental and Social Impact Assessment for escarpment road*, CNOOC.
- Environment Protection and Heritage Council and the Natural Resource Management Ministerial Council (2011). *National guidance for the management of acid sulfate soils in inland aquatic ecosystems*, Canberra, ACT.
- Eskew EA, Price SJ and Dorcas ME (2012). Effects of river-flow regulation on anuran occupancy and abundance in riparian zones. *Conservation Biology* 26(3): 504-512.
- Fahrig L (2001). How Much Habitat is Enough? *Biological Conservation* 100:65-74.





- Fahrig L (2003). Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology and Systematics* 34: 487-515.
- Ferraro SP and Cole FA (1992). Taxonomic level is sufficient for assessing a moderate impact on macrobenthic communities in Puget Sound, Washington, USA. *Canadian Journal of Aquatic Science*, 49:1184-1188.
- Findlay SCT and Bourdages J (2000). Response time of wetland biodiversity to road construction on Adjacent Lands. *Conservation Biology*. 14(1):86-94.
- Forest Department (2002). National Biomass Study, Technical Report of 1996-2002. Ministry of Water Lands and Environment, Kampala.
- Forman RTT (1995). *Land Mosaics: The Ecology of Landscapes and Regions*. Cambridge University Press. Cambridge, U.K.
- Formenty P, Karesh W, Froment JM and Wallis J (2003). Infectious Diseases in West Africa: A Common Threat to Chimpanzees and Humans. In: Kormos R, Boesch C, Bakarr MI and Butynski TM. *West African Chimpanzees. Status Survey and Conservation Action Plan IUCN/SSC Primate Specialist Group*. IUCN, Gland, Switzerland and Cambridge, UK. ix + 219 pp.
- Fujita S (2011). Health Monitoring. In: Matsuzawa T, Humle T and Sugiyama Y. *The Chimpanzees of Bossou and Nimba*. Springer. 465 pp.
- Gaston KJ (1991). The magnitude of global insect species richness. *Conservation Biology* 5:283-296.
- Gaston KJ and Lawton JH (1988). Patterns in distribution and abundance of insect populations. *Nature* 331:709-712.
- Gillison AN and Brewer KRW (1985). The use of gradient directed transects or gradsects in natural resource surveys. *Journal of Environmental Management* 20:103-127
- Gleeson J and Gleeson D (2012). *Reducing the Impacts of Development on Wildlife*. CSIRO Publishing, Collingwood.
- GBIF (2017). Global Biodiversity Information Facility. GBIF.org (13th November 2017) GBIF Occurrence. Download <https://doi.org/10.15468/dl.unota1>; Occurrence Download <https://doi.org/10.15468/dl.0igvvx>; Occurrence Download <https://doi.org/10.15468/dl.l7ljk9>.
- Global Climate Change Alliance (2012). *Adaption for climate change in Uganda*. Retrieved June 2014, from Global Climate Change Alliance: <http://www.gcca.eu/national-programmes/africa/gcca-uganda>.
- Godar J, Tizado E, Pokorny B and Johnson J (2012). Typology and characterization of Amazon colonists: a case study along the Transamazon Highway. *Human Ecology: An Interdisciplinary Journal*. 40(2):251 - 267.
- Golder Associates (Pty) Ltd (2014a). *Comments and Response Register - Public participation and stakeholder engagement for the Environmental and Social Impact Assessment for the proposed Kingfisher Field Development of CNOOC Uganda Ltd on the shore of Lake Albert, Hoima District, and along the proposed pipeline route*.
- Golder Associates (Pty) Ltd (2014c). *Scoping Report - Environmental and Social Impact Assessment for Kingfisher Development Area*.
- Golder Associates (Pty) Ltd (2018). *Influx Management Plan - Environmental and Social Impact Assessment for Kingfisher Development Area*.
- Gotelli NJ and Colwell RK (2011). Estimating Species Richness. Pages 39-54 in Magurran AE and McGill BJ (eds). *Biological Diversity: Frontiers in Measurement and Assessment*. Oxford University Press, New York.
- Government of Uganda (1995). *Wetlands Policy 1995*. Ministry of Natural Resources, Kampala, Uganda.



- Green J (1971). Associations of Cladocera in the zooplankton of the lake source of the White Nile. *Journal of Zoology*, 165: 373–414.
- Green RE, Taggart MA, Das D, Pain DJ, Kumar CS, Cunningham AA and Cuthbert R (2006). Collapse of Asian vulture populations: risk of mortality from residues of the veterinary drug diclofenac in carcasses of treated cattle. *Journal of Applied Ecology*, 43:949–956.
- Greenwood PH (1966). *The Fishes of Uganda*. The Uganda society, Kampala. Second Edition. 131 pp.
- Gregory RD, Gibbons DW and Donald PF (2005). Bird census and survey techniques. In: Sutherland WJ, Newton I and Green RE (eds). *Bird Ecology and Conservation: a handbook of techniques*, Oxford University Press, Oxford.
- Harris R (2013). The conservation of Accipitridae vultures of Nepal: a review. *Journal of Threatened Taxa*, 5(2):3603–3619.
- Happold M (ed) (2013a). *Mammals of Africa Volume III: Rodents, Hares and Rabbits*; Bloomsbury Publishing, London.
- Happold M (2013b). Genus *Miniopterus* Long-fingered Bats; pp 771-772. In: Happold M and Happold DCD (eds) (2013). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Happold M (2013c). *Tadarida trevori* Trevor's Free-tailed Bat; pp 537-536. In: Happold M and Happold DCD (eds) (2013). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Happold M (2013d). *Glauconycteris humeralis* Spotted Butterfly Bat; pp 570-571. In: Happold M and Happold DCD (eds) (2013). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Happold M and Happold DCD (eds) (2013). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Heckey RE (1993). The eutrophication of Lake Victoria. *Verhandlungen des Internationalen Verein Limnologie*, 25: 39-48.
- Henschel P, Breitenmoser-Wursten C and Sogbohossou E (2008). *Caracal aurata*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Henschel P, Hunter L, Breitenmoser U, Purchase N, Packer C, Khorozyan I, Bauer H, Marker L, Sogbohossou E and Breitenmoser-Wursten C (2008). *Panthera pardus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Heyer, WR, Donnelly M, McDiarmid RW, Hayek CL and Foster MS (eds) (1994). *Measuring and Monitoring. Biological Diversity Standard Methods for Amphibians*. Smithsonian Institution Press, Washington, DC.
- Hill D, Fasham, M, Tucker, G, Shewry, M and Shaw, P (2005). *Handbook of Biodiversity Methods*. Cambridge University Press.
- Hill D, Gorman M, Ounsted M, Keller V, Barker MAA and Hockin D (1992). Examination of the effects of disturbance on birds with reference with its importance in ecological assessments. *Journal of Environmental Management* 36(4): 253
- Hilton GM and Cuthbert RJ (2010). The catastrophic impact of invasive mammalian predators on birds of the UK Overseas Territories: a review and synthesis. *The International Journal of Avian Science* 152: 443- 458.
- Hockings K and Humle T (2009). *Best Practice Guidelines for the Prevention and Mitigation of Conflict Between Humans and Great Apes*. Gland, Switzerland: IUCN/SSC Primate Specialist Group. 40 pp.





Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M, Naeem S, Schmid B, Setälä H, Smstad AJ, Vandermeer J and Wardle DA (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecological Monographs* 75:3-35.

Howard PC, Viskanic P, Davenport TRB, Kigenyi FW, Baltzer M, Dickinson CJ, Lwanga JS, Matthews RA, and Balmford A (1998) Complementarity and the use of indicator groups for reserve selection in Uganda. *Nature* 394:472-475.

Hunter L, Henschel P and Ray JC (2013). *Panthera pardus* Leopard; pp159-168. Kingdon J and Hoffmann M (eds). *Mammals of Africa Volume V: Pigs, Carnivores, Pangolins, Equids and Rhinoceroses*; Bloomsbury Publishing, London.

Hutterer R (2013). *Crocidura selina* Ugandan Lowland Shrew; pp 130-131. In: Happold M and Happold DCD (eds). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.

Hutterer R, Oguge N, and Howell K (2008). *Crocidura selina*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.

International Council on Mining and Metals (2010). Good Practice Guidance for Mining and Biodiversity, <http://www.icmm.com/publications>.

International Erosion Control Association (2008). Best Practice Erosion and Sediment Control – Book 2. IECA, Australasia.

International Finance Corporation (2007a). Environmental, Health, and Safety Guidelines - General EHS Guidelines: Environmental - Air Emissions and Ambient Air Quality, <http://www.ifc.org>

International Finance Corporation (2007b). Environmental, Health, and Safety Guidelines for Onshore Oil and Gas Development, <http://www.ifc.org>

International Finance Corporation (2012a). Performance Standard 6: Biodiversity Conservation and Sustainable Management of Living Natural Resources. World Bank Group.

International Finance Corporation (2012b). Guidance Note 6: Biodiversity Conservation and Sustainable Management of Living Natural Resources. World Bank Group.

International Finance Corporation (2012c). Performance Standard 1: Assessment and Management of Environmental and Social Risks and Impacts. World Bank Group.

International Finance Corporation (2013). Good Practice Handbook: Cumulative Impact Assessment and Management – Guidance for the Private Sector in Emerging Markets. World Bank Group.

International Petroleum Industry Environmental Conservation Association (2005). A guide to developing biodiversity action plans for the oil and gas sector. <http://www.iecea.org>.

International Petroleum Industry Environmental Conservation Association (2007). An ecosystem approach to oil and gas industry biodiversity conservation. <http://www.iecea.org>.

International Petroleum Industry Environmental Conservation Association (2010). Alien invasive species and the oil and gas industry Guidance for prevention and management. <http://www.iecea.org>.

International Union for the Conservation of Nature (2001). IUCN Red List Categories and Criteria. Version 3.1 (second edition). <http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria>.

International Union for the Conservation of Nature (2010). Key Biodiversity Areas. [http://iucn.org/about/union/secretariat/offices/iucnmed/iucn\\_med\\_programme/species/key\\_biodiversity\\_areas/](http://iucn.org/about/union/secretariat/offices/iucnmed/iucn_med_programme/species/key_biodiversity_areas/). Downloaded on 26 March 2014.

International Union for the Conservation of Nature (2017). IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 15 December 2017.





International Union for the Conservation of Nature (2014). Integrated Biodiversity Assessment Tool. <[www.ibatforbusiness.org](http://www.ibatforbusiness.org)>. Downloaded on 1 December 2017.

International Union for the Conservation of Nature Species Survival Commission Amphibian Specialist Group (2014a). *Amietophrynus vittatus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 12 December 2017.

International Union for the Conservation of Nature Species Survival Commission Amphibian Specialist Group (2014b). *Amietia desaegeri*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 12 December 2017.

International Union for the Conservation of Nature Species Survival Commission East African Plants Red List Authority (2013a). *Afrothismia winkleri*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 20 December 2017.

International Union for the Conservation of Nature Species Survival Commission East African Plants Red List Authority (2013b). *Afrothismia winkleri* var. *budongensis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 20 December 2017.

Isabirye-Basuta G and Kasenene JM (1987). Small rodent populations in selectively felled and matured tracts of Kibale Forest, Uganda. *Biotropica* 19:260–266.

Isabirye-Basuta GM and Lwanga JS (2008). Primate populations and their interactions with changing habitats. *International Journal of Primatology*. 29:35-48.

Jenkins PT (2013). Invasive animals and wildlife pathogens in the United States: the economic case for more risk assessments and regulation. *Biological Invasions* 15: 243-248.

Jetz W, Thomas GH, Joy JB, Redding DW, Hartmann K and Mooers AO (2014). Global distribution and conservation of evolutionary distinctness in birds. *Current Biology* 24(9):9191-930.

Jetz W, Steffen J and Linsenmair KE (2003). Effects of light and prey availability on nocturnal, lunar and seasonal activity of tropical nightjars. *Oikos* 103:627–639.

John DM, Whitton BA and Brook AJ (2002). The freshwater algal flora of the British Isles: an identification guide to freshwater and terrestrial algae. Cambridge University Press, Edinburgh, UK.

Kalema J (2005). Diversity and distribution of vascular plants in wetland and savanna Important Bird Areas of Uganda. Makerere University, Kampala, Uganda.

Kalema J and Beentje HJ (2012). Conservation checklist of the trees of Uganda. Royal Botanic Gardens, Kew, England.

Kawata M, Sawada H, Hiroshi Y, Echenique-Díaz L and Ishibashi Y(2005). The effects of spatial habitat structure on the evolution of density-dependent growth and reproduction in freshwater snails. *Hydrobiologia*. 533(1-3):229-241.

Kemp AC and Christie DA (2013). Hooded Vulture (*Necrosyrtes monachus*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA and de Juana E (eds). Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52995> on 23 December 2017).

Kemp AC, Christie DA, Kirwan GM and Garcia EFJ (2014). African White-backed Vulture (*Gyps africanus*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA and de Juana E (eds). Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52995> on 23 December 2017).

Kemp AC, Kirwan GM, Christie DA and Marks JS (2014). Secretarybird (*Sagittarius serpentarius*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA and de Juana E (eds). Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52995> on 23 December 2017).

Kent M and Coker P (1992). Vegetation description and analysis. John Wiley & Sons, Chichester, England.  
Kielland J (1990). Butterflies of Tanzania. Hill House, Melbourne.



- Kingdon J and Hoffmann M (eds) (2013a). *Mammals of Africa Volume V: Pigs, Carnivores, Pangolins, Equids and Rhinoceroses*; Bloomsbury Publishing, London.
- Kingdon J and Hoffmann M (eds) (2013b). *Mammals of Africa Volume VI: Pigs, Hippopotamus, Chevrotian, Girraffes, Deer and Bovids*; Bloomsbury Publishing, London.
- Kingdon J and Hoffmann M (2013c). *Phataginus tricuspis* White-bellied Pangolin. In: Kingdon J and Hoffmann M (eds), *The Mammals of Africa. Volume V: Carnivores, Pangolins, Equids, Rhinoceroses*, Bloomsbury Publishing, London.
- Kingdon J and Hoffmann M (eds) (2013d), *The Mammals of Africa. Volume V: Carnivores, Pangolins, Equids, Rhinoceroses*, Bloomsbury Publishing, London.
- Kingdon J, Happold D, Hoffman M, Butynski, T, Happold M and Kalina N (eds) (2013). *Mammals of Africa Volume I: Introductory Chapters and Afrotheria*; Bloomsbury Publishing, London.
- Kingdon J, Hoffmann M and Hoyt R (2013). *Smutsia gigantea* Giant Ground Pangolin; pp 396-399. In: Kingdon J and Hoffmann M (eds), *The Mammals of Africa. Volume V: Carnivores, Pangolins, Equids, Rhinoceroses*, Bloomsbury Publishing, London.
- Kityo R, Plumptre AJ, Kerbis JP, Pilgrim J and Moyer D (2003). *Mammals*. pp 20-29. In: Plumptre AJ, Behangana M, Davenport T, Kahindo C, Kityo R, Ndomba E, Nkuutu D, Owuini I, Ssegawa P and Eilu G (eds). *The Biodiversity of the Albertine Rift. Albertine Rift Technical Reports No. 3. Wildlife Conservation Society, New York.*
- Klingel H (2013). *Hippopotamus amphibius* Common Hippopotamus. In: Kingdon J and Hoffmann M (eds) (2013b). *Mammals of Africa Volume VI: Pigs, Hippopotamus, Chevrotian, Girraffes, Deer and Bovids*; Bloomsbury Publishing, London.
- Kociolek AV, Clevenger AP, St Clair CC and Proppe DS (2011). Effects of road networks on bird populations. *Conservation Biology*. 25(2):241-249.
- Kolowski JM and Alonso A (2010). Density and activity patterns of ocelots (*Leopardus pardalis*) in northern Peru and the impact of oil exploration activities. *Biological Conservation*, 143: 917-925.
- Komarek J and Anagnostidis K (1999). *Cyanoprokaryota, 1. TeilChroococcales*. Jena: Gustav Fischer. pp 225-236.
- Kristensen TK and Stensgaard AS (2010). *Gabbiella africana*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 27 December 2017
- Kupferberg SJ, Palen WJ, Lind AJ, Bobzien S, Catenazzi A, Drennan J and Power ME (2012). Effects of flow regimes altered by dams on survival, population declines, and range-wide losses of California riverbreeding frogs. *Conservation Biology* 26(3) 513-524.
- Kusnierz PC, Stimmell SP and Leonard JBK (2014). Migration, Size, and Age Structure of Brook Trout (*Salvelinus fontinalis*) from Two Lake Superior Tributaries. *American Midland Naturalist*, 172(1):119- 130.
- Kyambadde R (2010a). *Gabbiella candida*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Kyambadde R (2010b). *Bellamya rubicunda*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Lalèyè P, Moelants T and Olaosebikan BD (2010). *Clarias camerunensis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 18 December 2017.
- Lamprey RH (2009). *BAT/TBA Uganda Forest Regeneration Project – Phase 2: Land Cover Land Cover Analysis of Survey Areas*.



Landsberg F, Treweek J, Stickler MM, Henninger N and Venn O (2013). Weaving ecosystem services into impact assessment: A Step-By-Step Method. World Resources Institute, Washington DC.

Langdale-Brown I, Osmaston HA and Wilson JG (1964). The Vegetation of Uganda. Government Printer, Entebbe.

Langen T, Gunson K, Scheiner C and Boulerice J (2012). Road mortality in freshwater turtles: identifying causes of spatial patterns to optimize road planning and mitigation. *Biodiversity & Conservation*. 21(12):3017-3034.

Langhammer PF, Bakarr MI, Bennun LA, Brooks TM, Clay RP, Darwall W, De Silva N, Edgar GJ, Eken G, Fishpool LDC, Fonseca GAB, Foster MN, Knox DH, Matiku P, Radford EA, Rodrigues ASL, Salaman P, Sechrest W and Tordoff AW (2007). Identification and Gap Analysis of Key Biodiversity Areas: Targets for Comprehensive Protected Area Systems. Gland, Switzerland: IUCN.

Lansdown RV, Knees SG and Patzelt A (2013). *Polypogon schimperianus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 20 December 2017.

Largen M and Howell K (2004). *Ptychadena christyi*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 12 December 2017.

Larsen TB (1991). *Butterflies of Kenya and Their Natural History*, Oxford University Press, Oxford.

Larsen TB (2005). *Butterflies of West Africa* 2 vols, 596 pp. Apollo Books.

Larsen TB (2011a). *Bicyclus jefferyi*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.

Larsen T (2011b). *Eurema brigitta*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.

Larsen TB (2011c). *Junonia oenone*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.

Larsen TB (2011d). *Zizina antanossa*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.

Larsen TB (2011e). *Micropentila bunyoro*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 21 December 2017.

Leblond M, Dussault C, Ouellet JP and Hayssen V (2013). Avoidance of roads by large herbivores and its relation to disturbance intensity. *Journal of Zoology*. 289(1):32-40.

Lehman JT, Litt AH, Mugidde R and Lehman DA (1998). Nutrients and Plankton biomass in the Rift Lake sources of the White Nile: Lakes Albert and Edward. In: Lehman JT (ed.) *Environment Changes and Response in East African Lakes*. Netherlands, Kluwer Academic Publishers.

Lewinsohn TM, Novotny V and Basset Y (2005). Insects on plants: diversity of herbivore assemblages revisited. *Annual Review of Ecology, Evolution & Systematics*, 36(1):597-620.

Lewison R and Oliver W (IUCN SSC Hippo Specialist Subgroup) (2008). *Hippopotamus amphibius*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 13 December 2017.

Lonsdorf E, Murray C, Lonsdorf E, Travis D, Gilby I, Chosy J, Goodall J and Pusey A (2011). A retrospective analysis of factors correlated to Chimpanzee (*Pan troglodytes schweinfurthii*) respiratory health at Gombe National Park, Tanzania. *EcoHealth*. 8(1):26-35.

Lowe S, Browne M, Boudjelas S and De Poorter M (2000). 100 of the World's Worst Invasive Alien Species. A selection from the Global Invasive Species Database. The Invasive Species Specialist Group (ISSG) a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN), 12 pp.



- Lussier SM, Enser RW, Dasilva SN, Charpentier M (2006). Effects of habitat disturbance from residential development on breeding bird communities in riparian corridors. *Environmental Management* 38(3): 504-521.
- Lytle DA and Poff NL (2004). Adaptations to natural flow regimes. *Trends in Ecology and Evolution* 19(2): 94-100.
- Madsen J (1998). Experimental refuges for migratory waterfowl in Danish wetlands. I. Baseline assessment of the disturbance effects of recreational activities. *Journal of Applied Ecology* 35:386–397.
- Mammides C, Cords M, and Peters MK (2009). Effects of habitat disturbance and food supply on population densities of three primate species in the Kakamega Forest, Kenya. *African Journal of Ecology*. 47(1):87-96.
- Mandahl-Barth G (1954). *The Freshwater Molluscs of Uganda and Adjacent Territories*. Danmarks Akvarium. Tervuren. 206pp.
- Margules CR and Pressey RL (2000). Systematic conservation planning. *Nature* 405: 243-253.
- Martínez-Vilalta A, Motis A and Kirwan GM (2014). Madagascar Pond-heron (*Ardeola idae*). In: del Hoyo J Elliott A, Sargatal J, Christie DA and de Juana E (eds). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/52702> on 29 December 2017).
- Martin TE and Blackburn GA (2010). Impacts of tropical forest disturbance upon avifauna on a small island with high endemism: implications for conservation. *Conservation & Society* 8(2):127-139.
- McDiarmid RW, Foster MS, Guyer C, Gibbons JW and Chernoff N (2012). *Reptile Biodiversity – Standard Methods for Inventory and Monitoring*. University of California Press, Berkeley.
- McGowan PJK and de Juana E (1994). Nahan's Francolin (*Francolinus nahani*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA and de Juana E (eds) (2013). *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona. (retrieved from <http://www.hbw.com/node/53406> on 23 December 2017).
- MvLennan MR (2008). Beleaguered chimpanzees in the agricultural district of Hoima, Western Uganda. *Primate Conservation*. 23:45-54.
- Meisingset EL, Loe LE, Brekkum Ø, Van Moorter B and Mysterud A (2013). Red Deer habitat selection and movements in relation to roads. *Journal of Wildlife Management*. 77(1):181-191.
- Mellin C, Bradshaw CJA, Fordham DA and Caley MJ (2014). Strong but opposing beta-diversity-stability relationships in coral reef fish communities. *Proceedings of the Royal Society B Biological Sciences* 281: 20131993.
- Merritt RW and Cummins KW (eds.) (1984). *An Introduction to the Aquatic Insects of North America*. Second Edition. Dubuque, IA: Kendall/Hunt Publishing Company. 722 pp
- Mickleburgh S, Hutson AM, Bergmans W, Fahr J and Racey PA (2008). *Eidolon helvum*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Mickleburgh S, Hutson AM, Bergmans W, Cotterill FPD and Fahr J (2008a). *Tadarida trevori*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Mickleburgh S, Hutson AM, Bergmans W and Cotterill FPD (2008b). *Tadarida ventralis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Mickleburgh S, Hutson AM, Bergmans W, Fahr J and Taylor PJ (2008c). *Otomops martiensseni*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Millennium Ecosystem Assessment (2005). *Ecosystems and Human Well-being: A Framework for Assessment*. Island Press



- Miller P and Miller K (2003). East African Dragonflies. The East African Natural History Society. Ministry of Energy and Mineral Development (2013). Strategic Environmental Assessment (SEA) of Oil and Gas Activities in the Albertine Graben, Uganda. Ministry of Energy and Mineral Development, Petroleum Exploration and Production Department (PEPD), National Environment Management Authority (NEMA), Kampala
- Molleman F (2012). Butterflies of Uganda: Kibale Forest, a Picture Guide to Uganda Forest Butterflies. Treasures of Uganda Series, Kampala.
- Monadjem A, Cotterill FPD, Schoeman-Malan MC and Corrie Schoeman, M (2010). Bats of Southern and Central Africa: A Biogeographic and Taxonomic Synthesis, Wits University Press.
- Morrison KL and Bothma JduP (1998). Wetland nesting sites of wattled and grey crowned cranes on the Steenkampsberg Plateau, South Africa. South African Journal of Wildlife Research 28(4): 110-120.
- Mugerwa F (2013). Encroachers invade Bugoma forest again. Daily Monitor. <http://www.monitor.co.ug/News/National/Encroachers-invade-Bugoma-forest-again/-/688334/2118976/-/tuqsyqz/-/index.html>. Retrieved 29 January 2014
- Mwebaza-Ndawula L, Kiggundu V and Pabire GW (2001). Invertebrate communities of Nabugabo lakes: A vital support resource for the fisheries and ecosystem diversity. In: Busuulwa H, Mafabi PG and Mwebaza-Ndawula L (eds.). A Compilation of Scientific Information on Nabugabo Ramsar Site, Uganda. Nabugabo.
- Mwebaza-Ndawula L, Kiggundu V and Pabire GW (2003). Diversity and abundance of invertebrates in the Victoria basin lakes and their role in fishery production. Uganda Journal of Agricultural Sciences, 8: 209-220.
- Mwebaza-Ndawula L, Kiggundu V and Pabire GW (2004). The status and significance of invertebrate communities. In: Balirwa JS, Mugudde R and Ogutu-Ohwayo R (eds.) Challenges for Management of the Fisheries Resources, Biodiversity and Environment of Lake Victoria. First ed. Jinja, Uganda, Fisheries Resources Research Institute.
- National Environment Management Authority (1996). State of the environment Report for Uganda. Kampala: National Environmental Management Authority.
- National Environment Management Authority (2002). National Biodiversity Strategy and Action Plan, First Edition, Kampala.
- National Environment Management Authority (2010). Environmental Sensitivity Atlas for Albertine Graben, Second Edition, Kampala.
- NatureUganda (2010). Important Bird Areas in Uganda, training manual for Important Bird Areas Site Monitors. The East Africa Natural History Society, Kampala.
- Ndang'ang'a PK and Sande E (Compilers) (2008). International Single Species Action Plan for the Madagascar Pond-heron (*Ardeola idae*). CMS Technical Series No. 20, AEWA Technical Series No. 39. Bonn, Germany.
- Nicholson E, Westphal MI, Frank K, Rochester WA, Pressey RL, Lindenmayer DB and Possingham HP (2006). A New Method for Conservation Planning for the Persistence of Multiple Species. Ecology Letters 9:1049-1060.
- Nummi P and Kuuluvainen T (2013). Forest disturbance by an ecosystem engineer: beaver in boreal forest landscapes. Boreal Environmental Research 18(suppl. A): 13-24.
- Oates JF, Tutin CEG, Humle T, Wilson ML, Baillie JEM, Balmforth Z, Blom A, Boesch C, Cox D, Davenport T, Dunn A, Dupain J, Duvall C, Ellis CM, Farmer KH, Gatti S, Greengrass E, Hart J, Herbinger I, Hicks C, Hunt KD, Kamenya S, Maisels F, Mitani JC, Moore J, Morgan BJ, Morgan DB, Nakamura M, Nixon S, Plumtre AJ, Reynolds V, Stokes EJ and Walsh PD (2008). Pan troglodytes. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 January 2018.





- Ochieng H (2006). Individual mouthpart deformities of macrozoobenthos and community indices as indicators of pollution stress in northern Lake Victoria (Uganda). MSc. Thesis, UNESCO-IHE, Delft, 91pp.
- Ochieng H, de Ruyter van Steveninck ED and Wanda FM (2008). Mouthpart deformities in Chironomidae (Diptera) as indicators of heavy metal pollution in northern Lake Victoria, Uganda. *African Journal of Aquatic Science*, 33:135-142.
- Odum EP (1985). Trends expected in stressed ecosystems. *Bioscience* 35:419-422.
- Oghenekaro NO (2011). Application of macroinvertebrate-based biomonitoring approaches to assess anthropogenic impacts in the Swartkops River, South Africa. Master's thesis, Rhodes University.
- Ogutu-Ohwayo R, Mugidde R and Balirwa JS (2002). Technical Guidelines for Management of Fisheries Resources, Biodiversity and Environment of Victoria Basin Lakes. First Edition. Fisheries Resources Research Institute. 70pp.
- Okello W and Kurmayer R (2011). Seasonal development of cyanobacteria and microcystin production in Ugandan freshwater lakes. *Lakes & Reservoirs: Research and Management* 16: 123–135.
- Okello W, Portmann C, Erhard M, Gademann K. and Kurmayer R (2010). Occurrence of microcystin-producing cyanobacteria in Ugandan freshwater habitats. *Environmental Toxicology*. 25: 367-380.
- Økland RH (1990). Vegetation ecology: theory, methods and applications with reference to Fennoscandia. *Sommerfeltia* supplement, Botanical Garden and Museum, University of Oslo.
- Olson DM and Dinerstein E (1998) The global 200: a representation approach to conserving the earth's most biologically valuable ecoregions. *Conservation Biology* 12, 502-515.
- Organisation for Economic Co-Operation and Development (1982). *Eutrophication of Waters: Monitoring, Assessment and Control*. 154 pp., Paris.
- Pain D, Fishpool L, Byaruhanga A, Arinaitwe J and Balmford A (2005). Biodiversity representation in Uganda's forest IBAs. *Biological Conservation*. 125(1): 133-138.
- Parks D, Shaffer A, Barry D (2013). Nearshore drift-cell sediment processes and ecological function for forage fish: implications for ecological restoration of impaired Pacific Northwest marine ecosystems. *Journal of Coastal Research* 29(4):984-997.
- Parren MPE and Byler D (2003). Logging in West Africa: Impacts on Chimpanzees. In: Kormos R, Boesch C, Bakarr MI and Butynski TM. *West African Chimpanzees. Status Survey and Conservation Action Plan* IUCN/SSC Primate Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK. ix + 219 pp.
- Pearson DL (1994). Selecting indicator taxa for the quantitative assessment of biodiversity. *Philosophical Transactions Royal Society of London B*. 345:75–79.
- Pearson DL and Carroll SS (1998). Global patterns of species richness: spatial models for conservation planning using bioindicator and precipitation data. *Conservation Biology*. 12(4): 809-821.
- Pennak RW (1953). *Freshwater Invertebrates of the United States*. Ronald Press Co. ix, 769pp.
- Petchey O and Gaston K (2006). Functional biodiversity: back to basics and looking forward. *Ecology Letters* 9(6): 741-758.
- Petzold A, Vargas- Ramirez M, Kehlmaier C, Vamberger M, Branch WR, Du Preez L, Hofmeyr MD, Meyer L, Schleicher A, Siroky P and Fritz U (2014). A revision of African helmeted terrapins (Testudines: Pelomedusidae: Pelomedusa), with descriptions of six new species. *Zootaxa* 3795(5):523-548. Picker M, Griffiths C and Weaving A (2004). *Field guide to insects of South Africa*. STRUIK, Cape Town. Pimentel D (2007). *Environmental and Economic Costs of Vertebrate Species Invasions into the United States*. In: *Managing Vertebrate Invasive Species: Proceedings of an International Symposium*. Witmer GW, Pitt WC and Fagerstone KA (eds.). USDA/APHIS/WS, National Wildlife Research Centre, Fort Collins, CO. Plam M,





Jorgensen A, Kristensen and Madsen H (2008). Sympatric *Biomphalaria* species (Gastropoda: Planorbidae) in Lake Albert, Uganda, show homoplasies in shell morphology. *African Zoology* 43(1):34- 44.

Plumptre AJ (2002). Extent and Status of the Forests in the Ugandan Albertine Rift. Wildlife Conservation Society. Plumptre AJ, Arnold M and Nkuutu D (2003). Conservation Action Plan for Uganda's Chimpanzees 2003- 2008. UWA.

Plumptre AJ, Behangana M, Ndomba E, Davenport T, Kahindo C, Kityo R, Ssegawa P, Eilu G, Nkuutu D and Owiunji I (2003). The Biodiversity of the Albertine Rift. Albertine Rift Technical Reports No. 3. Wildlife Conservation Society.

Plumptre AJ, Davenport T, Behangana M, Kityo R, Eilu G, Ssegawa P, Ewango C and Kahindo C (2005). Albertine Rift, in: Mittermeier RA, Gil PR, Hoffman M, Pilgrim J, Brooks T, Mittermeier TG, Lamoreux J, da Fonseca GAB, Seligmann PA and Ford H. Hotspots Revisited: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions. Conservation International.

Plumptre AJ, Davenport TRB, Behangana M, Kityo R, Eilu G, Ssegawa P, Ewango C, Meirte D, Kahindo C, Herremans M, Peterhans JK, Pilgrim JD, Wilson M, Languy M and Moyer D (2007). The biodiversity of the Albertine Rift. *Biological Conservation*. 134:178-194.

Plumptre A, Akwetaireho S, Hänni DC, Leal M, Mutungire N, Kyamanywa J, Tumuhamy D, Ayebale J and Isoke S (2010). Biodiversity Surveys of the Bugoma Forest Reserve, Smaller Central Forest Reserves, and Corridor Forests South of Bugoma. Wildlife Conservation Society.

Plumptre A, Akwetaireho S, Leal M, Mutungire N, Kyamanywa J, Tumuhamy D, Bamutura B, Ayebale J, and Mukundane R (2011). Biodiversity Surveys of the Corridor Forests East of Bugoma Forest Reserve up to Budongo Forest Reserve. Wildlife Conservation Society.

Plumptre AJ, Rose R, Nangendo G, Williamson EA, Didier K, Hart J, Mulindahabi F, Hicks C, Griffin B, gawa H, Nixon S, Pintea L, Vosper A, McClennan M, Amsini F, McNeilage A, Makana JR, Kanamori M, Hernandez A, Piel A, Stewart F, Moore J, Zamma K, Nakamura M, Kamenya S, Idani G, Sakamaki T, Yoshikawa M, Greer D, Tranquilli S, Beyers R, Furuichi T, Hashimoto C and Bennett E (2010). Eastern Chimpanzee (*Pan troglodytes schweinfurthii*): Status Survey and Conservation Action Plan 2010-2020. Gland, Switzerland: IUCN, 52 pp.

Pontoppidan M and Nachman G (2013). Spatial Amphibian Impact Assessment – a management tool for assessment of road effects on regional populations of Moor frogs (*Rana arvalis*). *Nature Conservation*. 5:29-52.

Poole J, Kahumbu P and Whyte I (2013). *Loxodonta africana* Savanna Elephant (African Bush Elephant) pp 181-194. In: Kingdon J, Happold D, Hoffman M, Butynski, T, Happold M and Kalina N (eds). *Mammals of Africa Volume I: Introductory Chapters and Afrotheria*; Bloomsbury Publishing, London.

Porensky LM and Young TP (2013). Edge-Effect interactions in fragmented and patchy landscapes. *Conservation Biology*. 27(3):509-519.

Porensky LM, Bucher SF, Veblen KE, Treydte AC and Young TP (2013). Megaherbivores and cattle alter edge effects around ecosystem hotspots in an African savanna. *Journal of Arid Environments*. 96:55-63.

Prevedello JA, Figueiredo MSL, Grelle CEV and Vieira MV (2013). Rethinking edge effects: the unaccounted role of geometric constraints. *Ecography*. 36(3):287-299

Rabanal LI, Kuehl HS, Mundry R, Robbins MM and Boesch C (2010). Oil prospecting and its impact on large rainforest mammals in Loango National Park, Gabon. *Biological Conservation*. 143:1017-1024.

Radiwan S and Popiolek B (1989). Percentage of rotifers in spring zooplankton in lakes of different trophy. *Hydrobiologia*, 186/187: 235-238.

Ramsar Convention Secretariat (2013). The Ramsar Convention Manual: a guide to the Convention on Wetlands (Ramsar, Iran, 1971), 6th ed. Ramsar Convention Secretariat, Gland, Switzerland.





Ramsar Convention Secretariat (2014). The Ramsar Convention on Wetlands. <http://www.ramsar.org>, downloaded 26 March 2014.

Ray JC and Butynski TM (2013). *Profelis aurata* African Golden Cat pp 168-173. In: Kingdon J and Hoffmann M (eds). *Mammals of Africa Volume V: Pigs, Carnivores, Pangolins, Equids and Rhinoceroses*; Bloomsbury Publishing, London

Reed DH, O'Grady JJ, Brook BW, Ballou JD and Frankham R (2003). Estimates of minimum viable population sizes for vertebrates and factors influencing those estimates. *Biological Conservation* 113(1):23.

Roise JP, Shear TH and Bianco JV (2004). Sensitivity analysis of transportation corridor location in wetland areas: a multiobjective programming and GIS approach. *Wetlands Ecology and Management*. 12:519–529.

Rodriguez JP, Rodriguez-Clark KM, Baillie JEM, Ash N, Benson J, Boucher T, Brown C, Burgess ND, Collen B, Jennings M, Keith DA, Nicholson E, Revenga C, Reyers B, Rouget M, Smith T, Spalding M, Taber A, Walpole M, Zager I and Zamin T (2011). Establishing IUCN Red List criteria for threatened ecosystems. *Conservation Biology* 25:21-29.

Romaire JR (1985). Water quality, Crustaceans and Mollusk, Aquaculture in United States. In: Huner JV and Brown EE (eds). Westport: AVI Publishing.

Rompré G, Boucher Y, Bélanger L, Côté S and Robinson WD (2010). Conserving biodiversity in managed forest landscapes: the use of critical thresholds for habitat. *Forestry Chronicle* 86(5): 589-596.

Rotholz E and Mandelik Y (2013). Roadside habitats: effects on diversity and composition of plant, arthropod, and small mammal communities. *Biodiversity & Conservation*. 22(4):1017-1031.

RPS Group Plc (2006). Environmental and Social Impact Assessment for the KF1 well. Heritage Oil Plc.

Ruggiero LF, Hayward GD, and Squires JR (1994). Viability Analysis in Biological Evaluations: Concepts of Population Viability Analysis, Biological Population, and Ecological Scale. *Conservation Biology* 8:364-372.

Rutner-Kolisko A (1974). Planktonic rotifers: Biology and taxonomy. Biological Station Lunz of the Austrian Academy of Science. E. Schweizerbart'scheVerlagsbuchhandlung.

Samways MJ, McGeoch MA and New TR (2010). *Insect Conservation: A Handbook of Approaches and Methods*. Oxford University Press, Oxford.

Sande E, Dranzoa C, Wegge P and Carroll JP (2009a). Breeding requirements of Nahan's Francolin, *Francolinus nahan*, in Budongo forest reserve, Uganda. *African Journal of Ecology* 48:615–620.

Sande E, Dranzoa C, Wegge P and Carroll JP (2009b). Home ranges and survival of Nahan's Francolin *Francolinus nahan* in Budongo Forest, Uganda. *African Journal of Ecology* 47: 457–462.

Sarkar S and Illoldi-Rangel P (2010). Systematic Conservation Planning: an Updated Protocol. *Natureza & Conservação* 8(1):19-26.

Sars GO (1895). An account of the Crustacea of Norway, Christiania and Copenhagen Alb. Cammermeyer Forlag.

Savić A, Randelović V, Branković S and Krpo-Četković J (2011). Mayfly (Insecta: Ephemeroptera) community structure as an indicator of the ecological status of the Nisava river (Central Balkan Peninsula). *Aquatic. Ecosystem Health & Management* 14(3):276-284.

Schlitter D (2008a). *Miniopterus inflatus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.

Schlitter D (2008b). *Glauconycteris humeralis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017



Schlitter D, Hutson AM, Racey PA and Ravino J (2008). *Miniopterus fraterculus*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.

Secretariat of the Convention on Biological Diversity, Netherlands Commission for Environmental Assessment (2006). Biodiversity in Impact Assessment, Background Document to CBD Decision VIII/28: Voluntary Guidelines on Biodiversity-Inclusive Impact Assessment, Montreal, Canada.

Sergio F, Newton I, Marchesi L and Pedrine P (2006). Ecologically justified charisma: preservation of top predators delivers biodiversity conservation. *Journal of Applied Ecology*. 43 (6): 1049-1055.

Simberloff D (1997). Eradication. In: *Strangers in Paradise*. Simberloff D, Schmitz DC and Brown TC (eds.). Island Press, Washington, DC. USA. Pp. 221-228

Smith MD, van Wilgen BW, Burns CE, Govender N, Potgieter ALF, Andelman S, Biggs HC, Botha J and Trollope WSW (2013). Long-term effects of fire frequency and season on herbaceous vegetation in savannas of the Kruger National Park, South Africa. *Journal of Plant Ecology*. 6(1):71-83

Soulé ME, Estes JA, Berger J and Del Rio CM (2003). Ecological effectiveness: conservation goals for interactive species. *Conservation Biology* 17(5): 1238-1251.

Spawls S, Howell KM, Drewes R and Ashe J (2004). *A Field Guide to the Reptiles of East Africa*. Bloomsbury Publishing Plc.

Stattersfield AJ, Crosby MJ, Long AJ and Wege DC (1998). *Endemic Bird Areas of The World*. BirdLife Conservation Series No. 7, BirdLife International, Cambridge.

Stevenson T and Fanshawe J (2002). *Birds of East Africa*. Helm Field Guides, Christopher Helm, London.

Stork NE (1988). Insect diversity: Facts, fiction and speculation. *Biological journal of the Linnean Society* 35: 321-337.

Strasser EH, Heath JA and Thompson D (2013). Reproductive failure of a human-tolerant species, the American kestrel, is associated with stress and human disturbance. *Journal of Applied Ecology*. 50(4):912-919.

Stuart C and Stuart T (2013). *Dologale dybowskii* Pousargues's Mongoose (Savanna Mongoose); pp 364-365. In: Kingdon J and Hoffmann M (eds), *The Mammals of Africa*. Volume V: Carnivores, Pangolins, Equids, Rhinoceroses, Bloomsbury Publishing, London.

Stuart C, Stuart T, and Hoffmann M (2008). *Dologale dybowskii*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.

Sutherland WJ, Newton I and Green RE (eds) (2004). *Bird Ecology and Conservation: a handbook of techniques*. Oxford University Press, Oxford.

Swift TL and Hannon SJ (2010). Critical thresholds associated with habitat loss: a review of the concepts, evidence, and applications. *Biological Reviews* 85, 35–53.

Taabu-Munyaho A, Mbabazi D, Nakiyende H, Bassa S, Muhumuza E, Nsega M, Byekwaso A, Amiina R and Balirwa JS (2012). Report of the Catch Assessment Survey of Lake Albert – Albert Nile Conducted in July 2012. ATAAS Project 5: Monitoring changes in stocks of the major commercial fish species in the five major waterbodies of Uganda (Victoria, Albert, Kyoga, Edward, George), National Fisheries Resource Research Institute.

Talbot MR, Jensen NB, Lædal T and Filippi ML (2006). Geochemical responses to a major transgression in giant African lakes. *Journal of Paleolimnology* 35(3):467-489.

Talling JF (1963). Origin of stratification in an African Rift lake. *Limnology and Oceanography*, 8: 68-78.

Tasevska O, Kostoski G and Guseska D (2010). Rotifer-Based Assessment of the Lake Dojran Water Quality. Balwois-Ohrid.



- Terborgh J (1992). Maintenance of diversity in tropical forests. *Biotropica* 24:283-292.
- The Energy and Biodiversity Initiative (2006). Integrating Biodiversity into Environmental and Social Impact Assessment Processes; [www.theebi.org](http://www.theebi.org).
- Thomas CD, Jordana D, Lewis OT, Hill JK, Sutcliffe OL and Thomas JA (1998). Butterfly distributional patterns, processes and conservation. (In) Mace GM, Balmford A and Ginsberg JR (eds.), *Conservation in a Changing World*. Cambridge University Press. pp 107–138.
- Thomas D and Henry M (2013). *Eidolon helvum* African Strw-coloured Fruit-bat pp 232-234. In: Happold M and Happold DCD (eds). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.
- Thompson ME and Wrangham RW (2013). Pan troglodytes Robust Chimpanzee (Common Chimpanzee) pp 55-64. In: Buttyński TM, Kingdon J and Kalina J (eds). *Mammals of Africa Volume II: Primates*; Bloomsbury Publishing, London
- Tortoise and Freshwater Turtle Specialist Group (1996). *Kinixys erosa*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 12 December 2017.
- Treweek J (ed) (1999). *Ecological Impact Assessment*. Blackwell Scientific, London.
- Tucker G (2005). Biodiversity Evaluation Methods. In: Hill D, Fasham, M, Tucker, G, Shewry, M and Shaw, P (2005). *Handbook of Biodiversity Methods*. Cambridge University Press.
- Tushabe H, Kalema J, Byarungha A, Asasira J, Ssegawa P, Balmford A, Davenport T, Fjelds J, Friis I, Pain D, Pomeroy D, Williams P and Williams C (2006). A nationwide assessment of the biodiversity value of Uganda's Important Bird Areas network. *Conservation Biology*, 20(1): 85-99.
- United Nations Educational, Scientific and Cultural Organisation (2014). The World Heritage Centre, <http://whc.unesco.org/>, downloaded 26 March 2014.
- United Nations Environment Programme (2014). The Species+ Website. Nairobi, Kenya. Compiled by UNEPWCMC, Cambridge, UK. Available at: [www.speciesplus.net](http://www.speciesplus.net). Accessed 11 December 2017.
- United Nations High Commission for Refugees (2014). Country Operations Profile – Uganda. <http://www.unhcr.org/cgi-bin/texis/vtx/page?page=49e483c06&submit=GO>
- Utermöhl H (1958). Zur Vervollkommnung der quantitativen Phytoplankton methodik. *Mitteilungen des Internationalen Verein Limnologie*. 2:1-38.
- Uzarski DG, Burton TM, Kolar RE, Cooper MJ (2009). The ecological impacts of fragmentation and vegetation removal in Lake Huron's coastal wetlands. *Aquatic Ecosystem Health & Management*. 12(1):45-62.
- van der Hoeven CA, de Boer WF and Prins HHT (2010). Roadside conditions as predictor for wildlife crossing probability in a Central African rainforest. *African Journal of Ecology*. 48(2):368-377.
- Vincent K, Mwebaza-Ndawula L, Makanga B and Nachuha S (2012). Variations in zooplankton community structure and water quality conditions in three habitat types in northern Lake Victoria. *Lakes & Reservoirs: Research & Management*, 17: 83-95.
- Vymazal J (2011). Plants used in constructed wetlands with horizontal subsurface flow: a review *Hydrobiologia*. 674(1): 133-156
- Walsh JC, Venter O, Watson JEM, Fuller RA, Blackburn TM, and Possingham HP (2012). Exotic species richness and native species endemism increase the impact of exotic species on islands. *Global Ecology & Biogeography*, 21(8): 841-850.





- Wandera SB (2000). The fisheries of the Ugandan sector of Lake Albert. A report on the current state of the fishery of the Ugandan sector of Lake Albert. Prepared for FAO sub-regional representative for Southern and Eastern Africa (SAFD), 29 pp.
- Wandera SB and Balirwa JS (2010). Fish species diversity and relative abundance in Lake Albert—Uganda. *Aquatic Ecosystem Health & Management*, 13(3):284–293.
- Waterman C, Pietersen D, Soewu D, Hywood L and Rankin P (2014). *Phataginus tricuspis*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Waterman C, Pietersen D, Hywood L, Rankin P and Soewu D (2014). *Smutsia gigantea*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- Wellcome TI, Kardynal KJ, Franken RJ and Gillies CS (2014). Off-road sampling reveals a different grassland bird community than roadside sampling: implications for survey design and estimates to guide conservation. *Avian Conservation & Ecology*. 9(1):1-13.
- Wenn CL (2008). Do freshwater macroinvertebrates reflect water quality improvements following the removal of point source pollution from Spennings Beck, West Yorkshire? *Earth & Environment*, 3:369-406.
- Wennergren EB, Whitaker MD (1976). Investment in roads and spontaneous colonization: additional evidence from Bolivia. *Land Economics*. 52(1):88-94.
- Wessels KJ, van Jaarsveld AS, Grimbeek JD and van der Linde MJ (1998). An evaluation of the gradsect biological survey method. *Biodiversity and Conservation* 7:1093-1121.
- Wetzel RG (2001). *Limnology of Lake and River Ecosystems*. Third Edition. Elsevier Science, California, USA. Wetzel RG and Likens GE (2000). *Limnology Analyses*. Third Edition. Springer-Verlag, New York, USA.
- White F (1983). *The vegetation of Africa: a descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa*. Natural Resources Research XX. UNESCO, Paris. 356 pp.
- Wigley BJ, Bond WJ and Hoffman MT (2009). Bush encroachment under three contrasting land-use practices in a mesic South African savanna. *African Journal of Ecology*. 47:62-70.
- Wittemyer G, Elsen P, Bean W, Burton C and Brashares J (2008). Accelerated human population growth at protected area edges. *Science*. 321:123–126.
- Wittemyer G, Northrup JM, Blanc J, Douglas-Hamilton I, Omondi P and Burnham KP (2014). Illegal killing for ivory drives global decline in African elephants. *Proceedings of the National Academy of Sciences* doi: 10.1073/pnas.1403984111
- Wilson EO (1985). The biological diversity crisis. *Bioscience* 35:700-706.
- Wilson DE, Cole FR, Nichils JD, Rudran R and Foster MS (1996). *Measuring and Monitoring Biological Diversity: Standard Methods for Mammals*. Smithsonian Institution Press, Washington D.C.
- Wilson ML, Balmforth Z, Cox D, Davenport T, Hart J, Hicks C, Hunt KD, Kamenya S, Mitani JC, Moore J, Nakamura M, Nixon S, Plumtre AJ and Reynolds V (2008). *Pan troglodytes ssp. schweinfurthii*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 22 December 2017.
- With KA and Crist TO (1995). Critical thresholds in species responses to landscape structure. *Ecology* 76:2446-2459.
- Woodley S, Kay J and Francis G (eds) (1993). *Ecological Integrity and the Management of Ecosystems*. St. Lucie Press.
- World Conservation Monitoring Centre (1998). *Milicia excelsa*. The IUCN Red List of Threatened Species. Version 2017-3. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 December 2017.





World Database on Protected Areas (2014). <http://www.protectedplanet.net>.

World Health Organisation (2014). Ebola in Uganda. [http://www.who.int/features/2012/ebola\\_uganda/en/](http://www.who.int/features/2012/ebola_uganda/en/)

Yalden DW and Happold M (2013). *Otomops martiensseni* Large-eared Giant Mastiff Bat; pp 480-482. In: Happold M and Happold DCD (eds). *Mammals of Africa Volume IV: Hedgehogs, Shrews and Bats*; Bloomsbury Publishing, London.

Zommers Z and MacDonald DW (2012). Protected Areas as Frontiers for Human Migration. *Conservation Biology*. 26(3):547–556.

Zoological Society of London (2017). Evolutionarily Distinct and Globally Endangered. <http://www.edgeofexistence.org/>, accessed 1 December 2017





# APPENDIX A

## Document Limitations



# **APPENDIX B**

## **Critical Habitat Approach and Method of Assessment**



# **APPENDIX C**

## **Terrestrial Ecology Baseline Study**

# **APPENDIX D**

## **Aquatic Ecology Baseline Study**

# **APPENDIX E**

## **Land Cover Classification**

# APPENDIX F

## Impact Assessment Approach and Methods







# APPENDIX G

## Critical Habitat Screening and Appraisal





CNOOC UGANDA LIMITED

# Ecosystem Services Review and Assessment for the Kingfisher Field Development Area in Hoima & Kikuube Districts, Uganda

**Submitted to:**  
CNOOC Uganda Limited



**Report Number:** 13615730-13379-18

**Distribution:**

1 Electronic copy - Project 13615730



### Executive Summary

This report consists of an Ecosystem Services Review and Impact Assessment for the proposed Kingfisher Field Development Area (KFDA) and Pipeline route to Kabaale, which CNOOC intends to develop (the Project). The objective of this study is to identify priority ecosystem services and goods currently supplied in the Project Area of Influence; qualify the relationship between ecosystem services, the ecosystems that provide them, and the condition of those systems and the current drivers of change of those systems; identify the beneficiaries who depend on priority ecosystem services; identify Project impacts on priority ecosystem services; and recommend appropriate mitigation measures.

The ecosystem services **Local Study Area** (LSA) for this assessment (Figure 4) generally aligns with the local study areas used for the socio-economic baseline assessment, which consisted of the Kingfisher Field Development Area study area (comprising 11 villages in the Buhuka Parish and villages on top of the escarpment), and the pipeline route study area (comprising 22 villages in the vicinity of the pipeline route).

Land cover mapping of the Local Study Area was carried out. A review of the biodiversity, socioeconomic and physical data and information gathered during the baseline phase of the ESIA was used to identify the specific ecosystem services associated with each of the mapped land cover types and determine the condition of the land cover types. This allowed a judgement on the condition of the land cover types to be made, and, therefore, their potential capacity to supply ecosystem services, based on the reported baseline condition of the habitat types supported therein. Ecosystem services provided by the various ecosystems within the Local Study Area were then listed and described. Priority ecosystem services upon which an impact assessment was conducted, were derived from this list of relevant ecosystem services. Priority ecosystem services are:

- Services for which Project impacts could affect beneficiaries' livelihoods, health, safety or culture (Type I); and
- Services that could prevent the Project from achieving operational performance (i.e., impact the Project) (Type II).

Beneficiaries for ecosystem services within the Local Study Area were defined as the Project, the inhabitants and herders of the Buhuka Flats, and subsistence farmers whose plots lie within the servitude of the proposed pipeline route. Priority ecosystem services supplied within the Local Study Area include:

- |   |  |
|---|--|
| ■ Grazing for livestock                                 | ■ Fresh Water (Type I)                   |
| ■ Capture fisheries                                     | ■ Fresh Water (Type II)                  |
| ■ Wild Foods  | ■ Regulating Air Quality (Type II)       |
| ■ Construction materials for traditionally-built houses | ■ Regulating Water Flows and Timing      |
| ■ Natural aggregates for Project facility construction  | ■ Water Purification and Waste Treatment |
| ■ Biomass Fuel  | ■ Ethical and spiritual values           |
|   | ■ Educational and inspirational values   |

The key Project activities affecting beneficiaries will be:

- Changes in land cover and associated reductions in the supply or quality of ecosystem services due to the proposed construction of the KFDA facilities, the escarpment road, the oil export pipeline to Kabaale, and all associated infrastructure.
- Population influx of people seeking jobs during construction and operation of the Kingfisher Field Development Area; people seeking to provide commercial services to the increasing population in the





## ECOSYSTEM SERVICES ASSESSMENT

vicinity of the Project, and the concurrent increase in demand for ecosystem services. This is likely to impact the quantity and quality of ecosystem service supply to existing beneficiaries.

- Water abstraction from Lake Albert to provide make-up water for the oil extraction process is proposed as part of the Project; this has the potential to affect water quantity and quality in Lake Albert and thus may affect the fisheries potential of those areas, affecting beneficiaries that rely on fishing in these areas for livelihoods.
- As a result of these Project activities, moderate to major impacts are predicted on priority **provisioning ecosystem services**, moderate impacts are predicted on priority **regulating ecosystem services**, and major impacts are predicted on priority **cultural ecosystem services**, within the Project Area of Influence.

Application of recommended mitigation measures is expected to reduce the severity of predicted impacts. Mitigation measures include:

- Resettlement Action Plan
- Livelihood Restoration Plan
- Community Development Plan
- Influx Management Plan
- Corporate social responsibility initiatives
- Support of scientific studies and monitoring programs
- Worker and community education programmes
- Basin-wide water management initiatives
- Avoidance of sites of cultural heritage importance

Pre- and post-mitigation impacts on priority ecosystem services are summarised in the table below.

Priority Ecosystem Service	Severity Pre-mitigation	Severity Post-mitigation
Grazing for Livestock	Major – 16	Moderate – 6
Capture Fisheries	Major – 16	Moderate – 8
Wild Foods	Moderate – 9	Moderate – 6
Construction materials for traditionally-built houses	Major – 12	Moderate – 6
Natural aggregates for Project facility construction	Moderate – 9	Minor – 2
Biomass Fuel	Major – 16	Minor – 4
Fresh Water (Type I)	Major – 16	Moderate – 6
Fresh Water (Type II)	Major – 12	Minor – 4
Regulating Air Quality (Type II)	Moderate – 8	Minor – 2
Regulating Water Flows and Timing	Moderate – 9	Moderate – 6
Water Purification and Waste Treatment	Major – 12	Moderate – 6
Ethical and spiritual values, and Inspirational values (Type I)	Major – 12	Moderate – 8
Ethical and spiritual values, and Inspirational values (Type II)	Major – 12	Moderate – 8

Potential impacts on the Project's social license to operate may occur due to negative public perception of the Project's *perceived* effect on air quality and fish stocks, despite lack of evidence to corroborate these





## ECOSYSTEM SERVICES ASSESSMENT

---

perceptions. Worker and community education programmes, as detailed in the mitigation measures, are deemed to be critical for the Project to maintain its social licence to operate over the lifetime of the Project.







## Table of Contents

<b>1.0 INTRODUCTION</b>	<b>1</b>
1.1 The Concept of Ecosystem Services	1
1.2 Ecosystem Services and the International Finance Corporation	1
1.3 Regional Ecosystem Services in the Context of the Project	2
1.4 Report Structure	2
<b>2.0 TERMS OF REFERENCE</b>	<b>3</b>
2.1 Objectives	3
2.2 Scope	3
2.3 Limitations	3
<b>3.0 LEGISLATIVE AND POLICY CONTEXT</b>	<b>4</b>
3.1 Ugandan Legislation and Policy relating to Ecosystem Services	4
3.1.2 Uganda Wildlife Act (1996)	5
3.1.3 Uganda Wildlife Policy (1999)	5
3.1.4 Uganda National Biodiversity Strategy and Action Plan (2002)	5
3.1.5 Uganda Oil Policy	6
3.2 IFC Performance Standards 2012	6
3.3 International Conventions	7
<b>4.0 METHODOLOGY</b>	<b>9</b>
4.1 Local Study Area	10
4.2 Approach Overview	10
4.3 Identification of Ecosystem Services Relevant to the Project	10
4.3.1 Land Cover Mapping and Condition Assessment	10
4.3.1.1 Land Cover Condition Assessment	11
4.3.2 Desktop Review	12
4.3.3 Listing of relevant Ecosystem Services	12
4.4 Step 2: Prioritisation of Ecosystem Services	12
4.4.1 Stakeholder Engagement Approach	13
4.4.2 Ecosystem Service Prioritisation Exercise	13
4.5 Step 3: Delineation of the Project Area of Influence	14
4.6 Step 4: Establishment of the baseline for priority ecosystem services	18
4.7 Step 5: Assessing Project Impacts on Priority Ecosystem Services within the Project Area of Influence	18





4.7.1	Impact Assessment Process.....	18
4.7.2	Description of Potential Project Impacts.....	19
4.7.3	Assessing significance of Project impacts on affected Stakeholders .....	20
4.7.3.1	Determination of Intensity of Impact .....	20
4.7.3.2	Determination of Sensitivity of a Receptor.....	21
4.7.3.3	Determination of Impact Severity/Significance .....	21
<b>5.0</b>	<b>LAND COVER MAPPING AND CONDITION ASSESSMENT .....</b>	<b>22</b>
5.1.1	Land cover condition .....	24
<b>6.0</b>	<b>REVIEW OF BASELINE BIODIVERSITY, SOCIOECONOMIC AND BIOPHYSICAL ENVIRONMENT OF THE LOCAL STUDY AREA.....</b>	<b>24</b>
6.1	Terrestrial and Aquatic Biodiversity .....	24
6.1.1	Vegetation Communities Providing Ecosystem Services.....	24
6.1.1.1	Invasive plant species.....	29
6.1.2	Faunal Communities Providing Ecosystem Services .....	30
6.1.2.1	Freshwater Fish Communities of Lake Albert .....	30
6.1.2.2	Terrestrial Fauna Hunted for Bush Meat.....	30
6.2	Surface Water Systems .....	31
6.2.1	Surface Water Resources in Local Study Area .....	32
6.2.2	Ecosystem Services Provided by Surface Water Systems in Local Study Area .....	32
6.2.3	Project Water Demand.....	33
6.3	Land Use, Land Capability and Soils .....	37
6.4	Baseline Socio-economic Setting .....	38
6.4.1	Fisheries .....	38
6.4.2	Agriculture.....	39
6.4.2.1	Buhuka Flats.....	39
6.4.2.2	Pipeline Route .....	40
6.4.3	Freshwater .....	40
6.4.4	Land tenure and use .....	40
6.4.5	Human health risk .....	40
6.4.6	Recreation and Tourism.....	41
6.5	Cultural Heritage Context .....	41
6.5.1	Spiritual Values .....	41
6.5.1.1	Sacred Sites .....	41





6.5.1.2	Cultural landscapes .....	42
6.5.2	Inspirational Values associated with Lake Albert .....	42
6.5.3	Intangible cultural heritage .....	43
6.6	Existing Drivers of Ecosystem Change .....	44
<b>7.0</b>	<b>ECOSYSTEM SERVICE REVIEW FOR IMPACT ASSESSMENT .....</b>	<b>47</b>
7.1	Step 1: Identification of Relevant Ecosystem Services .....	47
7.1.1	Land Cover Types / Ecosystems which could be impacted by the Project.....	47
7.1.2	Which Ecosystem Services could the Project impact?.....	48
7.1.3	Which Beneficiaries are potentially affected?.....	50
7.2	Step 2: Ecosystem Service Prioritisation .....	51
7.2.1	Type I Ecosystem Services: Priority Ecosystem Services according to Project Impact .....	52
7.2.1.1	Provisioning .....	52
7.2.1.2	Regulating .....	53
7.2.1.3	Cultural .....	54
7.2.2	Type II Ecosystem Services – Priority Ecosystem Services according to the Extent of Project Dependence.....	55
7.2.2.1	Provisioning Services .....	55
7.2.2.2	Regulating Services.....	55
7.2.2.3	Cultural .....	56
<b>8.0</b>	<b>PROJECT AREA OF INFLUENCE FOR PRIORITY ECOSYSTEM SERVICES.....</b>	<b>57</b>
<b>9.0</b>	<b>ASSESSMENT OF PROJECT IMPACT ON PRIORITY ECOSYSTEM SERVICES .....</b>	<b>59</b>
9.1	Food Provisioning .....	59
9.1.1	Grazing for livestock.....	59
9.1.2	Capture Fisheries.....	60
9.1.3	Wild foods .....	60
9.2	Biological Raw Materials.....	61
9.3	Biomass Fuel.....	62
9.4	Freshwater.....	62
9.5	Regulating Air Quality .....	67
9.6	Regulating Water Flows and Timing .....	67
9.7	Water Purification and Waste Treatment .....	68
9.8	Ethical and Spiritual Values; Educational and Inspirational Values .....	70
<b>10.0</b>	<b>RECOMMENDED MITIGATION AND MONITORING MEASURES .....</b>	<b>72</b>





<b>11.0 CONCLUSION</b> .....	<b>80</b>
<b>12.0 REFERENCES</b> .....	<b>81</b>

## TABLES

Table 1: Ecosystems services categories .....	9
Table 2: Criteria for assessing ecosystem integrity / land cover condition .....	11
Table 3: Types of Ecosystem Service Impact .....	20
Table 4: Intensity assessment rating scale.....	21
Table 5: Sensitivity assessment rating scale.....	21
Table 6: Determination of impact severity .....	22
Table 7: Land cover in the Kingfisher Field Development Area and proportion lost as a result of proposed land take .....	23
Table 8: Land cover in the Pipeline area of influence and proportion lost as a result of proposed land take .....	23
Table 9: Land cover classes, associated ecosystems, habitat types and condition .....	24
Table 10: Mammals recorded in the KFDA LSA that are bush meat sources .....	31
Table 11: Land use within the Local Study Area .....	37
Table 12: Traditional Plant names and uses – Appendix A, .....	43
Table 13: ES supplied within the Project Area of Influence .....	48

## FIGURES

Figure 1: The flow of ecosystem services to beneficiaries .....	9
Figure 2: Locations of beneficiaries of ecosystem services that may be affected by the Kingfisher Field Development Area.....	15
Figure 3: Locations of beneficiaries of ecosystem services that may be affected by the Pipeline route.....	16
Figure 4: Local Study Area for Ecosystem Service Assessment.....	17
Figure 5: How assessment of Project impacts on ecosystems leads to assessing impacts on beneficiaries of ecosystem services (Landsberg et al., 2013) .....	19
Figure 6: Land cover in the Project Area of Influence .....	22
Figure 7: Vegetation communities of the Buhuka Flats .....	25
Figure 8: Traditionally built house in process of being thatched.....	26
Figure 9: Focus on vegetation communities and aquatic habitats of the Buhuka Flats .....	27
Figure 10: Vegetation communities of the Local Study Area.....	28
Figure 11: Vegetation communities of the Escarpment.....	30
Figure 12: Regional location of the Project in relation to surface water features .....	34
Figure 13: Regional Topography .....	35
Figure 14: Local context of the Project in relation to surface water features and floodlines .....	36
Figure 15: Backyard farming in Nsonga, Buhuka Flats .....	37





Figure 16: Drying Nile perch at the shore of Lake Albert.....	39
Figure 17: Overview of cultural sites and objects identified in the Local Study Area.....	44
Figure 18: "Chutes" used for the transport of thatching grass harvested from the escarpment.....	46
Figure 19: Examples of invasive species in the Project Area of Influence.....	47
Figure 20: Project Area of Influence for Priority Ecosystem Services (includes Lake Albert – see inset).....	58

### **APPENDICES**

#### **APPENDIX A**

Document Limitations

#### **APPENDIX B**

Prioritisation of ES according to Project Impact

#### **APPENDIX C**

Prioritisation of ES according to Project Demand



### DEFINITIONS

Term	Definition
<b>Beach Management Units</b>	Legally empowered community organisations for planning and management of fisheries resources in partnership with national and local governments, established by the East Africa Community Partner States (Kenya, Tanzania and Uganda).
<b>Grassland carrying capacity</b>	The maximum population size of cattle that can be sustained indefinitely, given the amount of grass available in a given area
<b>Critical Cultural Heritage</b>	Includes natural areas with cultural and/or spiritual value such as sacred groves, sacred bodies of water and waterways, sacred trees, and sacred rocks. Natural areas with cultural value are equivalent to priority ecosystem cultural services as defined in Performance Standard 6
<b>Cultural ecosystem service</b>	The nonmaterial contributions of ecosystems to human well-being, such as recreation, spiritual values, and aesthetic enjoyment.
<b>Customary land tenure</b>	Customary land tenure/holdings are plots which are held by individual households. Rights over a plot include the right to build a house, cultivate and to graze animals. These rights can only be abrogated if the occupier abandons the property or commits a taboo. This system has led to land fragmentation and associated economic consequences.
<b>Ecosystem Integrity</b>	The structure, composition, and function of an ecosystem operating within the bounds of natural or historic range of variation.
<b>Ecosystem Resilience</b>	The capacity of an ecosystem to respond to a disturbance by resisting damage and recovering quickly.
<b>Ecosystem Function</b>	Refers to all of the natural ecological processes that occur within an ecosystem, and is dependent on the composition and extent of a particular habitat or combination of habitats; their integrity or intactness, and their resilience.
<b>Irreplaceability</b>	Relates to rarity or uniqueness of an ecosystem in the landscape.
<b>Local Study Area</b>	The spatial context for the study
<b>Priority Ecosystem Services</b>	Those services on which project impacts affect the livelihoods, health, safety, or culture of the ecosystem service beneficiaries, and those services that could prevent the project from achieving planned operational performance.
<b>Project Area of Influence</b>	The area relevant to the assessment of project impacts and dependencies on priority ecosystem services; it includes the ecosystems that supply the priority ecosystem services, and the locations where the Project and affected stakeholders access priority ecosystem services.



## ECOSYSTEM SERVICES ASSESSMENT

Term	Definition
<b>Provisioning Ecosystem Services</b>	The goods or products obtained from ecosystems, such as food, timber, fibre, and freshwater
<b>Regulating Ecosystem Services</b>	The contributions to human well-being arising from an ecosystem's control of natural processes, such as climate regulation, disease control, erosion prevention, water flow regulation, and protection from natural hazards.
<b>Social Licence to Operate</b>	Social Licence to Operate (SLO) refers to the acceptance within local communities of both companies and their projects. In order to obtain an SLO it is necessary to develop good relationships with all stakeholders, especially with local communities
<b>Supporting Ecosystem Services</b>	The natural processes such as nutrient cycling and primary production, which maintain the other services.
<b>Spiritual ecosystem services</b>	Sacred, religious, or other forms of spiritual inspiration derived from ecosystems.
<b>Vulnerability</b>	Refers to degree of threat to an ecosystem or species.





### 1.0 INTRODUCTION

This report consists of an Ecosystem Services Review and Impact Assessment for the proposed Kingfisher Field Development Area (KFDA) and Pipeline route to Kabaale, which CNOOC intends to develop (the Project). The report describes the ecosystem services supplied by the various land cover types in the Project Area of Influence, and the benefits that the local community gains from them under existing conditions. The services that the Project itself will depend upon are identified, and existing drivers of ecosystem change discussed. The potential impacts of the Project on Priority<sup>1</sup> Ecosystem Services (Landsberg, et al., 2013) are assessed, and mitigation measures proposed for any adverse impacts on identified Priority Ecosystem Services.

#### 1.1 The Concept of Ecosystem Services

Ecosystem services consist of all the natural products and processes that contribute to human well-being, as well as the personal and social enjoyment derived from nature (Landsberg, et al., 2013). For example, wetlands provide grazing for livestock and act as nursery areas for juvenile fish at the edges of large open water systems. Wetlands often support populations of waterfowl, which can provide tourism and recreation opportunities for bird watchers; they may also help to mitigate climate change by sequestering carbon, and help reduce floods by storing rainwater (Macfarlane, et al., 2008).

Since different ecosystems provide different ecosystem services, there are trade-offs and synergies amongst ecosystem services - for example, conversion of forest to agriculture lowers the wood supply and potentially the water flow regulation, but it increases food production from crops. On the other hand, restoring a wetland may remove more pollutants from drinking water supplies and increase recreation benefits for bird watching (Landsberg, et al., 2013).

The benefits of ecosystems are passed on at many levels, and to many different beneficiaries. Examples of the benefits provided at different scales include:

- Local scale: ecosystem services may be the basis for rural livelihoods and subsistence; particularly for the poor; for example, artisanal fishing of inland lakes provides both cash income and food for low-income families.
- Regional scale: the provision of water to communities and businesses from a forested watershed.
- Global scale: ecosystems regulate climate and act as a reservoir of biodiversity that underpins biological production of all types, including agriculture.

#### 1.2 Ecosystem Services and the International Finance Corporation

The International Finance Corporation's (IFC) Performance Standard 6 - Biodiversity Conservation and Sustainable Management of Living Natural Resources (PS6) (IFC, 2012a), and its Guidance Notes (IFC, 2012b) - defines ecosystem services as **the benefits that people, including businesses, derive from ecosystems**. The IFC define two types of ecosystem services:

- Type I Ecosystem Services: Ecosystem Services on which the Project operations are most likely to have an impact and, therefore, which result in adverse impacts to affected communities (beneficiaries); and
- Type II Ecosystem Services: Ecosystem Services on which the Project is directly dependent for its operations, for example, water.

Although ecosystem services are largely addressed by IFC PS 6, the assessment of ecosystem services is spread throughout the environmental and social Performance Standards (PS) because the potential effects of a project on ecosystem services relates to all aspects of peoples' relationship with the environment, including health and safety risks, land ownership or usage, and cultural heritage. The specific PS that contain provisions for ecosystem services assessment are Performance Standard 1: Assessment and

<sup>1</sup> Priority ecosystem services are those where the significance of the project impact on the ecosystem services is considered likely to be high







Management of Environmental and Social Risks and Impacts; Performance Standard 3: Resource Efficiency and Pollution Prevention; Performance Standard 4: Community Health, Safety, and Security; Performance Standard 5: Land Acquisition and Involuntary Resettlement; Performance Standard 7: Indigenous Peoples and Performance Standard 8: Cultural Heritage. These are described in Section 2.0.

### 1.3 Regional Ecosystem Services in the Context of the Project

The Albertine Graben is recognised as one of Africa's most important areas for biodiversity; it is an area of high endemism and threatened species, with over 50% of birds, 39% of mammals, 19% of amphibians and 14% of reptiles and plants of mainland Africa occurring in this region (Plumptre, 2002). The Graben is recognised as an area of global importance for conservation, which is reflected by the high density of areas protected or designated for biodiversity. As such, this biodiversity represents one of Uganda's most vital economic resources, with the services and products provided by biodiversity in the form of ecosystems, species and genetic resources contributing billions of shillings per year to Uganda's economy, and support some of the poorest and most vulnerable sectors of Uganda's population (NEMA, 2002). In particular, rural people, landless people, and women in certain areas are dependent on biological resource utilisation as a primary means of subsistence and livelihood, or as a supplementary resource during times of drought, or unemployment (NEMA, 2002).

The Kingfisher Field Development Area and pipeline route from there to Kabaale is located in the Buhuka Parish and, Kyangwali Sub-County, in Kikuube and Hoima Districts of Western Uganda. Whilst subsistence farming and small-scale commercial farming are the main economic activities in the Hoima & Kikuube Districts, inhabitants of Buhuka Parish villages directly depend on subsistence fishing activities as a source of food, livelihoods and a cash income. The majority (approximately 98.9%) of the population in Hoima and Kikuube Districts use wood fuel and charcoal as the dominant source of energy, which is locally harvested; locally harvested natural resources are also used to provide building materials. Evidence suggests an increasing population in the Project area - the District Development Plan (Hoima District Local Government, 2011) indicates that the population of urban dwellers has increased from 31,671 in 2002, to 42,813 in 2009 and is projected to increase to 51,741 in 2013. In addition, there is observational evidence of expansion and urban growth in the neighboring towns, attributed to the oil and gas exploration activities. This urbanisation rate is expected to increase as oil and gas exploration activities proceed, which is anticipated to increase demand for ecosystem services in the Project area.

It is worth noting at this point that, although oil is a natural resource, it is not considered to be an ecosystem service. Although fossil fuels and some minerals come from organic material that was alive millions of years ago, their quantity and quality do not depend on the living component of *existing* ecosystems, and so are not considered to be benefits derived from existing ecosystems (Hanson, et al., 2012).

### 1.4 Report Structure

The report is structured sequentially:

- The terms of reference are introduced in Section 2.0 which provide the context for the study.
- Relevant international and national legislation and policy in terms of the Project's obligations to take ecosystem services into account are summarised in Section 3.0.
- The methods used in the determination of the Local Study Area, identification of ecosystem services and beneficiaries within the Local Study Area, prioritisation of ecosystem services and impact assessment of those priority ecosystem services are detailed in Section 4.0.
- Sections 5.0 and 6.0 provide the results of the Land Cover Classification exercise and literature review in order to identify ecosystem services and beneficiaries within the study area.
- Section 7.0 provides the process and results of the ecosystem service review for the local study area, and the prioritisation of ecosystem services exercise.
- The Project Area of Influence for priority ecosystem services is defined in Section 8.0.





- Section 9.0 describes the assessment of Project impact on **Priority ecosystem services** within the Project Area of Influence.
- Recommended mitigation and monitoring measures are outlined in Section 10.0.
- Study conclusions are drawn in Section 11.0.
- References for the study are given in Section 12.0.

## 2.0 TERMS OF REFERENCE

In determining the requirements of the Ecosystem Service Review and Impact Assessment for the Project, reference was made to the international guidance document 'Weaving Ecosystem Services into Impact Assessment' (Landsberg et al., 2013), appropriate Ugandan legislation and guidance, as well as international standards and guidance. National policy and international standards pertaining to the Project are detailed in Section 4.0.

The ecosystem services impact assessment concentrates on assessing predicted changes in ecosystems and ecosystem function, physical and aesthetic changes in the Lake Albert landscape, and changes in human population dynamics within the Project Area of Influence; and the concomitant effects that these changes will have on ecosystem service supply and demand within the same area.

### 2.1 Objectives

The aim of this Ecosystem Services Review and Impact Assessment is to:

- Identify priority ecosystem services and goods currently supplied in the context of the area in which the Project will be located.
- Qualify the relationship between ecosystem services, the ecosystems that provide them, and the condition of those systems, and the current drivers of change of those systems.
- Identify beneficiaries of the services, that is, the Project and/or the people who benefit from the goods and services supplied, and their level of dependence on the ecosystem services.
- Identify potential impacts on priority ecosystem services arising from the Project and propose mitigation measures.
- Identify any necessary additional areas of investigation.

### 2.2 Scope

In order to address the above objectives, and in line with the KFDA Scoping Report, a description and regional contextualisation of the baseline ecosystem services supplied and utilised within the Project Area of Influence was undertaken.

Using available regional data on ecosystem services in Western Uganda, and pertinent data gathered from the biodiversity, socioeconomic, surface water and cultural heritage baseline studies, an assessment of the predicted Project effects on the ecosystem services of the Project's area of influence was conducted to meet the requirements of IFC PS6.

### 2.3 Limitations

This assessment is a desk-based study, informed only by the data gathered as part of the biodiversity, cultural heritage and socioeconomic baseline studies. Although specific ecosystem service utilisation related questions were provided to the socioeconomic baseline team, a dedicated ecosystem services data gathering team did not go into the field to gather specific data.

Other study limitations include the following:

- No ecosystem service supply and demand modelling was done for this study.



- No assessment of Project dependence on priority ecosystem services over the life of the Project was done for this study, as no detailed Project design was available at the time of writing.
- Limitations encountered by the specialist studies that pertain to the ecosystem service-related information that could be derived from the baseline data include:
  - Access limitations due to health and safety considerations and site conditions.
  - The information gathered in relation to traditional cultural places and intangible heritage is limited to that which the community was willing to share with the field team. Some of the recorded sites are considered 'secret', and there may be places known only to a small section of the community and/or some which are too sensitive to share. Consequently, there is a potential for unidentified features of cultural importance to exist within the **Local Study Area**.
  - Transient populations and migratory population groups may not have been present during the period of socio-economic data collection. However, based on available research, the data collection period between November and December 2013 appears to be an optimal timeframe to encounter most of these transient population groups.

Despite these limitations to baseline data, the conclusions contained within this report are based upon a robust and transparent procedure, and represent an accurate evaluation and assessment of likely impacts.

### 3.0 LEGISLATIVE AND POLICY CONTEXT

#### 3.1 Ugandan Legislation and Policy relating to Ecosystem Services

Currently, there is no specific legislation directly pertaining to ecosystem service utilisation in Uganda; however, in the wake of increasing ecosystem degradation and biodiversity loss, Uganda has enacted several legislative and policy interventions to conserve natural resources and ensure that these resources provide sustainable benefit to the local community (CRA, 2006). Those considered most relevant to this ecosystem services assessment are described below.

##### 3.1.1 The Laws and Acts of Uganda

###### 3.1.1.1 *The Constitution of the Republic of Uganda (as at 15 February 2006)*

The over-arching government policy on natural resource conservation, cultural heritage preservation and social and economic wellbeing in Uganda is provided for in the Constitution of the Republic of Uganda. The relevant constitutional provisions in the National Objectives and Directive Principles of State Policy include the following:

- **Principles of State Policy XXVII (iv):** mandates the State (both central and local government) to create and develop parks, reserves and recreational areas, and to ensure conservation and promote the rational use of natural resources so as to safeguard and protect the biodiversity of Uganda.
- **Article 237 (2) (b):** the Government or local government, as determined by Parliament by law, shall hold in trust for the people and protect, natural lakes, rivers, wetlands, forest reserves, game reserves, national parks and any land, to be reserved for ecological and touristic purposes for the common good of all citizens.
- **Article 245:** the utilisation of natural resources of Uganda shall be undertaken in such a way as to meet the development and environmental needs of present and future generations of Ugandans and, in particular, the State shall take all possible measures to prevent or minimise damage and destruction to land, air and water resources resulting from pollution and other causes.
- **Social and Economic Objective (XIV):** Under the general social and economic objective, the State shall endeavour to fulfill the fundamental rights of all Ugandans to social justice and economic development and shall, in particular, ensure that all developmental efforts are directed at ensuring the maximum social and cultural well-being of the people.



- **Cultural Objective (XXIV):** Cultural and customary values, which are consistent with fundamental rights and freedoms, human dignity, democracy, and with the Constitution, may be developed and incorporated in aspects of Ugandan life. The State shall promote and preserve those cultural values and practices which enhance the dignity and well-being of Ugandans.
- **Cultural Objective (XXV):** Preservation of Public Property and Heritage: The State and citizens shall endeavour to preserve and protect, and generally promote, the culture of preservation of public property and Uganda's heritage.

### **Project Relevance**

The constitution of the Republic of Uganda obliges the state, Government and local governments of Uganda to conserve and protect natural and cultural heritage resources for the social and economic wellbeing of the people of Uganda. This has significance for the Project in terms of gaining the appropriate authorisations/licences/permits from Government authorities for the Project to proceed – the Government authorities will need to be satisfied that the obligations of the Constitution of the Republic of Uganda are fulfilled.

#### **3.1.2 Uganda Wildlife Act (1996)**

The Uganda Wildlife Act defines two types of conservation areas: “wildlife protected” and “wildlife managed” areas. Allowances for the utilisation of natural resources within particular protected areas are specified in the Act:

##### **Wildlife Protected Areas**

- **Wildlife Reserve:** these are protected areas of importance for wildlife conservation and management and in which conservation of biological diversity, scenic viewing, recreation, scientific research, and regulated extractive utilisation of natural resources are permitted.

##### **Wildlife Management Areas**

- **Community Wildlife Areas:** these are wildlife management areas where wildlife is protected, whilst taking into account the continued use of the land and the sustainable exploitation of wildlife in the area by people and communities ordinarily residing there. Sustainable exploitation of the natural resources of the area, including by mining and other methods, is permitted - providing that it is in a manner compatible with the continued presence of wildlife in the area.

### **Project Relevance**

Where the Project activities have the potential to affect wildlife reserves or community wildlife management areas, the appropriate permits must be sought. The Project will need to demonstrate that Project activities can be conducted in a sustainable manner that makes provisions to maintain the continued presence of wildlife within the area.

#### **3.1.3 Uganda Wildlife Policy (1999)**

The Uganda Wildlife Policy generally promotes long-term conservation of wildlife and biodiversity in a cost-effective manner, which maximises the benefits to the people of Uganda in terms of ecology, economy, aesthetics, science and education. The policy aims at achieving this through promoting conservation and sustainable utilisation of wildlife throughout Uganda.

### **Project Relevance**

The Policy relates to maintaining the sustainable use of wildlife; therefore the Project must take into account the provisions of the policy, to ensure that sustainable wildlife use is not compromised as a result of the direct and indirect effects of Project activities.

#### **3.1.4 Uganda National Biodiversity Strategy and Action Plan (2002)**

Uganda's National Biodiversity Strategy and Action Plan (NBSAP) (NEMA, 2002) was produced to comply with Uganda's commitments as a signatory to the Convention on Biological Diversity (CBD), which aims for





the conservation of biodiversity, its sustainable use, and sharing of the benefits of biodiversity. The goal of the NBSAP is “*To enhance biodiversity conservation, sustainable utilisation and equitable sharing of its benefits at all levels*”.

The NBSAP recognises that the services and products provided by biodiversity in form of ecosystems, species and genetic resources contribute billions of shillings per year to Uganda’s economy, through economic output in the in the fisheries, forestry, tourism, agriculture and energy sectors; and support of the poorest and most vulnerable sectors of Uganda’s population, who are highly dependent on biological resource utilisation. It emphasises the need to safeguard important ecosystem services, and the importance of social considerations in biodiversity conservation, and particularly in people’s ownership of, or participation in, biodiversity management. Chapter Three of the NBSAP describes guiding principles relating to community participation in the conservation and wise use of biological resources.

### **Project Relevance**

Uganda’s NBSAP commits the Government to develop national strategies, plans or programmes for the conservation and sustainable use of biological diversity, which directly relates to the use of and impacts on ecosystem services. The Project will need to demonstrate alignment with these instruments in order to satisfy Government obligations as a signatory to the CBD.

#### **3.1.5 Uganda Oil Policy**

The Oil and Gas Policy (2008) recognises that many areas with potential for petroleum production coincide with areas of important biodiversity, including national parks, water bodies, and forest reserves. It also acknowledges the risk of the oil and gas industry leading to significant in-migration of people looking for work, and seeks to enforce regulations restricting population movements and settlements in wildlife protected areas, with only a minimum of required infrastructure being allowed in such areas.

### **Project Relevance**

The Project will need to demonstrate alignment with the objectives of the Uganda Oil Policy, through wise application of the mitigation hierarchy in areas of biodiversity importance, and development of appropriate population influx management planning for the lifetime of the Project.

#### **3.1.6 The Uganda National Culture Policy, 2006**

Cultural Heritage is defined in paragraph 2.2 of the Uganda National Culture Policy as:

*“The cultural heritage of Uganda includes artistic and cultural expressions. These are; language and literary arts, performing arts, visual arts and handicrafts, indigenous knowledge, cultural beliefs, traditions and values, cultural sites monuments and antiquities”.*

The Uganda National Culture Policy provides the framework for the promotion of cultural heritage; it is all-inclusive and advocates the rights of indigenous groups in Uganda. The core principle underlying this policy is respect for all cultures. The Culture Policy promotes social change and encourages new ideas and approaches within the laws of Uganda.

### **Project Relevance**

The policy directly relates to the supply of cultural, spiritual and intangible services provided by the Lake Albert and escarpment landscape within the Project Area of Influence, and the associated traditions, folklore and ways of life of the people local to the area. The Project will need to demonstrate alignment with the policy to gain the necessary permits to proceed.

## **3.2 IFC Performance Standards 2012**

At the project financing level, the assessment and management of ecosystem services is largely dealt with in PS 6 - Biodiversity Conservation and Sustainable Management of Living Natural Resources (IFC, 2012a); however, elements of PS 1, 3, 4, 5, 7 and 8 are also relevant to ES assessment. Relevant parts of the PSs are briefly summarised as follows.





### ***PS 6 – Biodiversity Conservation and Sustainable Management of Living Natural Resources***

PS 6 directly relates to the four types of ecosystem services, as one of the three major objectives of PS 6 is to maintain the benefits of ecosystem services. It establishes objectives and requirements to avoid, minimise and, where residual impacts remain, compensate/offset for risks and impacts to ecosystem services within a project's area of influence. It puts an onus on project developers (the 'client') to carry out a systematic review (including participation of beneficiaries) of all ecosystem services a project will impact, or is dependent upon, to identify priority ecosystem services, and avoid, minimise, and mitigate impacts on priority ecosystem services for which a client has direct management control or significant influence.

### ***PS 1 – Assessment and Management of Environmental and Social Risks and Impacts***

This PS requires that all reasonably expected risks and impacts related to ecosystem services are identified, and broader definition of a project's area of influence be used. Indirect project impacts on ecosystem services upon which beneficiaries' livelihoods are dependent should be included in the assessment.

### ***PS 4 – Community Health, Safety and Security***

This PS establishes the requirement for the assessment of impacts on priority ecosystem services that may result in adverse health and safety risks to beneficiaries.

### ***PS 5 – Land Acquisition and Involuntary Resettlement***

PS5 relates to project situations where restrictions on land use, access to natural resources, and use of natural resources, such as aquatic resources, timber products and fresh water, impact affected beneficiaries of ecosystem services. The client must assess impacts on, and compensate for, loss of provisioning ecosystem services resulting from land acquisition and involuntary resettlement.

### ***PS 7 – Indigenous Peoples***

PS7 addresses impacts on lands and natural resources that may be subject to traditional ownership, or under customary use. Such use may be seasonal/cyclical, and may be ceremonial, cultural, or economic in nature. PS7 requires that adverse impacts on affected Communities of Indigenous Peoples should be avoided where possible; or otherwise be subject to appropriate application of the mitigation hierarchy to minimise adverse impacts.

### ***PS8 – Cultural Heritage***

PS8 deals with the protection of tangible and intangible Cultural Heritage, and sets out requirements for avoidance, or the application of an appropriate mitigation hierarchy to minimise adverse impacts. When replicable cultural heritage is removed and avoidance is not possible, restoration measures including the maintenance of ecosystem services required to support the cultural heritage must be taken, either in situ or in a different location. Non-replicable cultural heritage should not be removed unless several specific conditions are met. The Project should not remove or significantly alter or damage critical cultural heritage.

### ***Project Relevance***

In the case of its direct investments (including project and corporate finance provided through financial intermediaries), the IFC requires its clients to apply the Performance Standards to manage environmental and social risks and impacts so that development opportunities are enhanced. Together, the Performance Standards establish standards that the Project is to meet throughout the life of an investment by IFC. As stated above, Performance Standards 1, 4, 5, 6, 7 and 8 have components that directly relate to ecosystem services and maintenance of their supply despite project impact. Therefore, in order to secure Project funding from IFC, the Project must demonstrate that it is in compliance with the requirements of each of the above-mentioned performance standards.

## **3.3 International Conventions**

### **3.3.1 The Convention on Biological Diversity (1992)**

Under the convention, each contracting party is expected to develop national strategies, plans or programmes for the conservation and sustainable use of biological diversity. Uganda is a signatory to the



CBD, which aims for the conservation of biodiversity, its sustainable use, and sharing of the benefits of biodiversity. Uganda's commitments as a signatory to the CBD are provided for in the Uganda NBSAP (ref. Section 3.1.4).

### ***Project Relevance***

As a signatory to the CBD, Uganda's Government is committed to develop national strategies, plans or programmes for the conservation and sustainable use of biological diversity, implemented through a National Biodiversity Action Plan (ref. Section 3.1.4). The Project will need to demonstrate alignment with the provisions of the NBSAP in order to satisfy Government obligations as a signatory to the CBD.

### **3.3.2 The Convention for the Protection of the World's Cultural and Natural Heritage (1972)**

Uganda is a signatory to the Convention on the Protection of the World Cultural and Natural Heritage (1972). To date, Uganda has three sites on the list of the World heritage sites namely: Kasubi tombs, enlisted in 2001; Bwindi Impenetrable Forest National Park; and Ruwenzori Mountains National Park. In 2005, UNESCO proclaimed the art of barkcloth making in Uganda a masterpiece of the oral and intangible heritage of humanity. Currently, five sites are on the World Heritage nomination list, including the ancient salt making sites at Kibiro that lies within the Albertine Graben, approximately 45 km north east of the Kingfisher Field development area.

### ***Project Relevance***

The Project will need to demonstrate alignment with the provisions of the convention in order to satisfy Government obligations as a signatory to the convention, through identifying and protecting cultural heritage by ensuring that internationally recognised practices for the protection, field-based study, and documentation of cultural heritage are implemented.

### **3.3.3 The Convention for the Safeguarding of the Intangible Cultural Heritage (2003)**

Uganda has been a signatory to the United Nations Educational, Scientific and Cultural Organisation's (UNESCO's) Convention for the Safeguarding of Intangible Cultural Heritage since 2009. The Convention seeks to raise awareness of threats to intangible heritage and encourages member states in the identification, protection and management of such assets, ensuring respect for those individuals and communities concerned.

### ***Project Relevance***

The Project will need to demonstrate alignment with the provisions of the convention in order to satisfy Government obligations as a signatory to the convention, through identifying and protecting intangible cultural heritage and cultural practices by ensuring that internationally recognised practices for the protection, field-based study, and documentation of cultural heritage are implemented.



### 4.0 METHODOLOGY

Ecosystem services are the benefits that people and/or a project (the beneficiaries) obtain from ecosystems. In the strictest sense, without those beneficiaries, there are no ecosystem services. The benefits gained can be either physical or psychological, and can be obtained actively or passively, directly or indirectly. For the purposes of this assessment, the definitions of ecosystem services were based on those developed by the Millennium Ecosystem Assessment (MA, 2005), (Table 1). These definitions were chosen to keep consistency with the IFC's Performance Standards, and because they are widely recognised.

Ideally, the Project should maintain the value and functionality of priority ecosystem services to those beneficiaries directly dependent upon them, through direct management control. As such, ecosystem services whose beneficiaries are at the global scale, and to a lesser extent, the regional scale, are not covered by this assessment.

Table 1: Ecosystems services categories

Broad categories	Definition
Supporting services	Natural processes essential to resilience, and functioning of ecosystems. e.g., primary production
Regulating services	Control of the natural environment e.g., maintenance of key ecological processes, protected areas, habitat of special value, groundwater recharge, catchments
Provisioning services	Supporting human needs e.g., traditional hunting grounds, medicinal plants and minerals, water sources, fishing grounds, fire wood
Cultural services	Aesthetic, spiritual, recreational, and other cultural values. e.g., sacred sites, recreation, sense of place

As mentioned, without the beneficiaries (that is, the local community (Type I) and the Project (Type II)), there are no ecosystem services. In terms of a project's setting, that is, its location, an understanding of the ecosystem processes occurring in the area is important, as it enables an understanding of how those processes affect the supply and demand of the ecosystem services arising from such processes, and the value the ecosystem services eventually offer to beneficiaries (that is, the supply side). A conceptual ecosystem services flow path illustrating these supply linkages, using photosynthesis and the functions, services and benefits that flow from it as an example, is shown in Figure 1.



Figure 1: The flow of ecosystem services to beneficiaries

Given the above, and given that the assessment of ecosystem services is also concerned with the social aspects of the benefits of services (that is, the demand side), the assessment of ecosystem services relied upon data gathered during the stakeholder engagement processes carried out as part of the socio-economic baseline study and the cultural heritage baseline study. Other information for the ecosystem services assessment was gathered from the various specialist inputs to the baseline for the ESIA, during the Desktop Review (Section 3.3).







### 4.1 Local Study Area

An over-arching ecosystem services assessment for the entire Albertine Graben region is currently being completed by independent consultants. Therefore, the focus of this assessment is only on ecosystem services at the local scale to the Project, specifically, the Kingfisher Field Development Area where the main project footprint will be located, the proposed escarpment road, and the proposed pipeline route (Figure 2, Figure 3). This ecosystem services Local Study Area for this assessment (Figure 4) generally aligns with the local study areas used for the socio-economic baseline assessment, which consisted of the Kingfisher Field Development Area study area (comprising 11 villages in the Buhuka Parish and villages on top of the escarpment), and the pipeline route study area (comprising 22 villages in the vicinity of the pipeline route). Throughout the report, the various areas of the Local Study Area are discussed in terms of its unit components, that is, the Buhuka Flats, the escarpment, and the pipeline route.

### 4.2 Approach Overview

The approach taken to conducting the ecosystem services review is based on Steps 1 to 3 of the method put forward by Landsberg et al. (2013). Given that the current assessment is a desk-based study, the method was adapted to preclude additional baseline data gathering for priority ecosystem services once the initial list of relevant ecosystem services list was refined to focus on priority ecosystem services only. Instead, baseline data on all ecosystem services (priority and non-priority) was gathered during stakeholder engagement (Step 2) in order to determine how all ecosystem services currently contribute to stakeholders' livelihoods, health, safety or culture.

The approach to impact assessment consisted of a combination of the Project impact assessment on priority ecosystem services method in Landsberg et al. (2013), and the prescribed impact assessment method being used for the ESIA (ref. Section 4.7).

### 4.3 Identification of Ecosystem Services Relevant to the Project

The ecosystem services that that Project could impact were identified by first defining which ecosystems could be affected, determining the ecosystem services supplied by and demanded from those ecosystems, and identifying the beneficiaries who use those services supplied by the ecosystems that could be affected, as per Step 1 of the guidance provided in (Landsberg, et al., 2013).

This was done by means examination of the land cover mapping exercise done during the biodiversity baseline assessment to identify the land cover types and thereby ecosystems that could be affected by the Project, and a desk study to identify the ecosystem services supplied by those land cover types, and the beneficiaries that use them.

#### 4.3.1 Land Cover Mapping and Condition Assessment

A land-cover dataset was produced, using existing multi-band, high-resolution satellite imagery (Figure 6). The locations of Project infrastructure and activities were mapped against the land cover types that potentially supply ecosystem services, to identify those land cover types may be impacted by the proposed construction, operation and decommissioning of the Project.

- The land-cover dataset provided the following information:
- Land-cover types that would be affected by loss in area to the Project footprint.
- The area/extent (hectares) of loss of each land cover type identified. The size of each land cover type was measured from aerial photography via GIS, as was the area of each land cover type being lost to the Project footprint – allowing the proportion of loss within the different Project Area of Influence to be calculated.
- Identification and location of communities dependent on services supplied within the Local Study Area.
- Together with information from the existing biodiversity/social/physical baseline data (Section 4.2.2), the land cover dataset was used to identify ecosystems that could be impacted by the Project, and





subsequently the ecosystem services supplied by the potentially impacted ecosystems could be identified.

**4.3.1.1 Land Cover Condition Assessment**

Condition assessments are widely adopted as regulatory indicators of ecosystem function, and for some services (e.g., habitat) links between condition and function are often direct (McLaughlin & Cohen, 2013)

Ecosystem integrity, or the condition of the land cover types (and therefore their capacity to supply ecosystem services – ref. Figure 5) was therefore assessed, primarily based on the ecosystem integrity assessment of the various habitat types identified in the biodiversity baseline report. The integrity of ecosystems was determined based on the criteria put forward Table 2 (after: (Kent & Coker, 1992), (Treweek (ed), 1999), (Tucker, 2005), and (Secretariat of the Convention on Biological Diversity, 2006)).

**Table 2: Criteria for assessing ecosystem integrity / land cover condition**

Criterion	Description
Composition	Diversity and complexity - what is there and how abundant (in a particular time frame) it is
Structure (or pattern)	How biological units are organised in time and space. Ecosystem ‘scale’ refers to the space it occupies and the way it changes over time. The structure and interactions that shape the flow of energy and the distribution of biomass.
Linkages and corridors	To habitat of the same or different ecosystems, which provide an important ‘playing field’ for ecological processes and enable persistence. These linkages are in contrast to a highly fragmented landscape where patches of natural habitat are effectively isolated.
Key processes (including ecosystem function)	Which natural (that is, physical and/or biological) and/or human-induced processes are of key importance for the creation and/or maintenance of ecosystems. These are termed drivers of change, and include direct and indirect drivers. Examples of direct drivers include: changes in land use and land cover; fragmentation and isolation; extraction, harvest, or removal of species; external inputs such as emissions, effluents, chemicals; disturbance; introduction of invasive, alien and/or genetically modified species; and restoration. Examples of indirect drivers of change include: demographic; economic; socio-political; cultural; and technological processes or interventions.
Representativeness in the landscape	The uniqueness of the ecosystems within the LSA and the wider landscape; this rarity factor is related to the concepts of irreplaceability and vulnerability.
Resilience and stability	The ability of the ecosystem to absorb change, persist, and maintain the same form.

Based on the assessment of these criteria, the condition of the ecosystems and habitats was estimated and assigned a subjective class, as defined in the biodiversity baseline assessment:

- Pristine
- Near-pristine
- Slightly-degraded
- Moderately-degraded
- Heavily-degraded
- Note that not all ecosystem services supply dynamics are affected to the same degree by the condition of the land cover type; for example, harvest of wood for fuel takes place both in forest habitats in good condition, and in bushland that might be considered to be in a degraded condition due to overgrazing.





However, the bushland may be more important in terms of supply as a result of proximity to settlements and ease of access.

### 4.3.2 Desktop Review

A review of the biodiversity, socioeconomic and physical data and information gathered during the baseline phase of the ESIA was conducted, and a full literature review of available publications, reports and data relating to ecosystem services and biological resource use in the Albertine Graben and Western Uganda was done. Relevant information from the review was collated to identify the specific ecosystem services associated with each of the mapped land cover types and determine the condition of the land cover types, and thus their potential/capacity to supply the ecosystem services,

The socioeconomic baseline data enabled the identification of the people who depend on those potentially impacted ecosystem services for their livelihoods, health, safety, and culture, i.e. the beneficiaries. Beneficiaries are defined as those individuals, communities, institutions, and companies (including CNOOC) that could be positively or negatively affected as a result of Project impacts on ecosystem services (Landsberg, et al., 2013).

The baseline data for the Project footprint reviewed and presented in this report was sourced from the following specialist reports done for the baseline study for the Project ESIA:

- Cultural Heritage Baseline
- Socio-Economic Baseline
- Surface Water Baseline,
- Soils and Land Capability Baseline
- Biodiversity baseline report

Numerous reports relating to ecosystem service supply and natural resource utilisation in the Albertine Graben and Western Uganda were used to inform the study; these are referenced throughout the text.

### 4.3.3 Listing of relevant Ecosystem Services

- A comprehensive ecosystem services supply and demand list was then developed based on relevant information collated from the desktop review. This included: the land cover types and associated ecosystem services that could be directly impacted by the Project; the natural resources that the Project will require for its operation; information on water supply, energy use, economic activities and population movement and migration derived from the baseline socioeconomic study; and the cultural and spiritual traditions and beliefs of people in the Project area detailed in the baseline cultural heritage study.
- The ecosystem services used, and beneficiaries of those services, were described. No ranking of importance of the ecosystem services was done at this stage; instead, the list was carried through to the next step where ecosystem services were prioritised (Section 4.4).

## 4.4 Step 2: Prioritisation of Ecosystem Services

Priority ecosystem services, upon which the impact assessment was focused, were selected from the list of relevant ecosystem services generated in Step 1. Priority ecosystem services are:

- Services for which Project impacts could affect beneficiaries' livelihoods, health, safety or culture (Type I);
- Services that could prevent the Project from achieving operational performance (i.e. impact the Project) (Type II).



### 4.4.1 Stakeholder Engagement Approach

Collection of stakeholder input to the prioritisation of ecosystem services for which Project impacts could affect beneficiaries was undertaken via surveys of local residents within the area of influence, as part of the socioeconomic and cultural heritage baseline data gathering fieldwork. The interviews gathered information on Type I priority ecosystem services as defined by the IFC (2012a).

The information gathered during the interviews was used to:

- Identify what ecosystem services were being used by beneficiaries, and gain an understanding of how much of the particular services were used, how far people had to travel to obtain it, and the importance of that service to their livelihood, wellbeing, and culture.
- Identify existing drivers of ecosystem change.
- Assess current supply of priority ecosystem services, the degree of dependence that beneficiaries have on priority ecosystem services, and whether the services are readily substitutable, compensable, irreplaceable etc.
- Estimate foreseeable supply of ecosystem services and their contribution to beneficiaries' well-being in the absence of the project.

The interviews were carried out by the social and cultural heritage specialists in December 2013 to April 2014, during the baseline data gathering phase of the ESIA. All conversations were facilitated by CNOOC's Community Liaison Officer (CLO).

#### **Socio-economic baseline**

Primary data collection for the socio-economic baseline study was conducted in the Local Study Area during December, 2013 for the KFDA local study area, and March 2014 for the pipeline study. Data collection included focus group discussions, key stakeholder interviews, a sample household socio-economic survey and a land-use constraints mapping ground-truthing exercise.

#### **Cultural Heritage baseline**

A non-invasive field survey to record all cultural heritage sites within the Local Study Area was conducted between 20 January and 2 February 2014. During the field work, consultation with the affected communities (those villages within the respective discipline-specific LSAs) was undertaken. Consultation was done via transcribed interviews in order to capture places of local cultural and/or sacred importance (for example, ritual sites, burial grounds, churches and mosques) and any related intangible heritage practice (taboo, oral history, traditional plant and medicinal plant usage etc.).

### 4.4.2 Ecosystem Service Prioritisation Exercise

- The ecosystem service prioritisation exercise was carried out systematically, using the WRI Impact and Dependence Scoping tools, and current guidance regarding conducting an Ecosystem Services Review (Landsberg, et al., 2013). In addition, ecosystem services guidance specifically pertaining to the oil and gas industry was applied for the assessment of potential dependencies and impacts of oil and gas projects and operations (IPIECA, 2011).
- The list of priority ecosystem services supplied in the Project's area of influence was developed by identifying priority ecosystem services; that is, those where the significance of the Project's impact is considered high, specifically:
- Type I priority ecosystem services were identified and defined by:
  - Identifying potential Project-caused drivers of ecosystem change.
  - Identifying potentially impacted ecosystems and associated ecosystem services, and potentially affected beneficiaries.
  - Assessment of Project's impact on the ecosystem services.





- Type II priority ecosystem services were identified and defined based on the Project's requirements as outlined in the Project Description.
- The importance of each ES to communities, and their level of dependence upon the supply of the ecosystem services, was established via engagement with the socio-economic, cultural heritage, biodiversity and surface water specialists.
- Identification of the availability of alternatives for the supply of identified ecosystem services, including the existence of, and access to, those alternatives, based on land cover mapping, stakeholder interviews and information derived from the baseline studies conducted for the ESIA.

### 4.5 Step 3: Delineation of the Project Area of Influence

The **Project area of influence** is the area relevant to the assessment of project impacts and dependencies on priority ecosystem services; it includes the ecosystems that supply the priority ecosystem services, and the locations where the Project and affected stakeholders access priority ecosystem services (Landsberg, et al., 2013).

The Project area of influence was set by mapping the locations of Project infrastructure and activities against the land cover types that supply priority ecosystem services to identify those land cover types that may be impacted by the proposed construction, operation and decommissioning of the Project. The locations where the beneficiaries of the identified priority ecosystem services (Figure 2, Figure 3) access those services were then mapped and used to define the boundary of the Project Area of Influence for Impact Assessment.



# ECOSYSTEM SERVICES ASSESSMENT




Figure 2: Locations of beneficiaries of ecosystem services that may be affected by the Kingfisher Field Development Area





# ECOSYSTEM SERVICES ASSESSMENT




Figure 3: Locations of beneficiaries of ecosystem services that may be affected by the Pipeline route





# ECOSYSTEM SERVICES ASSESSMENT




Figure 4: Local Study Area for Ecosystem Service Assessment







### 4.6 Step 4: Establishment of the baseline for priority ecosystem services

As mentioned previously, Landsberg *et al.*'s (2013) ecosystem service review method was adapted to preclude additional baseline data gathering for priority ecosystem services once the initial ecosystem services list was refined (Step 4). Instead, baseline data on all ecosystem services (priority and non-priority) was gathered during stakeholder engagement (ref. Section 4.4.1) in order to determine how all ecosystem services currently contribute to stakeholders' livelihoods, health, safety or culture.

### 4.7 Step 5: Assessing Project Impacts on Priority Ecosystem Services within the Project Area of Influence

The impact assessment identifies the intensity of a particular impact from the Project and then compares that intensity with the sensitivity of the receiving environment to derive an overall severity for the impact. This method relies on a detailed description of both the impact and the ecosystem service valued component that is the receptor. The intensity of an impact depends on its characteristics, which includes factors such as its magnitude, duration, reversibility, area of extent, and nature in terms of whether positive, negative, direct, indirect or cumulative.

One of the main purposes of the impact assessment is to provide answers to questions that people have about how a project could affect something that matters to them, such as a valued component. To focus this assessment, and ensure that the impact assessment clearly addressed the key issues raised by the stakeholders (see Section 4.4.1), and the objectives set for this impact assessment (see Section 2.1), questions were formulated that captured the concerns relative to a particular issue. In this report, those concerns are expressed as a 'key question', which forms the basis of the investigations of potential effects and impacts of the Project:

#### 1) What impact could the Project have on the supply of priority ecosystem services to beneficiaries?

In order to answer the key question in relation to ecosystem services, the impact assessment involved the following steps:

- 1) Review and identification of the trends and external, non-project-related threats to current ecosystem services supply - current pressures on ecosystems and the resilience inherent in those ecosystems were identified.
- 2) Assessment of the replaceability/substitutability of ecosystem services, and/or spatial alternatives for confirmed vulnerable beneficiaries.
- 3) An impact significance assessment (ref. Section 4.7.3.3) was conducted on Type I and II priority ecosystem services before and after mitigation.
- 4) Identification of social/operational/financial/regulatory/reputational risks associated with the residual impacts.
- 5) Identification of potential alternatives to supply of services.

#### 4.7.1 Impact Assessment Process

The impact assessment process compares the intensity of the impact with the sensitivity of the receiving environment. This method relies on a detailed description of both the impact and the environmental or social component that is the receptor. The intensity of an impact depends on its characteristics, which may include such factors as its duration, reversibility, area of extent, and nature in terms of whether positive, negative, direct, indirect or cumulative.

The impact assessment process was aligned with the World Resources Institute (WRI) approach (Landsberg, et al., 2013), consisting of a combination of those workers' approach to assessment of Project impact on priority





ecosystem services and thereby assessment of impact on beneficiaries (Figure 5); and the prescribed impact assessment method being used for the ESIA (ref. Section 4.7.3).




Figure 5: How assessment of Project impacts on ecosystems leads to assessing impacts on beneficiaries of ecosystem services (Landsberg et al., 2013)

### 4.7.2 Description of Potential Project Impacts

Interactions between the proposed Project activities and priority ecosystem services were identified through a review of the current Project Description in the context of the identified baseline environment in the Local Study Area (Section 6.6.0). In summary, Project activities will change the physical landscape and socio-economic context of the Project Area of Influence, which will result in direct and indirect impacts to priority ecosystem services.

- The key Project impacts affecting beneficiaries will be:
- Changes in land cover and associated reductions in the supply of ecosystem services due to the proposed construction of the KFDA facilities, the escarpment road, the oil export pipeline to Kabaale, and all associated infrastructure:
  - The physical presence of the Project in the landscape will directly change the land surface and will potentially interact with cultural heritage features - these are 'direct impacts', which are likely to affect both beneficiaries within or adjacent to the Project footprint, and beneficiaries from further afield who may travel to avail of cultural heritage ecosystem services intrinsically linked with the Lake Albert and Escarpment landscape.
  - Activities that will not affect the land surface directly may indirectly alter the setting in which a site is experienced (for example, by related dust and noise disturbance) or limit the supply of provisioning ecosystem services (e.g., oil development areas may be fenced off which could restrict access to grazing lands for cattle) – these are 'indirect impacts'. These are likely to affect beneficiaries within close proximity to the development.
- Population influx of people seeking jobs during construction and operation of the Kingfisher Field Development Area, and people seeking to provide commercial services to the increasing population in the vicinity of the Project, and the concurrent increase in demand for ecosystem services; this is likely to impact the quantity and quality of ecosystem service supply to existing beneficiaries.
- Water abstraction from Lake Albert to provide make-up water for the oil extraction process is proposed as part of the Project; this has the potential to affect water quantity and quality in areas of Lake Albert and thus may affect the fisheries potential of those areas, affecting beneficiaries that rely on fishing in these areas for livelihoods.



The types of potential Project impacts considered appropriate for the ecosystem services assessment are summarised in Table 3.

Table 3: Types of Ecosystem Service Impact

<b>Direct Impact</b>	Impacts that result from a direct interaction between a planned Project activity and the receiving environment/receptors (e.g., destruction of a sacred site as a result of construction of well pads, loss of an ecosystem’s capacity to supply an ecosystem service due to degradation from over-grazing).
<b>Indirect impact</b>	Secondary impacts that result from project activity and affect the environment in which the receiving receptor is experienced (e.g., job-seeking population influx to the area and concurrent increase in demand for ecosystem services).
<b>Cumulative impact</b>	Impacts that act together with other impacts (including those from concurrent or planned activities from other projects) to affect the same resources and/or receptors as the Project.

4.7.3 Assessing significance of Project impacts on affected Stakeholders

4.7.3.1 Determination of Intensity of Impact

Magnitude or intensity describes the severity of the effect. To classify magnitude using an ordinal scale (that is, negligible, low, medium, or high) in a manner meaningful for ecosystem services, the effect size (loss of land cover class that supplies the ecosystem services to the Project footprint) must be placed in the context of the availability of the land cover class and thus available supply of the ecosystem services within the Project’s Area of Influence. The intensity of the potential impact was gauged by considering the following factors:

Direction

Direction describes the trend of the effect compared with baseline conditions. There are three options for direction:

- Adverse – effect is worsening or is undesirable.
- Neutral – effect is not changing compared with baseline conditions and trends.
- Positive – effect is improving or is desirable.

Geographic Extent

Geographic extent describes the quantitative measurement of area within which an effect occurs. Effects are described in terms of whether they are limited to the Project Footprint, Project Area of Influence, or extend farther:

- Local – effect is limited to the Project footprint.
- Regional – effect extends beyond the Project footprint, but is limited to the Project Area of Influence Local Study Area.
- Beyond regional – effect extends beyond the Project Area of Influence.

Duration

Duration refers to how long an effect lasts. Duration is described in relation to the phases of the development of the Project within the RSA, although effects may last longer than the phases of the Project for some valued components. The following framework was used:

- Short-term – effect is limited to the construction period (~2 years), or the period of decommissioning activities (~2 years).
- Medium-term – effect extends throughout the project operations, that is, 25 years.





- Long-term – effect extends beyond the 25 years of operation.
- Far future – effect extends more than 30 years after closure.

**Reversibility**

This criterion describes whether the effect is reversible or not. This can be associated with duration, as many effects eventually could be considered to be reversible (that is, in geological time). However, the extinction of a species can be considered as irreversible.

**Table 4: Intensity assessment rating scale**

Criterion	Rating scales
<b>Negligible</b>	Where the impact affects the environment in such a way that natural, and /or cultural and social functions and processes are negligibly affected and valued, important, sensitive or vulnerable systems or communities are negligibly affected.
<b>Low</b>	Where the impact affects the environment in such a way that natural, and/or cultural and social functions and processes are minimally affected and valued, important, sensitive or vulnerable systems or communities are minimally affected. No obvious changes prevail on the natural, and / or cultural/ social functions/ process as a result of project implementation
<b>Medium</b>	Where the affected environment is altered but natural, and/or cultural and social functions and processes continue albeit in a modified way, and valued, important, sensitive or vulnerable systems or communities are moderately affected.
<b>High</b>	Where natural and/or cultural or social functions and processes are altered to the extent that they will temporarily or permanently cease, and valued, important, sensitive or vulnerable systems or communities are substantially affected. The changes to the natural and/or cultural / social- economic processes and functions are drastic and commonly irreversible.

**4.7.3.2 Determination of Sensitivity of a Receptor**

Sensitivity for each Ecosystem Service supplied and/or demanded ranged from very low to high according to increasing level of threat (Table 5).

**Table 5: Sensitivity assessment rating scale**

Criterion	Rating scales
<b>Negligible</b>	None of the below
<b>Low</b>	Ecosystem service is readily substitutable or replaceable
<b>Medium</b>	Ecosystem service is substitutable or replaceable
<b>High</b>	Ecosystem service is not substitutable and/or irreplaceable

**4.7.3.3 Determination of Impact Severity/Significance**

Once the intensity of the impact and the sensitivity of the receiving environment were described, the severity of the potential impact was determined. The determination of significance of an impact is largely subjective and primarily based on professional judgment.

To provide a relative illustration of impact significance, it is useful to assign numerical descriptors to the impact magnitude and receptor sensitivity for each potential impact. Each is assigned a numerical descriptor of 1, 2, 3, or 4, equivalent to very low, low, medium or high. The significance of impact is then indicated by the product of the two numerical descriptors, with significance being described as negligible, minor, moderate or major, as in Table 6. This is a semi-quantitative method designed to provide a broad ranking of the different impacts of a project.





Table 6: Determination of impact severity

		Sensitivity of receptor			
		Negligible	Low	Medium	High
		1	2	3	4
Intensity of Impact	Negligible	1 Negligible	2 Minor	3 Minor	4 Minor
	Low	2 Minor	4 Minor	6 Moderate	8 Moderate
	Medium	3 Minor	6 Moderate	9 Moderate	12 Major
	High	4 Minor	8 Moderate	12 Major	16 Major

### 5.0 LAND COVER MAPPING AND CONDITION ASSESSMENT

The land cover types within the Local Study Area are illustrated in Figure 6.




Figure 6: Land cover in the Project Area of Influence

The area of each land cover category within the Local Study Area in relation to estimated land take from Project components (that is, footprint, escarpment road and pipeline) are shown on Table 7 and Table 8.

Land cover is treated separately for the Kingfisher Field Development Area of the Local Study Area (the Project footprint the escarpment road, and the pipeline servitude from the Buhuka Flats to the top of the





escarpment) and the pipeline route above the escarpment, as the dominant land cover types in these two areas are quite different.

**Table 7: Land cover in the Kingfisher Field Development Area and proportion lost as a result of proposed land take**

Kingfisher Field Development Area		Land take Area (Ha)	% of total
Land cover Class	Total Area (Ha)		
Dense Bush	248.5033	12.2915	4.95
Grassland	700.4368	62.3896	8.91
Open Bush	522.8448	33.9714	6.50
Settlement	29.0651	-	0.00
Subsistence cropland	19.4155	0.6740	3.47
Water	145.6080	-	0.00
Wetland	99.0343	0.5505	0.56
Woodland	133.6616	6.6261	4.96
<b>Total:</b>	<b>1898.5694</b>	<b>116.5031</b>	<b>29.34</b>

Grassland on the Buhuka Flats and open bush of the escarpment will suffer proportionately the greatest loss to the Project (Table 7); while subsistence cropland will be the most affected land cover type by the pipeline route servitude, beyond the escarpment (Table 8).

**Table 8: Land cover in the Pipeline area of influence and proportion lost as a result of proposed land take**

Pipeline Route		Estimated land take by 30 m pipeline servitude	
Land cover Class	Area (Ha)	Area (Ha)	% of total
Commercial cropland	6.6552	-	-
Degraded Bush	4502.7576	37.856	0.84
Dense Bush	565.3455	0.9282	0.16
Grassland	456.9595	2.7049	0.59
Mining	11.3849	-	-
Open Bush	4483.8372	28.0981	0.63
Plantation	4.8232	-	-
Settlement	20.9645	0.2666	1.27
Subsistence cropland	5314.6965	46.3545	0.87
Water	0.1651	-	-
Wetland	10.7274	-	-
Woodland	2930.6862	17.8371	0.61
<b>Total:</b>	<b>18309.0028</b>	<b>134.0454</b>	<b>4.98</b>





**5.1.1 Land cover condition**

The ecosystem condition of the various land cover types within the Local Study Area was estimated. Table 9 shows each of the affected mapped land cover categories as they relate to habitat types recorded within the Project Area of Influence, and their condition. This allowed a judgement on the condition of the land cover types to be made (and, therefore, their potential capacity to supply ecosystem services), based on the reported baseline condition of the habitat types supported therein.

**Table 9: Land cover classes, associated ecosystems, habitat types and condition**

Land cover class	Ecosystems	Dominant Habitat types	Estimated condition
Dense Bush Open Bush Grassland	Escarpment vegetation corridors	Open wooded grassland Dense wooded grassland Dense bushland Riverine bushland	Slightly degraded to moderately degraded
Degraded Bush Settlement Subsistence cropland	Settlement and cultivation areas	Open bushland and shrubland Grassland with thicket Open grassland	Heavily degraded
Wetlands	Wetlands	Permanent wetlands, seasonally flooded grassland	Slightly degraded to moderately degraded
Woodland	Bugoma Central Forest Reserve	Woodlands, Wooded bushlands	Slightly degraded to moderately degraded
Water	Near-shore habitats of Lake Albert	Shallow river-associated waters, open sandy shores, lagoons, large bays, rocky escarpments, open-water habitats	Near-Pristine

**6.0 REVIEW OF BASELINE BIODIVERSITY, SOCIOECONOMIC AND BIOPHYSICAL ENVIRONMENT OF THE LOCAL STUDY AREA**

The Project will be located on the south-eastern flank of the Lake Albert Basin, which is part of the western arm of the East African Rift System. This area is commonly known as the Buhuka Flats, situated in the administrative boundary of Kyangwali Sub-County in Kikuube District (Figure 2). The Project also includes the pipeline route to Kabaale (Figure 3).

This section presents a description of the existing environment within the Local Study Area, in relation to the supply of, and demand for, ecosystem services. In particular, it presents a summary of the relevant information distilled from the biodiversity and surface water baseline reports, in order to put the proposed Project area’s provisioning, supporting and regulating ecosystem services supply capability in context. Terrestrial and aquatic biodiversity is discussed in terms of vegetation communities and habitats and their condition, which directly relates to the results of the land cover condition assessment discussed in Section 5.0 above. This section also addresses drivers of ecosystem change that already exist within the Study Area, in the absence of the Project.

**6.1 Terrestrial and Aquatic Biodiversity**

Surveys of flora and fauna were conducted in the Project footprint and along the pipeline route, for the baseline biodiversity assessment (Eco & Partners, 2014). The data gathered on vegetation communities and fauna is presented in the following sections in the context of ecosystem service provision within the Local Study Area.

**6.1.1 Vegetation Communities Providing Ecosystem Services**

■ **Buhuka Flats**





The vegetation communities that dominate the Buhuka Flats include thicket-grassland mosaic, open grassland, wooded bushland and wetlands (Figure 7, Figure 9). These areas are a source of fodder for grazing cattle, and are over-exploited for grazing in many parts. The wetlands are a source of thatching material, wattle and mud 'daub' for traditionally built houses (Figure 8). Papyrus culms are also harvested from wetlands and used as a construction material for houses (NEMA, 2002).



*Thicket-grassland*



*Open grassland*



*Woodland (riparian areas)*



*Seasonally-flooded wetlands on shore of Lake Albert*

*Figure 7: Vegetation communities of the Buhuka Flats*





*Figure 8: Traditionally built house in process of being thatched*







### ■ **Escarpment**

The slopes of the Escarpment are characterised by wooded grassland, bushed grassland and bush-land and shrub-land (Figure 10, Figure 11). These vegetation types lend themselves to ecosystem services provision including wood for charcoal production, and subsistence hunting.

Tamarind (*Tamarindus indica*) is fairly common on the escarpment in woodland and wooded grassland; this species is commercially used for food in parts of northern Uganda (Katende, et al., 1995) though it is unclear whether it is used for this purpose in the Local Study Area. Due to its heavy exploitation, it is of conservation concern in Uganda and is on the Uganda Reserved Tree Species List of the National Forest Authority (Eco & Partners, 2014).

The total length of the proposed escarpment road is approximately 7 km and 9 metres wide, including shoulders. The construction of the road is likely to influence the supply and demand of ecosystem services. The loss of the wooded grassland and woodland will reduce the supply of ES such as availability of wood for charcoal production; the construction of the road will enhance access to wooded areas and this may increase demand for ES supplied by woodlands in these areas.

### ■ **Pipeline route**

The pipeline route to the refinery area and Kabaale is approximately 46 km in length. The majority of this part of the Local Study Area has been subjected to high intensity, subsistence agriculture, which has altered much of the original natural landscape (Forest Department, 2002); the dominant vegetation type along the pipeline route is cultivated land associated with settlement (Figure 10).

Widespread cattle grazing and charcoal manufacture have put significant pressure on the few natural vegetation communities in the vicinity of the pipeline route, which consist of fairly degraded seasonal wetlands, riverine bushland along Hohwa River and pockets of natural woodlands. The effects of large-scale conversion of natural habitats to subsistence agricultural fields are particularly noticeable between the escarpment proper and Bugoma Central Forest Reserve.

#### **6.1.1.1 Invasive plant species**

A number of invasive plant species are present throughout the Local Study Area. *Mimosa pigra* (Giant Sensitive Tree), *Lantana camara* (Lantana), and *Eichhornia crassipes* (Water Hyacinth) were the commonest species recorded, predominantly on the Buhuka Flats and the shore of Lake Albert, which are recognised as some of most noxious weeds in the world (Lowe, et al., 2000).

Several of the invasive species recorded in the Local Study Area provide ecosystem services to local communities and are planted specifically for this purpose, including Castor oil (*Ricinus communis*), Neem (*Azadirachta indica*), *Jatropha* (*Jatropha curcas*), and *Parkinsonia* (*Parkinsonia* sp.):

**Castor oil:** planted and used for medicinal purposes in the LSA.

**Jatropha:** its non-edible seeds are harvested for biodiesel production; it is also planted on graves in the LSA

**Neem:** used in East Africa for various medicinal purposes including the treatment of scabies and head lice, and its hard wood is harvested for use in construction (BioNET-EAFRINET, 2011).

**Parkinsonia:** may be used for firewood, charcoal production, medicine, fodder, shade, mulch, as a 'live' fence, and as a windbreak for soil stabilisation (BioNET-EAFRINET, 2011).



Wooded Grassland



Bushland and shrubland



Bushed grassland



Wetlands associated with Escarpment ravines

Figure 11: Vegetation communities of the Escarpment

### 6.1.2 Faunal Communities Providing Ecosystem Services

#### 6.1.2.1 Freshwater Fish Communities of Lake Albert

Nile Perch (*Lates niloticus*), *Ragoge* (*Brycinus nurse*), *Ngassa* (*Hydrocynus forskahlii*), Shield-head Catfish (*Synodontis schall*), Black Nile Catfish (*Bagrus bajad*), *Muziri* (*Neobola bredoi*) and *Angara* (*A. baremoze*) made up the majority of the wet and dry season baseline fish survey catch, and are some of the most commercially important species in Lake Albert (Taabu-Munyaho, et al., 2012).

In general, the fish community in the near-shore zone of the LSA is composed of a fairly uniform, multispecies mix of various ages in good condition. The diversity, age classes and condition of the species assessed is a reflection of adequate food and a healthy environment.

The near-shore artisanal fishery is dominated by gillnets, and is mostly focused on Nile Perch, *Ragoge*, *Ngassa*, and *Angara*; whilst *Muziri* features strongly in the seine net fishery. The condition of the near-shore aquatic habitats that support the fishery is considered Near-Pristine (ref. Section 5.1.1); therefore the capacity of this ecosystem to continue to supply this ecosystem service is high, based on its current status. Further information on the fisheries value of the Lake Albert fish community to beneficiaries is provided in Section 6.4.1.

#### 6.1.2.2 Terrestrial Fauna Hunted for Bush Meat

A study of bush meat consumption in Uganda found that bush pig (*Potamochoerus larvatus*), cane rat (*Thryonomys* sp.), guinea fowl (*Numida meleagris*) and kob (*Kobus kob*) were the main sources of bush



meat in the Murchison Falls area (Olupot, et al., 2009), which is the nearest (approx. 150 km northeast) studied area to the ecosystem services Local Study Area. Other species taken for bushmeat in Uganda include bushbuck (*Tragelaphus scriptus*), duikers (*Cephalophinae*), oribi (*Ourebia ourebi*), buffalo (*Syncerus caffer*), hippopotamus (*Hippopotamus amphibius*), redtail monkey (*Cercopithecus ascanius*) and black-and-white colobus monkey (*Colobus* sp.). However Olupot et al. (2009) reported that bush meat was eaten by a comparatively low number of respondents (5-32%) when compared to the (reported) consumption of livestock meat and fish (94-100%).

Due to the heavily human-impacted nature of the Local Study Area, vegetation cover is compromised and the potential of the area to support terrestrial mammal species is considered limited (Eco & Partners, 2014); however, a few medium sized mammals were recorded during the biodiversity baseline studies. These tended to be thicket and dense bushland specialists, such as bushbuck and duiker. The low populations and diversity of these species could also be a reflection of increased pressure for bush meat from the local human population, which has increased markedly over the last ten years (AECOM, 2012).

The socio-economic baseline report makes reference to hunting occurring in several villages, and hunting of 'rats' as an alternative livelihood in the village of Kamukeduke; these, together with mammal species that were reported (during interviews with locals, and trapping and observation of track/sign during field work) in the biodiversity baseline study of the LSA, that are known bush meat sources in the area, are outlined in Table 10.

Table 10: Mammals recorded in the KFDA LSA that are bush meat sources<sup>2</sup>

Common name	Species
Hippopotamus	<i>Hippopotamus amphibius</i>
Bush Duiker	<i>Sylvicapra grimmia</i>
Lesser Cane-rat	<i>Thryonomys gregorianus</i>
Northern savanna multimammate rat	<i>Mastomys hildebrandti</i>
Black rat	<i>Rattus rattus</i>

The biodiversity baseline report identifies bush meat hunting as a driver of change in the Local Study Area, particularly in Bugoma Central Forest Reserve (NEMA 2010, Plumptre et al. 2010). Between 2011 and 2012, increased immigration into the areas surrounding the forest resulted in increased deforestation, with an estimated 5,000 ha of the forest subject to encroachment by about 1,000 families and pit-sawyers (AECOM, 2013). In Bugoma CFR, trapping of large mammals now appears to have declined, which is attributed in part to the decline in large mammals to such an extent that hunting is not very productive any more (Plumptre et al., 2010).

A study of nearby Budongo Central Forest Reserve (Zommers & MacDonald, 2012), identified that of the local communities that hunted bush meat in the forest, nearly 73% were immigrants to the area; and furthermore, that the households of immigrants were also more likely to be involved with deforestation.

Overall, the indication is that bush meat is a resource utilised by beneficiaries within the Local Study Area, and may be of greater importance to immigrant populations than residents who have established subsistence crops or livestock grazing areas. Pressure on faunal species that are hunted for bush meat is thus expected to increase as a result of population influx associated with development of the Project.

## 6.2 Surface Water Systems

Lakes and rivers provide provisioning services in the form of water supply, food (fish), and vital regulating services such as groundwater recharge, water storage, flood control and water purification/waste assimilation (IPIECA, 2011).

<sup>2</sup> Based on WCS study (Olupot, et al., 2009) and ESIA survey 2014





The information presented in the following sections contextualises the baseline information presented in the Surface Water baseline report in terms of supply of fresh water to the Project and to beneficiaries, and regulating ecosystem service provision within the Local Study Area.

### 6.2.1 Surface Water Resources in Local Study Area

The location of the proposed Project in relation to regional surface water features and topography is shown in Figure 12 and Figure 13 respectively. The local surface water features in relation to the Project are illustrated in Figure 14.

Hydrologically, the Project is located within the Lake Albert catchment, which drains westwards from the escarpment into the south-eastern shores of Lake Albert. Lake Albert's catchment is strongly associated with the adjacent escarpment, draining into the lake via several streams flowing westwards. Surface water bodies within the Project's Area of Influence include the Kamansinig and Masika Rivers. Various other streams also flow off the escarpment and either join the main rivers mentioned above (such as Masika) or gradually and independently feed Lake Albert. The area below the escarpment is approximately 13 km<sup>2</sup> and, besides the rivers mentioned, is characterised by relatively scattered wetlands at an elevation level associated with most Project infrastructure (Figure 13).

The escarpment catchment generates runoff during the rainy season that discharges onto the Buhuka flats via ravines. High energy ravine flows are quickly dissipated as the slope meets the flats and the bushy vegetation at the bottom of the escarpment further slows the flow of water. Streams that are large enough slowly make their way through densely vegetated wetlands to Lake Albert. Some of the smaller streams disappear from the surface a few hundred metres away from the bottom of the escarpment; this shows that the zone at the bottom of the escarpment is an important zone of recharge of water into the soil.

Water quality results for the baseline assessment were compared with the local Ugandan Acceptable Standards for drinking standards (NEMA, 1996), and the World Health Organisation (WHO) for Drinking Water (WHO, 2011). Overall, the water quality in the Buhuka Flats area during the dry season is generally good; during the wet season there is potential for humic acids (from surrounding land areas such as wetland systems) to increase pH levels and introduce metals into Lake Albert.

### 6.2.2 Ecosystem Services Provided by Surface Water Systems in Local Study Area

The zone at the bottom of the escarpment is an important zone of recharge of water into the soil; therefore, this area has a role in provision of regulating services in the Buhuka Flats area, including groundwater recharge, water storage and flood control. Scattered wetland areas in the Buhuka Flats consist of riparian floodplains, and permanent wetlands at the Lake's edge. These wetland areas also provide regulating ecosystem services including flood attenuation and sediment retention.

Main water sources within the villages along the pipeline route include springs, streams and boreholes, which are used for drinking and domestic purposes as well as animal watering. These water sources provide typically poor quality drinking water, which commonly causes various illnesses in the local communities.

The communities of the Buhuka Flats region source drinking and bathing water from either Lake Albert or from the gravity flow scheme (which is contaminated with human waste). These households also dispose of solid waste and waste water into the Lake; therefore, the lake has importance both as a source of drinking water, and for provision of some waste assimilation services.

The construction and operation of the Project has potential to influence or change the processes that drive these systems and thus their capacity to supply ecosystem services. For example, the road leading from the foot of the escarpment and the borrow pit has noticeably influenced the flow regimes and drainage patterns of the seasonally flooded grasslands associated with the Kamansinig River, resulting in the alteration of the wetland on the western side of the road and associated loss of function. Construction of the pipeline route is expected to intercept several permanent wetland systems which may alter flow regimes and ultimately affect the capability of the wetlands to provide services such as flood attenuation or nutrient (waste) assimilation.



### 6.2.3 Project Water Demand

During the operational phase, the Project will require between 520 m<sup>3</sup>/day and 7 315 m<sup>3</sup>/day of water, which equates to between 191 260 MI/annum and 2 669 245 MI/annum. This equates to between 0.00048% and 0.00654 % of the average inflow into Lake Albert. This is much less than the monthly variations observed naturally at Lake Albert. Therefore, the supply of fresh water provided by the Lake is not expected to be significantly affected by the water required for Project operation.









# ECOSYSTEM SERVICES ASSESSMENT




Figure 13: Regional Topography





# ECOSYSTEM SERVICES ASSESSMENT




Figure 14: Local context of the Project in relation to surface water features and floodlines





### 6.3 Land Use, Land Capability and Soils

Land uses in the Local Study Area were assessed as part of the baseline study for the ESIA. Land capability was determined based on the types of soil present in the areas investigated.

In the Buhuka Flats, 'natural grassland' consisting of thicket-grassland mosaic, open grassland, and seasonally flooded grassland vegetation categories, is the dominant existing land use (Table 11). This area is currently used for grazing cattle. The land potential of this area is appropriate for arable crop production. Wetlands (approx. 19%) and settlements (approx. 16%) account for the majority of the remainder of land use in the Buhuka Flats area.

Land use along the pipeline route is dominated by 'cultivated land' (Table 11). Patches of permanent wetland, seasonally flooded grassland, and open bushland and shrubland occur along the pipeline route, these are classified as 'natural grassland' in terms of land use category. The land potential of the areas classified as 'natural grassland' is also considered suitable for arable crop production.

Table 11: Land use within the Local Study Area

Area	Land Use	Surface Area (ha)	% of Total
Buhuka Flats	Existing CNOOC Base & Airstrip	24	3.16
	Natural Grassland	468	61.57
	Cultivated Land	5	0.66
	Villages and immediate surrounds	120	15.79
	Wetlands	143	18.82
Pipeline Route	Natural Grassland	396	17
	Cultivated Land	1,923	83

Examination of the land use categories in the Local Study Area, shows that livestock grazing opportunities provided by natural grassland and wetland categories accounts for approx. 80% of the land use of the Buhuka Flats area. This highlights the importance of this area for provision of grazing ecosystem services. Almost no crop cultivation occurs in this area.

Land use along the pipeline route is dominated by cultivated lands (83%). These areas are used for subsistence farming and small-scale commercial farming, which are the main economic activities along the pipeline route. Production is carried out on small farm holdings less than 1 acre in size (Figure 15), and crops are mostly used for household consumption or sale in community markets.



Figure 15: Backyard farming in Nsonga, Buhuka Flats





### 6.4 Baseline Socio-economic Setting

The information in the following sections is largely derived from the interviews done as part of the socio-economic baseline study. Other relevant information sources are referenced as appropriate throughout the text.

The Local Study Area is starkly different from the general district and national trends in terms of primary economic activity. While agriculture is a major economic activity in the district, the Local Study Area is mostly characterised by fishing and livestock (mainly cattle) keeping as economic activities. The villages in the Buhuka Flats are predominantly engaged in fishing and cattle farming as a livelihood, whilst villages on top of the escarpment engage in agricultural crop farming activities as a main livelihood activity. A substantial amount of processing and trade across Lake Albert occurs with the fish produce, whilst trade from the top of the escarpment to the Buhuka Flats is somewhat limited due to poor access. More details regarding economic activities are provided in the following sections.

The total estimated household count in the Buhuka Flats is 2,831. On average, households have 8 members, indicating that the approximate population size in the Buhuka Flats area is 22,000 persons. This average household size is much larger than the regional average household size of 4.9 persons per household. There has been a substantial increase in population over the past 10 years (between 99% and 184%), driven by a multitude of factors such as regional instability, attractive livelihood opportunities to engage in fishing on Lake Albert and more recently, interest in capitalising from opportunities related to oil and gas developments.

#### 6.4.1 Fisheries

At least 75% of the households in Buhuka Parish villages directly depend on subsistence fishing activities as a source of food, livelihoods and a cash income. The fish caught in the district is spread over 68 landing sites in the district which are distributed quite evenly along the shoreline. Drying and salting of fish along the lake shore (Figure 16) is undertaken before selling the fish at markets in Hoima and Kikuube Districts; however, according to focus group discussions, relatively little of the fish catch goes up the escarpment for trade (about 25%). Village residents from the top of the escarpment are more likely to travel to the lake to catch their own fish for household consumption and transport it back to their houses than to purchase fish from the markets.

Most of the catch that is traded, is channelled into the Democratic Republic of Congo (DRC) and distant trading posts in the north (Panyimur, Ntoroko and Bwera), which have easier access routes by boat or roads than transporting fish up the escarpment for sale to villages at the top. The fishing trade (including both fishermen and traders) forms part of the informal economic sector, and therefore earnings are largely undocumented. What is clear, is that earnings are unpredictable in size and occurrence. Fishing has greatly influenced social and economic development in the sub-county of Kyangwali where the Project will be located.

There are numerous reports of declining fish numbers, driven by unsustainable fishing practises and an increasing population engaging in fishing activities. The fishing sector is also being threatened by declining catches, mainly due to the use of destructive and illegal fishing methods, fishing in breeding areas, non-compliance with regulations and inadequate control of catches. Poor fishing gear and techniques result in taking of immature fish before they have the chance to reproduce and maintain the fish population. Other factors that affect fishing activity on the lake shore include restrictions on illegal fishing methods and poaching put in place by **Beach Management Units (BMUs)**; however, limited funds and inadequate coordination between BMUs and fisheries authorities limits the conservation capacity of these organisations (Uganda Nile Discourse Forum, 2013). Most fisheries are formally open-access with no legal controls in place to prevent entry to fisheries (Scullion, 2007). The impact of oil exploration activities on fishing is thought to be the least detrimental impact on current amounts of fish caught. Indirect effects of oil exploration activity, such as population influx, is likely to increase in the number of people involved in fishing, putting fish stocks under increasing pressure which may result in fish stock decline.





Figure 16: Drying Nile perch at the shore of Lake Albert

## 6.4.2 Agriculture

Subsistence farming and small-scale commercial farming are the main economic activities in the Kikuube and Hoima Districts. The majority of people along the pipeline route are dependent on agriculture, with crop production as a major economic activity, followed by poultry and livestock. However, the Buhuka Flats is notably different, with households being heavily dependent on fisheries rather than agriculture. Where subsistence farming is engaged in, the dominant activity in this area is livestock keeping, rather than the growing of crops.

### 6.4.2.1 Buhuka Flats

Very few households grow agricultural produce in the Buhuka Flats, and instead purchase this from Ngoma on top of the escarpment. The steep escarpment footpath (which is the only access route between the escarpment and the Buhuka Flats) discourages people to travel to Ngoma to buy agricultural produce; therefore, the diet of the local population is comprised mostly of fish and lacking in vegetables and other foodstuffs.

Households in the Buhuka Flats that do grow crops, do so on small backyard plots. Crops typically consist of cassava, maize, beans, sweet potatoes, tomatoes and *matooke* (bananas). Some of the food crops are sometimes sold in order to raise money needed for other household needs. Besides cultivating on open spaces in the backyard of their homesteads, approx. 33% of Buhuka Flats respondents reported that they also had access to arable land away from their homesteads, the majority of which (approx. 70%) is located on top of the escarpment and the rest comprising plots of land in the Buhuka Flats. The average size of land currently being cultivated according to the respondents is 3 acres and the most common size is 1 acre.

On the Buhuka Flats, livestock keeping is undertaken by a large percentage of households (approx. 80%). Livestock numbers are large. Carrying capacity calculations suggest that there is a high degree of



overgrazing (approx. double the carrying capacity) already in both the Buhuka Flats and neighbouring areas into which cattle keepers range to find alternative grassland sources.

In recent times, there has been an influx of cattle keepers from as far as Tanzania and Kasese areas leading to a tremendous increase of cattle in the Kyangwali sub-county, including the Buhuka Flats. In Buhuka, the cattle keepers were attracted mainly by the abundance of water from the lake and the open grasslands that are conducive for grazing animals.

### 6.4.2.2 Pipeline Route

Most of the pipeline route is covered by cultivated areas (Figure 6). The most common subsistence crops include bananas (for food), bananas (for beer), cassava, sweet potatoes, Irish potatoes, cotton, soybeans, groundnuts, pigeon peas, beans, sorghum and maize, whilst perennial crops including coffee, banana and sugar cane plantations and tree plantations (pine wood, eucalyptus), which are typically commercial crops. Beekeeping for honey production is practised in a number of villages along the pipeline route.

Both the crop farming and livestock sectors in the Kyangwali sub-county are faced with a number of challenges and these include: unpredictable weather changes, vermin attacks, crop diseases, poor farming practices which is leading to shortages of land for cultivation, and environmental degradation/deforestation because of the increase in clearing of land for farming practices.

### 6.4.3 Freshwater

On the Buhuka Flats, the main source of water for household consumption is piped water from the gravity flow scheme. The water is sourced from open water sources at the top of the escarpment and is then piped down to the Buhuka Flats area. Approx. 79% of the Buhuka population buy their water from taps and 15% draw their water from the lake (Environmental Assessment Consult (U) Ltd., 2013). Poor quality drinking water is a challenge for most villages in the Local Study Area, (for example, in Kyakapere village, tap water is reportedly polluted so water is drawn from the lake and boiled) and water-borne diseases are prevalent.

In the vicinity of the pipeline route, villages obtain water from sources including boreholes, wells, water holes, streams/rivers and swamps. In many cases, animals/livestock use the same sources for drinking water, and the bad quality of drinking water contributes to the spread of various illnesses. These water sources are considered sensitive where construction of the pipeline will take place.

### 6.4.4 Land tenure and use

Both Hoima and Kikuube districts cover a total land area of 3,612.17 km<sup>2</sup>. Out of this, 2,853.48 km<sup>2</sup> (79.1%) is under agriculture, settlement and other miscellaneous land uses. The remaining 758 km<sup>2</sup> (20.9%) is under protected areas, including forest reserves and wildlife conservation areas. Protected areas are classified as Public Land and include forest reserves, wetlands and any other land outside customary, free-hold or lease-held land.

On private land, customary land tenure (both individual and communal) is the most widely practiced system. The lack of a uniform land tenure system, presents management challenges particularly with regard to land speculation that has been exacerbated by the discovery of oil in the district. This could cause land use conflicts and result in landless households and communities in the district, as land purchases and delineation of previously communally owned land continue to take place. Such conflicts have the potential to affect the Project's social license to operate.

### 6.4.5 Human health risk

Most urban populations in Uganda lack water-borne sewage systems, and domestic wastes often flow directly into swamps and wetlands, which provide important water purification services (NEMA, 2002).

No formal waste disposal services or facilities exist in either Kikuube or Hoima District, and, therefore, by implication, the Local Study Area. This lack of sanitation and refuse disposal systems has implications for health conditions of communities. It also increases the importance of the role played by swamps and wetland in absorption of nutrients and waste assimilation, in the absence of formal municipal treatment schemes.



### 6.4.6 Recreation and Tourism

Ecosystems and biodiversity play an important role for many kinds of tourism, which in turn provides considerable economic benefits and can be a vital source of income for many countries (TEEB, 2010). Tourism is typically based on the use of both natural and cultural ecosystem services (PANParks, unknown). In addition, people derive recreational pleasure from natural or cultivated ecosystems (Landsberg, et al., 2013)). In Uganda, tourism is focused in the approximately 21,000 km<sup>2</sup> of gazetted wildlife protected areas (NEMA, 2002), where tourists pay to view wildlife in remote and pristine natural settings. The aesthetic value of the Lake Albert landscape offers cultural ecosystem services to tourists, in the form of inspirational experiences.

The development of accommodation establishments around Lake Albert (for example, Lake Albert Safari Lodge and Lake Albert Guest House) has been attributed to the developing oil industry, the employees of which are thought to be boosting tourism in the area (Solomon & George, 2012). Although Lake Albert has potential for community tourism, with activities such as sport fishing and canoeing/kayaking considered feasible; most landing sites on the lake shore have not been developed for tourism and lack the necessary facilities and amenities (Solomon & George, 2012). However, oil industry-related activities may negatively impact tourism potential through land take and associated impacts on flora and fauna, potential pollution and primarily, visual intrusion. In order to assess the impacts of oil activities on tourism; number of species, number of tourists, tourism revenue, visual impact and habitat quality would need to be monitored.

## 6.5 Cultural Heritage Context

Due to the confidential nature of some of the cultural sites (for example, secret sites) their exact locations were not given in the baseline report. An overview of cultural sites present within the Project Area of Influence is provided in Figure 17.

### 6.5.1 Spiritual Values

People attach spiritual, religious, aesthetic values to ecosystems, landscapes and species (Landsberg, et al., 2013), which are non-material contributions of ecosystems to human well-being. The ecosystems, sites and landscapes that contribute to this ecosystem service in the Local Study Area are discussed in the following sections.

#### 6.5.1.1 Sacred Sites

Sacred sites within the Project Area of Influence fell into four broad categories:

- Ritual Sites
- Sacred Rivers
- Sacred Trees and/or Cultural Trees
- Ritual Objects.

#### *Ritual Sites*

- 'Luzira' is a lagoon close to the Project footprint. It is an active place of worship and the historic centre of cultural activity, where, traditionally, pilgrims would travel some distances to stay within the swamp for nine consecutive days, to make offerings and conduct ritual activities.
- Kasonga beach, near the village of Nsonga, is used for ritual ceremonies specifically related to fish catches – when lake stocks appear low or when fishermen have troubles.
- A secret sacred pool is located on the River Masika. It is well known by elders and taboo for younger members of the community. The site is utilised for rituals during cholera outbreaks in particular, and a specific ceremony for sick babies is also conducted there.





### Sacred River

- The River Masika in itself is considered sacred. Areas on the river bank are used for ceremonies to improve fish catches and occasionally to cure sick children.

### Sacred, Cultural and Barkcloth Tree

- A secret tree (species unidentified) at one of the lakeshore villages has a number of taboos associated with it. Another such tree is present near the escarpment road.
- A Barkcloth Tree (*Ficus natalensis*) near the pipeline route at Kaseeta – barkcloth making is an ancient craft listed on UNESCO's intangible world heritage list.
- A tree in Nsonga village is the village assembly tree.

### Ritual Objects

- These were recorded at three locations - a village in Buhuka Flats, escarpment road and along the pipeline route. They consisted of stones used for feet washing, rituals and worship.

#### 6.5.1.2 Cultural landscapes

Both Lake Albert and the escarpment are iconic features of the natural landscape, defining the local (communal) sense of place and apparent (traditional) cultural associations of the natural features (rivers, lakes, trees). Both sites provide a strong historic and religious focus for the lakeside communities in particular, evident within the oral traditions and the sacred places associated with both locations (Box 1).

*“There is a tradition that (as a sign of respecting the fish and ensuring their continued supply from the lake) if a woman comes from the escarpment top with cassava flour, it’s up to her husband to prepare it to accompany a fish. When the woman leaves the lake shore to return she is then given a fish to take back up the escarpment top”*

**Box 1: Example of Oral Tradition associated with Escarpment**

#### 6.5.2 Inspirational Values associated with Lake Albert

People in the Local Study Area have derived various elements of information from the ecosystems surrounding them in Lake Albert, and used this information for aspects of culture, traditions, art and story-telling. These practices that are intrinsic to Lake Albert contribute to the local people's sense of place.

##### Local legends and revered animals

Throughout the lakeside villages consulted during the community survey, some common beliefs and practices were identified. Some traditional practices call on a giant snake (seen to bring good fortune) or a giant crocodile (can bring or take away fish shoals).

##### Beliefs associated with Lake Albert

A number of taboos and beliefs are specifically associated with Lake Albert, such as ceremonies to cure sick children and bless new boats and nets, beliefs that evil spirits roam the area at night time, and conducting of certain activities during the night is a sign of disrespect to the ancestors, and taboos relating to women bathing in the Lake, amongst others.

##### Oral history and village naming

The names of many of the lakeshore villages are directly derived from natural features (e.g. Nsunzu is the name of the type of grass that is good for feeding cattle).



### Medicinal plants

About 80% of the population in Uganda depend on traditional plant medicines derived from >300 plant species (CRA, 2006). Medicinal/traditionally used plant species observed by the cultural heritage field team are summarised in Table 12; specific knowledge on the exact identification (taxonomy) of these plants was not gathered. 'Secret plants' were also mentioned to be used by women in labour, specifically to address complications with the afterbirth.

**Table 12: Traditional Plant names and uses – Appendix A,**

Local Plant name	Medicinal Use
Omwoyante	Malaria (boiled leaves)
Kyangwe	Used for sponges; leaves for ringworm
Tengo	Backache; bilharzia; teeth: use the roots; remove poison: fruits; eibisebe: flowers
Omulisana	Ringworms, hook worms
Omukoma	Constipation; allergy: the stem
Ekiryabiruku	Cough
Kibeere	Cough, making the placenta stable
Omususa	Allergy, syphilis
Omupeera (guava)	Cough (leaves), asthma (roots)
Omusheshe	Allergy (leaves and roots), syphilis (stem),
Omukwatange	Fibroids (stems and roots)
Akagando	Wounds locally called ebironda (leaves), asthma (roots), Allergies, Teeth
Omululuza	Worms, malaria, wounds, and burns. Note: it has to be picked early in the morning when the chlorophyll is in the leaves but if the roots are to be used it is advisable to pick them in the evening.
Omushebashebe	Pancreas (stem to be drunk).

### 6.5.3 Intangible cultural heritage

A number of intangible heritage practices were identified during interviews with the local community within the Local Study Area that can be associated with certain ecosystem services.

#### Hand-crafted ghee storage gourds

Ghee-making is traditionally practised by the Balalo pastoralist community ('cattle keepers') who were interviewed near the Project during the cultural heritage field work. The hand-crafted ghee storage gourds also traditionally made by this group were displayed by the community members to the cultural heritage field team. The use of biological raw materials for fashioning ornamental/functional items, such as a ghee storage gourd, is a recognised ecosystem service (Landsberg, et al., 2013)

#### Making bark-cloth

Bark-cloth making is an ancient craft performed by the Baganda people from the Buganda kingdom in south Uganda (UNESCO, 2005). Although no direct evidence of bark-cloth making within the Local Study Area was gathered, a tree that can be used for bark-cloth making (*Ficus natalensis*) was identified in proximity to the pipeline route; however, there was no evidence of use for this purpose on that individual.

#### Traditionally built houses

Hand built shelters for ducks and chickens were observed within the villages surveyed by the field team. Traditionally built houses, without any manmade materials, were also prevalent amongst the isolated communities on the Buhuka Flats. The practice of constructing a house is done by men only, with women only allowed to smear the house to finish off the building. Materials used in construction of these shelters and houses are mud (daub) and 'wattle', which is wood harvested from trees. It is unclear exactly which tree species are harvested to provide the timber for the wattle. One estimate for Uganda suggests that if the stock of wattle and daub housing in Uganda was to be maintained, 136,000 trees per year would be needed



to supply enough wood (European Commission, 2001). Grasses and reeds are used to thatch the roof (Figure 8).




Figure 17: Overview of cultural sites and objects identified in the Local Study Area

### 6.6 Existing Drivers of Ecosystem Change

Ecosystems are directly affected in two ways; by natural processes (for example, desertification) that cause continual change in species composition and habitat structure across regions; and by human activities and





increased human population pressure. Five direct drivers of ecosystem change directly attributable to human activity have been identified (Landsberg, et al., 2013); changes in local land use and land cover, harvest and resource consumption, pollution, introduction of invasive species, and climate change. These are considered to have the greatest effects on ecosystem health and condition, and therefore the supply of ecosystem services. In summary, the existing land cover within the Local Study Area is already modified by the five main drivers of ecosystem change, which are directly attributable to human activity and behaviour.

### ■ *Changes in local land use and land cover*

The natural vegetation of the Buhuka Flats region has been substantially modified, largely due to over-grazing and uncontrolled harvesting of trees for wood and charcoal production, and house construction.

Along the pipeline route, the natural vegetation has been replaced by a mosaic of commercial crop production, subsistence farming, with remaining non-cultivated areas consisting of heavily-modified grassland, wetland and woodland habitats. The occurrence of frequent fires was also evident on the escarpment. Over-frequent fire is known to detrimentally affect the functioning and processes of savanna ecosystems (Smith, et al., 2013), thus may be a driver of change in land cover condition, and, therefore, ecosystem condition and ecosystem service delivery in this part of the Local Study Area.

As mentioned in the next section, population influx has increased demand for ecosystem services such as grazing for cattle; livestock grazing has now been extended into areas that might not previously have been typically grazed, such as dense reed stands in wetlands and difficult-to-access open bushland areas of the escarpment. Such changes in land use and land cover are likely to affect spiritual sites, as these sites have typically been selected due to their remoteness, natural setting, and difficulty to access. Further changes in land use and land cover in the LSA as a result of the Project are anticipated.

### ■ *Harvest and resource consumption*

Fishing activities based out of both the fishing villages in the Buhuka Flats and villages further up the escarpment, puts pressure on the local fish populations. Apart from the published accounts of the decrease in commercial fish stocks (Wandera, 2000), (Wandera & Baliwra, 2010); Taabu-Munyahu et al. 2012), anecdotal accounts from the local fishermen also identify noticeable decreases in catches of fish per unit effort over the years. The main drivers of change influencing the vegetation communities along the escarpment are from livestock grazing, fuel wood harvesting, charcoal manufacture and the conversion of natural vegetation for subsistence agriculture. This is particularly noticeable along the pipeline route.

The keeping of livestock forms a substantial component of the local socio-economic structures in the Local Study Area, and particularly on the Buhuka Flats. Livestock numbers are large and there is strong evidence for overgrazing on the Buhuka Flats extending up onto the escarpment.

Large trees on the escarpment are becoming rarer as these individuals are selectively harvested for the manufacture of charcoal, which is typically then sold. Smaller woody species are regularly harvested for fuel wood used directly in the fishing villages. The harvest of fibre and other house construction materials is common on the escarpment. For example, thatching grass is regularly harvested on the escarpment and transported to the local fishing villages of the Buhuka Flats (Figure 18).

There has been a substantial increase in the local population over the past 10 years, driven by factors such as regional instability, livestock grazing opportunities, attractive livelihood opportunities to engage in fishing on Lake Albert, and more recently, interest in capitalising from opportunities related to oil and gas developments. This population influx is expected to exacerbate demand for numerous ecosystem services already used by the local population, including fisheries, fuel and construction wood, and bush meat, which will have knock-on effects on the extent and condition of the ecosystems that supply them.





Figure 18: "Chutes" used for the transport of thatching grass harvested from the escarpment

### ■ **Pollution**

Despite the human impacts evident throughout the Project Area of Influence, industrial/commercial facilities and/or concrete hard-standing typically associated with pollution potential are generally absent. Pollution of surface water and groundwater resources and air pollution has not been highlighted as a major issue in the baseline biophysical reports; however Wandera (2000) and Wandera and Balirwa (2010) have identified that agricultural run-off is having real effects on increasing the nutrient levels of Lake Albert, contributing to eutrophication of the lake's waters. In addition, the presence of approximately 22,000 people on the Buhuka Flats and other nearby villages who do not have access to running water and sanitation, will put large pressures on the nutrient loading of the inflowing waters of Lake Albert.

Many ecosystem services are reduced when inland waters and coastal ecosystems become eutrophic; water from lakes that experience algal blooms is more expensive to purify for drinking or industrial uses, fish populations can be reduced or eliminated, and many of the cultural services provided by lakes can be lost, for example, odours of rotting algae, slimes, and toxic chemicals produced by some blue-green algae during blooms keep people from swimming, boating, and otherwise enjoying the aesthetic value of lakes (Nelson, 2005).

### ■ **Introduction of invasive species**

A number of invasive plant species have been observed throughout the Local Study Area, including lantana, castor oil, neem, jatropha and parkinsonia. Some species e.g. castor oil are proliferating in uncultivated areas of the escarpment. Invasive aquatic plants including water lettuce (*Pistia stratiotes*) and water hyacinth (*Eichhornia crassipes*) have been recorded on the shores of Lake Albert itself (Figure 19).

Many invasive plant species do have the capacity to change vegetation community composition and the ecosystem services provided by those vegetation communities. This is particularly the case for water hyacinth, which can reduce biological diversity through reduction of water oxygen levels (e.g., oxygen-sensitive fish species may be affected), blocking sunlight to native plants, and blocking access to water to some animal communities (Center for Aquatic and Invasive Plants, University of Florida, 2014). Water hyacinth has already become dominant of the marginal ecotones of Lakes Kyoga and Victoria in the Rift Valley system, which has resulted in negative impacts on aquatic biodiversity and fishing activity in marginal



habitats (Twongo, 1996). Spread of water hyacinth in Lake Albert could therefore become a major driver of change in the near-shore aquatic habitats of the Lake in the future.



*Invasive castor oil plant on escarpment*



*Invasive Pistia stratiotes on shore of Lake Albert*

Figure 19: Examples of invasive species in the Project Area of Influence

### ■ **Climate change**

Observed changes in climate, especially warmer regional temperatures, has already effected changes in species distributions, population sizes, and the timing of reproduction or migration events, as well as an increase in the frequency of pest and disease outbreaks, especially in forested systems (Nelson, 2005).

Uganda is expected to experience more extreme periods of intense rainfall, and erratic onset and cessation of the rainy seasons and more frequent episodes of drought due to future climate change (Global Climate Change Alliance, 2012). Recent increases in the range of water level fluctuations in Lake Albert have been attributed to climate change (International Lake Environment Committee Foundation, 1999). For example, recent flooding on the Buhuka Flats (2012) which may be attributable to climate change, caused the death of 10 people, the loss of 70 homes and other household losses.

## **7.0 ECOSYSTEM SERVICE REVIEW FOR IMPACT ASSESSMENT**

### **7.1 Step 1: Identification of Relevant Ecosystem Services**

The Project's area of influence hosts numerous ecosystem services. The following sections characterise the supply of ecosystem services within the Project's area of influence, their quality, and their level of use or value to beneficiaries.

Beneficiaries considered most likely to be affected by the Project are the inhabitants and herders of the Buhuka Flats, and subsistence farmers whose plots lie within the servitude of the proposed pipeline route.

#### **7.1.1 Land Cover Types / Ecosystems which could be impacted by the Project**

Given the limited amount of non-degraded natural habitat types within the Project footprint, the extent (Table 7, Table 8) and ecosystem condition (Table 9) of the various land cover types within the Project footprint was estimated. The ecosystems that could be impacted by the Project are discussed according to the relevant area of the Local Study Area.

##### ***Buhuka Flats and Escarpment***

- The dominant land cover classes in the Buhuka Flats (grassland) and the Escarpment (open bush and dense bush) will have the greatest proportionate loss to the Project footprint
- The condition of the escarpment vegetation corridors is slightly to moderately degraded, therefore, its capacity to supply provisioning ecosystem services such as fuel wood, construction wood and inspirational services such as spiritual sites is considered to be moderate to high.



- The Buhuka Flats have extensive settlement areas, which have resulted in over-exploitation of livestock grazing resources available in the open grasslands, grasslands with thickets and open bushland and shrublands that dominate the land cover of the Buhuka Flats. As a result, the condition of these ecosystems is considered heavily degraded, largely due to overgrazing. In addition, carrying capacity calculations indicate that these areas are being grazed by more than twice the amount of cattle than they are capable of.
- However, as stated previously (Section 4.3.1.1 ), not all ecosystem services supply dynamics are affected to the same degree by the condition of the land cover type. In this case, the capacity of degraded grasslands to supply livestock grazing ecosystem services is still considered high; however, ongoing poor grazing management practises are likely to result in ploughing of the ground by cattle’s hooves in wet conditions, and overgrazing in dry conditions, ultimately leading to soil erosion, bush encroachment and exotic species invasion. The Project and associated population influx will likely create additional pressure on these grazing resources. These expected changes in land cover will reduce the capacity of these ecosystems in the Buhuka Flats to supply ecosystem services in the future.

**Pipeline**

- The dominant land cover classes are subsistence farmland and degraded bush; accordingly these will have the greatest proportionate loss to the pipeline servitude footprint
- Although the ‘ecological’ condition of subsistence cropland land cover class is considered poor due to the negative ecological impacts of cultivation, its functionality/ability to supply ES is an effect of its cultivation; therefore, its capacity to supply ecosystem services is considered high as it supports crop production.
- The condition of degraded bushland is heavily-degraded largely as an effect of clearance of bush for cultivation, and probably over-grazing by browsing livestock. Its ability to supply its associated ecosystem services (Table 13) is thus considered Low.

**7.1.2 Which Ecosystem Services could the Project impact?**

The ecosystem services supplied within the Local Study Area are listed according to land cover type in Table 13. These ecosystem services are supplied by ecosystems that will have a loss in area or access as a result of the Project, and thus could potentially be impacted by the Project.

**Table 13: ES supplied within the Project Area of Influence**

Ecosystem Service	Land Cover Type	Definition of Service
<b>Provisioning</b>		
Food	Subsistence cropland	<b>Crops:</b> Subsistence crops associated with individual households Small-scale commercial plantation crops along pipeline route
	Grassland Open bush	Grazing for livestock Cattle and goat herders in the Buhuka Flats are reliant on the availability of grazing resources
	Water (Lake Albert)	<b>Capture Fisheries:</b> Fish is the main source of nutrition for the people in Buhuka Parish





## ECOSYSTEM SERVICES ASSESSMENT

Ecosystem Service	Land Cover Type	Definition of Service
	Degraded bush Open bush Dense bush	<b>Wild foods:</b> Subsistence hunting for bush meat was reported for several villages, mostly along the pipeline route No reports of wild food (fruits, roots) use were recorded during baseline. One study of wild food use in Uganda reported stigma attached to use of wild foods (indicative of poverty and laziness) (Agea, et al., 2011); therefore, actual levels of use of wild foods may have been under-reported.
Biological raw materials	Wetland Woodland Grassland	Reeds and grasses are harvested from wetlands and escarpment grasslands for use as thatching material Papyrus culms may be harvested from wetlands for construction of traditionally-built houses Timber is harvested from trees for use as 'wattle' in construction of mud and wattle houses
	All categories	Extraction of local natural materials (e.g. aggregates) for Project road and camp construction etc.
Biomass Fuel	Degraded bush Open bush Dense bush Woodland	Wood is harvested for firewood and charcoal production
Freshwater	Water Wetland	Freshwater for consumption and irrigation is taken from Lake Albert, boreholes, wells, rivers, streams and swamps throughout the area
Medicinal plants	Degraded bush Open bush Dense bush Woodland Grassland	Numerous medicinal plants are harvested for various purposes within the local study area ( <b>Error! Reference source not found.</b> )
<b>Regulating</b>		
Regulating air quality	Water Wetland	Baseline air quality is generally considered good but may deteriorate periodically as a result of biomass burning – Lake Albert and its associated sedimentary flats and wetlands would have a role as a 'sink' for any atmospheric pollutants
Regulating climate	Woodland	The topography of the escarpment plays a role in the local climate, particularly in rainfall patterns in the Buhuka Flats, which are lower than those on the top of the escarpment
Regulating water flows and timing	Grassland Open bush Wetland	Approximately half (56%) of the soils within the Kingfisher Field Development Area footprint are readily permeable, facilitating aquifer recharge Wetlands in the Buhuka flats retain water and contribute to reduced flooding frequency
Control of erosion	Grassland Open bush Wetland	Vegetation cover within the footprint reduces soil loss and prevents erosion. In areas where storm water infrastructure has been improperly installed, severe erosion was observed
Water purification and waste treatment	Wetland	In the absence of municipal water treatment schemes, wetlands within the Project Area of Influence likely have an important role in partial water purification







Ecosystem Service	Land Cover Type	Definition of Service
Pollination	Subsistence cropland Degraded bush	Beekeeping is practised in several villages along the pipeline route Subsistence agriculture reliant on pollination by bees for fruit and vegetable growth
<b>Cultural</b>		
Recreation and ecotourism	Water (Lake Albert)	Some holiday lodge facilities have been constructed in the vicinity of Lake Albert to cater for tourism thought to be largely driven by staff of the local oil industry
Ethical and spiritual values	Water (Lake Albert, rivers) Wetland Woodland (escarpment)	Local communities of the Buhuka Flats have strong spiritual ties with the lands. The Buhuka Flats area has numerous sacred sites, and has a rich, intangible cultural heritage that includes folklore and taboos related to the Lake, traditional crafts (e.g. making ghee, making bark-cloth), and oral histories
Educational and inspirational	Water Wetland Grassland Open bush Dense bush Woodland	The communities of the Buhuka Flats area have stories and taboos relating to and inspired by the landscapes of the Lake and the escarpment
<b>Supporting</b>		
Habitat	Wetland Water Woodland	Wetland areas are nurseries for juvenile fish Lake Albert provides habitat for fish and waterfowl Wooded areas on the escarpment support terrestrial fauna Forested areas contribute to landscape connectivity for terrestrial fauna through their role as wildlife corridors (Akwetaireho, et al., 2011)
Nutrient cycling	Subsistence cropland	The majority of the pipeline route is considered to have agricultural potential as arable land, due to the productive nature of the soils
Primary production	Grassland Wetland	The Buhuka Flats is an important grazing area as a result of the combination of water availability and productive soils which promote the development of grasslands and wetland vegetation
Water cycling	Grassland Wetland	The Buhuka Flats' hydrological system is different to that beyond the escarpment and plays a role in aquifer recharge. Wetland systems in the Buhuka Flats and on top of the escarpment have importance for water storage and flood attenuation

### 7.1.3 Which Beneficiaries are potentially affected?

The beneficiaries who use those services supplied by the ecosystems that could be affected by the Project (Section 7.1.2) were identified, and fall into two categories:

- The local community – the people that benefit from ecosystem services (Type I) supplied by the land cover types of the area that will be affected by the Project. This consists largely of the rural subsistence population, including subsistence farmers, pastoralists, fishermen, and the unemployed.
- The Project (Type II):



- the Project is dependent on the abstraction of appropriate quantities of fresh water from Lake Albert for make-up water as part of its operation;
- the Project is dependent on the regulation of water quality of Lake Albert; if water quality decreases then treatment costs may be prohibitive;
- the Project is dependent on flood attenuation in the Buhuka Flats due to aquifer recharge and wetland water storage;
- the Project is dependent on a **social licence to operate**.

The demand for services by beneficiaries varies between the different Project-affected areas, largely depending on the dominant land cover type in that area.

### ***Buhuka Flats***

In the Buhuka Flats, demand for services (Type I) arises principally from the following beneficiaries:

- The communities of the lakeshore villages that depend on the Lake Albert fisheries as a primary source of livelihood, income and nutrition.
- The communities of the lakeshore villages that depend on obtaining thatching material from wetland areas on the Buhuka Flats and grassland areas on the escarpment; and construction material for building mud and wattle houses from trees in woodland and bushland both on the escarpment and above it.
- Subsistence farmers resident in the lakeshore villages that use the Buhuka Flats for grazing livestock.
- Migratory herders that bring their cattle to the Buhuka Flats to avail of grazing opportunities there.
- The communities of the lakeshore villages that have strong spiritual attachment to the Lake, the river Malika and the Escarpment.

In addition, ecosystem services are demanded by the Project itself (Type II) - clean fresh water in the necessary quantities and of appropriate quality, which will be abstracted from Lake Albert and used in the oil extraction process during Project operations; as well as the social licence to operate from local communities, who may see their sense of place change as a result of the Project.

### ***Pipeline Route***

Beyond the escarpment, demand for services arises principally from the communities of the villages along the pipeline route who depend on:

- Cultivation of subsistence and small-scale commercial crops as a primary source of livelihood, income and nutrition.
- Raising livestock as a primary source of meat and dairy products.
- Obtaining freshwater from rivers, streams, swamps, wells and boreholes.
- Harvest of timber for wood fuel for cooking, charcoal production and use in home construction.
- Harvest of grass for use in thatching traditionally built homes.
- Hunting of bush meat as a dietary supplement in times of hardship.

## **7.2 Step 2: Ecosystem Service Prioritisation**

Priority ecosystem services related to the Project were identified following an ecosystem service review (Landsberg, et al., 2013). The services were prioritised in two phases, and were aligned with the two types of Priority ecosystem services defined by IFC PS 6.



As *Supporting* ecosystem services have no specific/direct beneficiaries, and impacts to these are captured within the *Provisioning*, *Regulating* and *Cultural* categories for this project, they are not included in the prioritisation exercise.

### 7.2.1 Type I Ecosystem Services: Priority Ecosystem Services according to Project Impact

Ecosystem services were prioritised according to project impact, by answering the three key questions put forward by (Landsberg, et al., 2013):

- 1) Could the Project affect the ability of others to benefit from this ecosystem service?
- 2) Is the ecosystem service important to beneficiaries' livelihoods, health, safety or culture?
- 3) Do beneficiaries have viable alternatives to this ecosystem service?

The results of the prioritisation exercise for Type I ecosystem services are detailed in Appendix B. All Type I ecosystem services and the reasoning behind their determination as Priority ecosystem services (or not) are discussed in the following sections.

#### 7.2.1.1 Provisioning Food – Subsistence Crops

The majority of land cover along the pipeline route consists of subsistence cropland, which is a primary source of livelihood, income and nutrition for the beneficiaries in this part of the Local Study Area. Approximately 1.3% of this land cover will be lost to the footprint of the pipeline and its 30 m servitude (Table 8). The average size of farm currently being cultivated according to the respondents is 3 acres (1.2 ha) and the most common size is 1 acre (0.4 ha). Therefore, average maximum land loss to the pipeline servitude is expected to range from 0.18 ha or 46% of the most common sized farms, to 0.33 ha or 27% of average farms.

Although this will impact the ability of farmers whose land is intercepted by the pipeline and servitude to benefit from this ecosystem service, alternatives may be provided to affected beneficiaries; if this is not possible, then fair and adequate compensation should be offered. The compensation should be in line with IFC Performance Standard 5 and will be addressed in an appropriate Resettlement Action Plan (RAP).

Food from subsistence crops is therefore classed as a non-priority ecosystem service for the purposes of this assessment.

#### Food – Grazing for livestock

Some residents of the Buhuka Flats raise cattle for subsistence purposes, so access to **grazing** areas is considered a priority ecosystem service. The availability of alternative locations to those that may be lost to the Project footprint is uncertain, as carrying capacity for grazing animals in the Buhuka Flats area is already being exceeded by approximately double the amount of livestock present. In addition land tenure in the area is uncertain.

Both residents of the Buhuka Flats and the villages near the pipeline route, and migratory pastoralists, raise **livestock** for subsistence purposes. The extent to which these beneficiaries rely on the grazing opportunities provided by the grasslands in the Project Area of Influence is likely of high importance to their livelihoods and wellbeing. As the availability of alternative sources of grazing is likely to be limited, given the current rates of overstocking, this is considered a **Priority ES**.

#### Food – Capture fisheries

Although **capture fisheries** are not anticipated to be negatively affected by the Project, there is a perception amongst residents of the Buhuka Flats that the Project will affect fish stocks and catch success. Capture fisheries are therefore included as a **Priority ES** for which potential impacts must be assessed.

#### Food – Wild foods



Bush meat hunting and beekeeping were recorded in communities along the pipeline route. The importance of these ES for beneficiaries is uncertain, and the availability of viable alternatives to these ES is unclear, so **wild foods** are included as **Priority ES** based on the precautionary principle.

### **Biological Raw Materials – Construction Products**

The majority of houses in the communities of both the Buhuka Flats and the pipeline route are traditionally built, using grasses from on top of the escarpment as thatching material, mud daub on walls and wood or papyrus culms as construction material for walls. Grasslands, wetlands, woodlands are the primary source of these materials, small areas of which will be lost to the Project footprint in the context of the LSA.

Given the proportionately small amount of expected loss of this land cover type in the context of the Project Area of Influence, there should be viable alternative sources of these raw materials in other unaffected locations. However, increased demand for these materials as a consequence of population influx is unknown. In addition, Project effects on wetland functioning beyond the footprint are uncertain. Supply of biological raw materials is therefore considered a **Priority ecosystem service**.

### **Biomass Fuel**

The majority (approximately 98.9%) of the population in the two Districts use **wood fuel** as the dominant source of energy. In some villages along the pipeline route, a mixture of firewood, banana leaves and grass are used as a source of fuel for cooking. However, firewood has become scarce and many people have resorted to using charcoal which is expensive, and therefore is not considered a viable (affordable) alternative for beneficiaries. This is therefore considered a **Priority ecosystem service**.

### **Freshwater**

The provision of **freshwater** is considered to be a **Priority ES**, due to its importance for beneficiaries throughout the Project Area of Influence, stakeholder perception that the Project may impact the availability and/or quality of freshwater supply, and the lack of viable alternatives to this ES.

### **Medicinal Plants**

Given the proportionately small areas of land take by the Project relative to the available areas within the Local Study Area, it is expected that alternative areas that support medicinal plant species are readily available to beneficiaries. This ES is therefore not considered as priority.

## **7.2.1.2 Regulating**

### **Regulating Air Quality**

Woodland and bushland vegetation of the escarpment may contribute to extraction of atmospheric chemicals (e.g., near roadways), and Lake Albert plays a role as a sink for air emissions of compounds from the burning of fires.

The Project is unlikely to push the regulation of air quality across a sustainability or regulatory threshold, and emissions are expected to be within the standards required by the IFC. This ecosystem service is not considered to be in short supply relative to demand in the Local Study Area, given the baseline of very little industrial or commercial enterprises in the area. Regulation of air quality is therefore not considered to be a priority ecosystem service in terms of Project impact for this assessment.

### **Regulating Climate**

Escarpment vegetation, wetlands of the Buhuka Flats and sediments of Lake Albert within the Local Study Area may contribute to climate regulation through their role as a carbon sink. The loss of the relatively small areas of escarpment vegetation and wetlands to the project footprint, in the context of total available alternative areas in the Project Area of Influence is minimal. However, the effect of increased pressure on these systems due to greater demand for services by the increased population is uncertain. Given the current uncertainty in relation to climate change and possible scenarios, as well as increasing human pressures, how important these habitats will become in the future in terms of climate regulation is uncertain. Indications are that they will increase in importance (Ayebare, et al., 2013), provided human pressures do not overwhelm them. Overall predictions of Project impacts on the ecosystems involved in regulation of





climate range from moderate on escarpment vegetation and wetlands; to no impacts on Lake Albert deep water areas.

The Project is unlikely, however, to push the regulation of climate across a sustainability or regulatory threshold; neither is this service in short supply relative to demand in the LSA, given the minor loss of the ecosystems that supply this ES to the expected Project impacts. Regulation of climate is therefore not considered to be a priority ecosystem service for this assessment.

### ***Regulating Water flows and timing***

The Buhuka Flats has a unique hydrological system which is not fully understood (for example, water supply to 'Luzira' lagoon, aquifer recharge in the Flats from escarpment streams, and wetland water storage capacity). The potential Project impacts on the hydrological system (crossing of drainage lines) are considered to still have a moderately severe impact post-mitigation. Therefore, the Project could affect the ability of others to benefit from this important ecosystem service in the Buhuka Flats area. Viable alternatives to this hydrological system are not evident; therefore regulating water flows and timing is a **Priority ecosystem service**.

### ***Regulating Soil Stability and Erosion Control***

Potential Project impacts on the regulation of soil stability and associated erosion control are considered likely, because clearance of vegetation for construction works in the Buhuka Flats, the escarpment road, and along the pipeline route will increase the vulnerability of soils in these areas to erosion by wind and water. The greatest impacts to soils typically occur during the construction phase. However, mitigation measures and construction management controls are expected to be adhered to during construction of the Project infrastructure, therefore the Project is not expected to impact on this ecosystem service in such a way that the ability of others to benefit from this service would be affected. Therefore this ecosystem service is not considered a priority for this assessment.

### ***Water Purification and Waste Treatment***

Kyangwali sub-county, within which the Project Area of Influence is located, has low safe water coverage (approx. 47%); and many beneficiaries obtain their water supply directly from rivers, streams and swamps. This heightens the importance of the role that wetlands play in the removal of harmful pollutants such as metals and organic materials from surface water systems; this is thus considered a **Priority ecosystem service**.

#### **7.2.1.3 Cultural**

##### ***Recreation and ecotourism***

This ES is not of importance to local beneficiaries, as tourism facilities are non-existent at Lake Albert within the Local Study Area. Although some tourism accommodation facilities (safari lodges) have been developed in the Districts to accommodate low-level tourism by oil workers, there is no evidence that the local communities of the Buhuka Flats or the pipeline route benefit in terms of livelihoods. This ES is thus not considered a priority ES.

##### ***Ethical and spiritual values***

Sacred sites and intangible cultural heritage, evident throughout the Project Area of Influence, are intrinsically linked with natural ecosystems such as wetlands, rivers, lakes and forests. Changes in natural ecosystems arising from Project land take, and changes in the appearance of the landscape due to the visual presence of the Project are likely to affect the ability of local communities to benefit from this ES. This ES is important to beneficiaries as it substantially contributes to their sense of identity. It has thus been identified as a **Priority ecosystem service**.

##### ***Educational and inspirational values***

The Lake Albert and Escarpment landscapes inspire folklore, myths and taboos, thereby contributing to beneficiaries' sense of heritage and identity. Changes in appearance of the landscape are likely to affect the ability of beneficiaries to retain the benefit of this service that is considered important in terms of cultural heritage and identity; in addition access to these areas may be increased (thus impacting the remote quality



of the landscape) or restricted as a result of the Project (e.g., the lagoon near the Kingfisher Field Development Area) therefore this ecosystem service is considered a **Priority ecosystem service** for this assessment.

### 7.2.2 Type II Ecosystem Services – Priority Ecosystem Services according to the Extent of Project Dependence

The outcomes of the prioritisation exercise for Type II ES are detailed in Appendix C. The Type II ecosystems and reasoning behind classification as priority/non-priority ecosystem services are discussed in the following sections.

#### 7.2.2.1 Provisioning Services

##### *Freshwater*

The Project will depend on the abstraction of fresh water from Lake Albert for successful performance (oil extraction), and there are no viable alternatives to water abstraction. There is concern amongst beneficiaries that the Project may impact the quality of water; in addition, the quality of water in Lake Albert is already being pressured by elevated nutrient inputs and associated eutrophication. The Project is reliant on the quality of freshwater resources remaining constant throughout its lifetime, both in order to maintain its social license to operate and in order for operation to remain cost-effective - treatment of eutrophic waters to an acceptable standard for processing use may be cost-prohibitive for the Project.

Although the amount of water required by the Project is considered minimal in the context of the currently available resource, the total demand for Lake Albert's water from other exploration areas and other users has not been defined, and future changes in the lake's water levels over the life of the Project are unknown. Therefore, freshwater provision is considered a Type II **Priority ecosystem service**.

##### *Aggregates for construction*

The Project depends on the extraction of locally-sourced aggregates for construction of facilities, such as camps and access roads, in order to reduce resource costs by using locally available materials, and reduce carbon emissions.

Existing demand for locally extracted aggregate by other beneficiaries is unclear. Local beneficiaries in the Local Study Area currently mostly live in wattle and mud houses but it is unknown how construction practises may change in the future, if economic standards were to improve. The Project has no viable alternative to this ecosystem service; therefore, it is considered a **Priority ecosystem service** in terms of Project dependence.

#### 7.2.2.2 Regulating Services

##### *Regulating air quality*

Air emissions from oil and gas development activities include combustion sources from power and heat generation and use of engines, emissions resulting from flaring and venting of hydrocarbons and fugitive emissions. Stakeholders in the Project Area of Influence perceive that the Project may impact the air quality; therefore, the Project is reliant on the quality of air remaining constant throughout its lifetime in order to maintain the social license to operate.

Although emissions from the Project are expected to be in compliance with the specified standards, and the service is not considered to be in short supply relative to demand in the Local Study Area, given the baseline of very little industrial or commercial enterprises in the area, public perception is that the Project will impact air quality within the Local Study Area. Therefore this is a **Priority ecosystem service** in terms of Project dependence on maintenance of a social license to operate.

##### *Regulating climate*

Flooding of the shores of Lake Albert in 2012 resulted in damage and destruction of homes, and some loss of life. The predicted changes in climatic conditions over the lifetime of the Project could lead to a rise in





Lake Albert's water level, which would result in flooding of the flats, which could affect operational performance.

To counter this risk, the Project infrastructure will be designed to stand at a raised elevation from the actual ground level to avoid potential impacts of flooding over the Buhuka Flats, and will also include additional designed and engineered controls.

These engineering design measures can be considered to be viable alternatives to the climate regulation service provided by ecosystems within the local study area, therefore regulating climate is a non-priority ecosystem service in terms of Project dependence.

### ***Regulation of water timing and flows***

The Buhuka Flats is a zone of aquifer recharge from escarpment streams, and the wetlands in the Buhuka Flats area play a role in water storage. The potential Project impacts on the hydrological system (crossing of drainage lines) are considered to have a moderately severe impact post-mitigation for the construction and operation phases, and this together with ecosystem changes external to the Project (such as increased flooding due to climate change) could potentially change this ecosystem service in a way that would prevent the Project from achieving operational performance.

However, the Project design takes into account such potential impacts, and will put in place appropriate stormwater and flood management engineered measures to prevent the predicted potential impacts taking place. These engineering design measures are viable alternatives to the ecosystem service.

Therefore, regulating water flows and timing is a non-priority ecosystem service in terms of Project dependence.

### ***Regulating soil stability and erosion control***

The predicted rise in the level of Lake Albert over the Project lifetime could lead to an increase in erosion of the shoreline, thereby reducing the width of the Flats and increasing flood likelihood; therefore, the Project depends upon continued supply of this ES for its operational performance.

Engineered measures for the control of erosion arising from vegetation removal are considered sufficient to minimise the impacts of vegetation clearance. The Project, in compliance with the requirements of IFC Performance Standards 1 and 3, has undertaken predictive modelling to ensure that the Project's operational performance will not be put at risk by rises in Lake Albert's level. Therefore, regulating soil stability and erosion control is not considered to be a Priority ecosystem service according to operational risk to Project Performance.

### ***Water Purification and Waste Treatment***

The Project is reliant on the availability of fresh water of a certain quality standard from Lake Albert, both for use as drinking water for Project staff, and for Project oil and gas activities and processes. The role that wetland systems and the lake itself play in water purification through nutrient assimilation may change over the lifetime of the Project, both in terms of wetland and lake ecosystem extent and condition which could potentially decrease in this time, due to existing drivers of change.

However, the Project incorporates a waste water treatment facility, which could be a viable alternative for this ecosystem service, should abstracted fresh water need to be treated to achieve appropriate standards prior to use. In addition, the Project's water management specification commits the Project to discharge waters that are treated to acceptable environmental standards. Water purification and waste treatment is thus not considered a priority ecosystem service in terms of Project dependence.

### **7.2.2.3 Cultural *Ethical and spiritual values***

The Project may depend on the availability of this ES remaining constant throughout its lifetime, in order to maintain its social license to operate. However, the presence of the Project in the landscape may affect beneficiaries' affinity to sacred sites for example, which are most frequently associated with areas of natural



beauty, and that also have less obvious qualities, such as being remote or isolated or quiet. Maintenance of such features and their associated ecosystem services is considered significant in order to maintain the Project's social license to operate, so these are classified as Type II **Priority ecosystem service**.

### *Educational and inspirational*

As with the provision of ethical and spiritual values, the Project may depend on the availability of this ecosystem service remaining constant throughout its lifetime, in order to maintain its social license to operate. Similarly, the actual physical presence of the Project may reduce the inspirational value of the Lake Albert and Escarpment landscapes; and Project provision of an alternative inspirational resource would likely prove impossible. This is therefore considered a **Priority ecosystem service**.

## 8.0 PROJECT AREA OF INFLUENCE FOR PRIORITY ECOSYSTEM SERVICES

- The Project Area of Influence for Priority Ecosystem Services was set, based on mapping the land cover types that supply Priority Ecosystem Services against the Local Study Area, within which the beneficiaries of those ecosystem services are encompassed. The Project Area of Influence for Priority Ecosystem Services is illustrated in Figure 20. It is in this context that impacts on priority ecosystem services are assessed.





# ECOSYSTEM SERVICES ASSESSMENT




Figure 20: Project Area of Influence for Priority Ecosystem Services (includes Lake Albert – see inset)



### 9.0 ASSESSMENT OF PROJECT IMPACT ON PRIORITY ECOSYSTEM SERVICES

The following sections summarise expected impacts on Priority ecosystem services. Three of the four identified Type II ecosystem services overlap with Type I ecosystem services; in these cases, they are discussed under the same heading.

The impact assessment is not separated into construction/operation/decommissioning phases, as the ecosystem services are generally tied to land cover types and associated loss to the Project footprint (especially provisioning and regulating ecosystem services), or the presence of the Project in the landscape (cultural ecosystem services), which will be in effect for the lifetime of the Project. However, where potential impacts on ecosystem services are considered specific to a particular Project phase (for example, regulation of air quality is more likely to be affected during the operational phase of the Project), this is stated at the outset.

The impact severity ratings presented in Table 14, Table 15 and Table 16 are based on the anticipated impacts on ecosystem services, before specific mitigation measures have been applied. Specific mitigation measures relating to ecosystem services set out by relevant specialist studies are discussed in Section 0. In cases where the specialist studies do not address mitigation of impacts on ecosystem services (that is, food and biofuel provisioning ecosystem services), or where residual impacts on ecosystem services remain following application of specialist recommendations, additional mitigation measures to address such impacts are also provided in Section 0 Mitigation Measures.

#### 9.1 Food Provisioning

The potential impacts on food provisioning within the Project Area of Influence will extend throughout the construction, operation and decommissioning phases in the Project, due to the presence of the Project and associated loss of land cover to its footprint, and the increased demand on food supply that is expected to occur in tandem with population influx, once the Project commences construction and operation. The potential impacts will also extend throughout the lifetime of the Project along the Pipeline route, because the servitude will remain in place for maintenance and access to the pipeline itself.

The impacts on all Provisioning ecosystem services are discussed in the following Sections and summarised in Table 14.

##### 9.1.1 Grazing for livestock

Potential Project impacts on this ecosystem service are related to the loss of available area for subsistence farming, and increased loss of cropland associated with increased risk of soil erosion in areas cleared (particularly along the pipeline servitude) grazing livestock. Grassland areas that are currently used for grazing livestock will be reduced in extent as a result of land-take for the Project footprint in the Buhuka Flats, and will be further pressured by population influx to the Buhuka Flats and concomitant increased demand for livestock grazing resources.

The loss of this land constitutes a more significant impact in comparison to the loss of cropped land, because grazing pressure in the Buhuka Flats is severe and the loss of areas of grassland will place more pressure on the remaining grazing lands. Pastoralists travel to the Buhuka Flats specifically to graze their livestock, indicating a paucity of alternative grazing lands in the region. Population influx once the Project commences construction, and operation activity may further contribute to increased grazing pressure in the Buhuka Flats. In addition, land tenure in the Project Area of Influence is not always well defined, which may affect the success of compensation schemes for beneficiaries.

The intensity of potential Project impacts on supply of grazing for livestock was considered to be **high**, due to the current level of overgrazing in the Buhuka Flats, and its potential compoundment by loss of grazing lands to the Project footprint, and the Project population influx, which is expected to increase pressure on the availability of grazing for livestock. The geographic extent of the impact may reach to beneficiaries beyond the footprint, such as pastoralist herders who travel to the Flats specifically to access grazing lands.





The sensitivity of this ecosystem service is considered **high**, due to the apparent lack of alternative grazing resources elsewhere in the region. As with subsistence cropping, land tenure in the area is not always clear, and there are beneficiaries (for example, the Balalo pastoralist communities) grazing livestock in the Buhuka Flats that do not have the right land tenure, and therefore may be difficult to compensate for their loss.

The potential impact severity on this ecosystem service is thus considered **high**. With the application of the recommended mitigation measures, the intensity of impact on this ecosystem service will be reduced; with appropriate resettlement, compensation and community development planning the sensitivity of the ecosystem service to impact can also be reduced, resulting in an impact of **moderate** severity, post-mitigation.

### 9.1.2 Capture Fisheries

The Project itself, in terms of its footprint and operations, is expected to have minor impacts on the near-shore habitats of Lake Albert post-mitigation. The near-shore aquatic habitat is an important fish breeding area, therefore it can be projected that construction of the Project and its operation will have minor impacts on the Lake Albert fishery. However, the majority of beneficiaries in the Buhuka Flats rely on Lake Albert fishery as a sole source of livelihood, income and food, and there is concern amongst beneficiaries that the Project may impact fish stocks or the size of the fish catch. In addition, population influx associated with the Project and increased easy access to the lakeshore via the proposed escarpment road will place additional strain on the supply of natural resources and may contribute to over-fishing in the locality.

The intensity of potential Project impacts on this ecosystem service is, therefore, considered **high**, largely due to population influx. Population influx is expected to contribute to a negative feedback loop that will contribute to overfishing - as the villages on top of the escarpment become more accessible to fish sellers as a result of improved access via the escarpment road, more favourable conditions for trading fish on the escarpment (instead of at boat-accessible locations) are expected to develop, resulting in stimulation of economic growth in the Project Area of Influence – which in turn is expected to attract more people to the Project Area of influence – which may ultimately have dire consequences on the long-term viability of the Lake Albert fishery. These positive and negative socio-economic impacts will be addressed by the livelihood management plan. The effect of population influx would likely occur from construction through to the medium-term, that is, the lifetime of the project operations (25 years).

The sensitivity of the ES is **high**, as it is not substitutable or replaceable – beneficiaries are heavily dependent on this ES, and have no obvious alternate livelihoods, incomes or even food sources. The severity of the potential Project impact on the supply of this ES is thus considered **Major**. Following the application of the recommended mitigation measures, including influx management planning and provision of mess facilities, the intensity of impacts is reduced to Low, leaving an impact of **moderate** severity, post-mitigation.

### 9.1.3 Wild foods

Bush meat hunting (specifically hunting for rats) was recorded in several villages near the pipeline routes. Bush meat hunting is a known pressure in Bugoma Central Forest Reserve in the Project Area of influence. The value of this ecosystem service to beneficiaries within the Project Area of Influence is unclear from the baseline data, however, the literature suggests that bush meat hunting may form an important 'bridge' or 'safety-net' resource for beneficiaries, particularly during times when food is scarce e.g. during times of unemployment, crop failure or drought (Akwetaireho, et al., 2011), (CRA, 2006).

Reductions in land cover types (woodland, dense and open bush) that support the supply of this ecosystem service due to Project impact will negatively affect the supply. In addition, the construction of the escarpment road and upgrade of the Hoima-to-Ikamiro Road will increase and enhance ease of access to the escarpment vegetation and Bugoma Central Forest Reserve, which, together with the expected population influx to the area, is expected to significantly increase demand for these ecosystem services. This could therefore likely affect the condition of the ecosystems and their capacity to supply ecosystem services. However, the demand for bush meat within the Project Area of Influence is uncertain. Bush meat is not thought to be a major source of food within the Project Area of Influence. A **medium**-intensity Project impact on this ecosystem service is expected for beneficiaries of this ES within the Project Area of Influence.



The sensitivity of this ES is ranked as medium – the supply of bush meat that would be affected by Project land take and population influx may not be easily substitutable. As mentioned earlier declines in poaching in Bugoma Forest Reserve were attributed to the lack of animals left to hunt, rather than changes in hunters' attitudes.

The severity of potential Project impacts on the supply of this ES is thus considered **moderate**. These impacts will still be of moderate severity following mitigation measures, but are predicted to occur at a lower intensity as a result of promotion of sustainable farming and other activities that provide alternative food sources, and community education programmes.

## 9.2 Biological Raw Materials

### *Construction materials for traditionally built houses*

Almost all beneficiaries in the Local Study Area (approx. 98%) live in houses constructed with mud-and-wattle walls with earth floors; a very small percentage live in houses built with fired bricks. The majority have thatched roofs (approx. 76%), with the remainder having iron sheet roofs. As a sub-set of the Local Study Area, this pattern is also expected in the Project Area of Influence.

Thatching materials are sourced from grasslands on the escarpment, and wattle is timber harvested from trees in woodland and bushland areas. Increases in the population in the Project Area of Influence is expected to increase demand for these raw construction materials, which could affect their supply; both in terms of immediate availability, and in the long-term, degradation of the ecosystems that supply these ecosystem services may also contribute to reduced supply capacity. At present however, there is no baseline evidence that suggests that availability of these resources is under pressure. The intensity of potential Project effects on the supply of raw materials for traditional house construction is thus considered **medium**.

Alternative housing materials (fired bricks and iron roof sheeting) are available for purchase in the Project Area of Influence; however, purchase of these materials is probably not a viable alternative to harvesting grass and wattle timber, which is free. Therefore, the sensitivity of the ecosystem service is **high**.

The severity of potential Project impacts on the supply of this ecosystem service is considered **major**. Provision of housing for workers employed by the Project during construction and operation may alleviate the demand for these materials in the Project Area of Influence, which would reduce the intensity of the impact, however the sensitivity of the ecosystem service remains unchanged as population influx will also include people who do not work at the Project, for whom the ecosystem service will not be substitutable. Post-mitigation impacts are thus expected to be of **moderate** severity.

### *Natural aggregates for Project facility construction*

The scale of extraction of aggregates for Project facility construction is unclear. It is not expected that the Project will affect the ability of other beneficiaries to benefit from this ecosystem service, as other beneficiaries within the Project Area of Influence do not typically rely on aggregate extraction as a priority/non-priority ecosystem service; however, construction practises may change over the lifetime of the Project, and local beneficiaries may come to depend on this ecosystem service. Based on current evidence, the intensity of potential Project effects on the availability of natural aggregates is considered **medium**.

The Project is reliant on the availability of locally-sourced aggregates to maximise cost efficiency and carbon footprint reduction; however, alternatives in the form of imported resources are expected to be available to the Project. The sensitivity of the ecosystem service is thus considered **medium**.

The severity of potential Project impacts on the supply of this ecosystem service is considered **moderate**. Avoiding extraction of aggregate in areas important for ecosystem service supply, especially sites of cultural heritage importance, and Project development of a procurement strategy for using locally-sourced aggregates, reduces the intensity of impact and sensitivity of receptors, reducing potential impacts on this ecosystem service to **minor**.



### 9.3 Biomass Fuel

The majority of the population in Kikuube and Hoima district and thus the Project Area of Influence use wood fuel as their main source of energy. However, firewood has become scarce and many beneficiaries have resorted to using charcoal, which is expensive. In addition, charcoal is derived from timber that is also harvested from woodland and bush areas.

Reductions in land cover types (woodland, degraded bush, open bush, dense bush) that supply this ecosystem services due to Project impact may negatively affect the supply of this ecosystem services. In addition, population influx to the area associated with the Project is expected to significantly increase demand for firewood and charcoal, which will have a direct impact on the condition of the woodlands and bush, and thus their capacity to continue to supply timber for use as fuel. The intensity of potential Project effects on the supply of biomass fuel is thus considered **high**, as the ecosystems and their capacity to supply firewood may be altered to the extent that supply will temporarily or permanently cease.

The sensitivity of the ecosystem service is **high**, because there are almost no viable alternatives to the use of firewood and charcoal by beneficiaries, it is not easily replaceable – although non-wood products, such as banana leaves and grass, are used in some parts as a source of fuel for cooking to supplement firewood, such fuels could not replace firewood in terms of burning time and heat generation.

The severity of potential Project impacts on the supply of this ecosystem service is considered **major**. If the recommended mitigation measure of CNOOC, that is, investigating the feasibility of provision of cheap gas to local communities is enacted, the impact would be reduced in intensity and the ecosystem service sensitivity would be lowered due to the availability of an affordable substitute to charcoal, resulting in **minor** impacts, post-mitigation.

### 9.4 Freshwater

Freshwater falls under both Type I and Type II priority ES – the Project may impact the supply of this ES and the Project is also dependent on both the quantity and the quality of supply of this ES.

#### *Freshwater as a Type I Priority ES*

The Project footprint may impact the supply of Freshwater for beneficiaries, particularly in the vicinity of areas where the infrastructure will intercept drainage lines, streams, rivers and/or swamps. Pressure from increased populations in the Project Area of Influence may contribute to increased nutrient load and eutrophication of the lake, with concomitant effects on water quality. Although unlikely, an accidental spill of oil or process water in Lake Albert cannot be discounted as a potential impact on the quality of the freshwater supply. Impacts on the quality and quantity of water supply may ensue.

The intensity or magnitude of potential effects could extend throughout the Project Area of Influence and beyond, but would most likely be short-term in duration. Potential impacts on water quality would most likely occur during the construction phase due to potential sediment release during earthworks activities, and may also occur during the medium-term operational life of the Project, in the event that inflow of untreated human waste and agricultural runoff continues and will increase due to population influx. Impacts on quantity of freshwater supply are considered unlikely, given the amount of water being abstracted from Lake Albert for the Project operations in the context of the available surface water resource. Nonetheless, should water quality be impacted negatively, for example, by eutrophication or an industrial accidental leakage, the intensity of impacts could be **high**.

The sensitivity of the ecosystem service is **high**, as freshwater supply in the necessary quantities and to the required quality standards is not easily substitutable. In addition impacts on Lake Albert water quality from potential eutrophication would also directly affect the fish population and thus the capacity to support capture fisheries, which is also a highly sensitive ecosystem service. The severity of potential Project Impacts on the supply of this ecosystem service is thus considered **major**. The intensity of impacts can be reduced by appropriate water management and monitoring mechanisms and influx management planning, reducing the predicted impacts post-mitigation to **moderate** severity.





### *Freshwater as a Type II Priority ES*

The Project is dependent on the supply of Freshwater from Lake Albert for operational performance, as water use is a necessary part of the oil extraction process. Potential impacts on the quantity of this ES may arise as a result of changing water levels in Lake Albert, and the cumulative effects of other oil development areas also abstracting water from Lake Albert for the same purpose. Water quality deterioration may occur in Lake Albert as a result of eutrophication from nutrient loading of streams and rivers flowing into the lake. The Project is dependent on the lake's capacity to assimilate this waste loading to avoid the financial implications of having to treat intake water to an appropriate standard for use in operations.

The intensity of potential Project effects on this ecosystem service is considered **medium**. The amount of water demanded by the Project in terms of the available water resource is negligible, and will be even more so if as predicted the water level in Lake Albert rises due to predicted effects of climate change. However, ongoing nutrient-loading to the lake is likely to result in water quality deterioration, which may require that the Project treat abstracted water before it can be used. However, if necessary, treatment of intake water should be a relatively inexpensive process for the Project.

The sensitivity of the ES is **high**, as the ES is not viably substitutable for the Project. The severity of potential impacts on this ES in terms of Project demand is thus considered **major**. Reducing volumes of water intake through treatment and recycling, and participation in water catchment management activities in the Lake Albert basin can reduce the Project impact intensity to negligible, reducing overall impacts on this ecosystem service post-mitigation to **minor**.





**Table 14: Impacts on Provisioning ecosystem service within the Project Area of Influence**



## ECOSYSTEM SERVICES ASSESSMENT

Ecosystem Service	Potential Impacts	Pre-mitigation			Post-mitigation		
		Intensity	Sensitivity	Severity	Intensity	Sensitivity	Severity
Grazing for Livestock	<ul style="list-style-type: none"> <li>Loss of grassland to Kingfisher Field Development area</li> <li>Increased erosion potential of grassland adjacent to cleared areas</li> <li>Population influx-associated pressures on already over-utilised resource</li> <li>Land tenure not clearly defined – there are pastoralist herders present on the Buhuka Flats who do not have land tenure and thus would not be compensated for loss</li> </ul>	High - 4	High - 4	Major - 16	Medium - 3	Medium - 3	Moderate - 6
Capture Fisheries	<ul style="list-style-type: none"> <li>Perception amongst beneficiaries that Project may impact fish stocks/fish catch</li> <li>Minor impacts on near-shore aquatic environment expected, which may affect fish breeding</li> <li>Population influx will increase demand on Lake Albert fisheries</li> </ul>	High - 4	High - 4	Major - 16	Low - 2	High - 4	Moderate - 8
Wild Foods	<ul style="list-style-type: none"> <li>Bush meat hunting takes place in some villages along the pipeline route and in Bugoma Central Forest Reserve;</li> <li>Woodland and bush that supports hunted species will be reduced in area by pipeline servitude land-take, and will be made more accessible to hunters by upgrade of road and clearance of pipeline servitude</li> <li>Population influx will increase demand for bush meat</li> </ul>	Medium - 3	Medium - 3	Moderate - 9	Low - 2	Medium - 3	Moderate - 6
Construction materials for traditionally-built houses	<ul style="list-style-type: none"> <li>Reductions in land cover types (grassland, wetland, woodland and bushland) that supply this ES due to Project land-take</li> <li>Population influx may increase demand for housing construction materials</li> </ul>	Medium - 3	High - 4	Major - 12	Low - 2	Medium - 3	Moderate - 6
Natural aggregates for Project facility construction	<ul style="list-style-type: none"> <li>Reductions in land cover types that supply this and other ES due to extraction of aggregates</li> <li>Local shortages in aggregates could lead to higher costs for other beneficiaries if importation becomes necessary</li> </ul>	Medium - 3	Medium - 3	Moderate - 9	Negligible - 1	Low - 2	Minor - 2





## ECOSYSTEM SERVICES ASSESSMENT

Biomass Fuel	<ul style="list-style-type: none"> <li>Reductions in land cover types that supply this ES due to Project land-take, particularly along pipeline route and escarpment road</li> <li>Population influx expected to increase demand for firewood and charcoal</li> </ul>	High - 4	High - 4	Major - 16	Low - 2	Low - 2	Minor - 4
Fresh Water (Type I)	<ul style="list-style-type: none"> <li>Impacts on quantity and quality of water supply where Project infrastructure intercepts drainage lines, streams, rivers and/or swamps</li> <li>Impacts on water quality of Lake Albert should an accidental spill occur</li> <li>Impact on water quality due to increased nutrient loading from population influx</li> </ul>	High - 4	High - 4	Major - 16	Low - 2	High - 4	Moderate - 6
Fresh Water (Type II)	<ul style="list-style-type: none"> <li>Quantity of fresh water supply to the Project may be affected by climate change and change in Lake Albert levels</li> <li>Cumulative effect of other oil development areas also abstracting water from Lake Albert</li> <li>Water quality deterioration may necessitate treatment of water being used in Project activities, increasing Project operation costs and affecting operational performance</li> </ul>	Medium - 3	High - 4	Major - 12	Negligible - 1	High - 4	Minor - 4





### 9.5 Regulating Air Quality

This is a Type II priority ecosystem service – the Project is also dependent on the supply of this ES in order to maintain its social licence to operate, and maintain operational performance by not triggering regulatory responses, for example, by exceeding emission guidelines.

#### *Regulation of Air quality as a Type II Priority Ecosystem Service*

The Project depends on the ongoing provision of this ecosystem service, as although Project emissions will be within the recommended limits set out in the relevant guidelines, there will still be some level of emissions by the Project to the air. Stakeholders perceive that the Project may affect air quality, so any changes in air quality (though not necessarily caused by the Project itself) might be attributed by beneficiaries to the Project, thereby affecting its social license to operate.

The principle ecosystem delivering air quality regulation services within the Project Area of Influence is Lake Albert's sediments and its associated flats and wetlands, which would have a role as a 'sink' for any atmospheric pollutants, as well as carbon sequestration. The escarpment vegetation and forest areas, particularly Bugoma CFR, grasslands and bushlands are also expected contribute to air quality regulation.

The severity of Project impacts to the Lake Albert and wetland ecosystems providing this service is expected to be minor following mitigation; however moderate-major effects remain on the escarpment vegetation and Bugoma CFR respectively, after mitigation. The intensity of potential Project impact on the provision of this ecosystem service is considered to be **medium**, given the limited amount of loss of escarpment vegetation and woodland area, and the minor effects anticipated on Lake Albert and wetlands post-mitigation.

The sensitivity of the ecosystem service is **high**, as although measures can be enacted to mitigate direct Project impact on this ecosystem service, the perception of local stakeholders may remain unchanged. The severity of potential future impacts on the Project is **major**, as a result of its dependence on this ES for operational performance and maintenance of its social license to operate.

Following the application of recommended mitigation measures, particularly community education programmes and corporate social responsibility initiatives, the negative public perception of potential Project impacts on air quality and thus the sensitivity rating for this ecosystem service can be reduced to low; and intensity to negligible,

### 9.6 Regulating Water Flows and Timing

The effects of placing Project infrastructure within and intercepting wetlands, rivers, streams and drainage lines will both reduce the surface area of these land cover types, reducing their ability to regulate water flows, and alter their hydrological properties (e.g., subsurface flow through soils) and ecological integrity, which may affect their capacity to regulate water flows.

The intensity of potential Project construction impacts on wetlands and drainage lines that supply this ecosystem service in the Project Area of Influence is expected to be **medium**, as although the wetlands will be altered, natural processes are expected to continue in impacted wetlands, albeit in a modified way.

The sensitivity of the ecosystem service is considered **medium**, as the proposed mitigation measures for wetlands that are directly affected by the Project should ensure that the provision of the ecosystem service will be maintained over the lifetime of the Project. A potential Project impact of **moderate** severity on the supply of this ecosystem service is predicted.

The application of the recommended mitigation measures, particularly the incorporation of engineered design features to ensure that water flows in impacted wetland systems and sub-surface flows are maintained, will reduce the extent of any potential impacts and limit their duration, however the sensitivity of the ecosystem service will remain medium; a potential Project impact of **moderate** severity is predicted for this ecosystem service, post-mitigation.





### 9.7 Water Purification and Waste Treatment

Many beneficiaries within the Project Area of Influence obtain their drinking water directly from sources including Lake Albert, rivers, streams, wetlands and swamps. The role that ecosystems such as wetlands play in the removal of harmful pollutants such as metals and organic materials from surface water systems is important in the context of the lack of formal water treatment systems. Lake Albert also plays a role in the assimilation of nutrients in surface water systems associated with the lack of human sanitation facilities in the region.

The amount of wetland cover being directly lost to the footprint is minimal in the context of the available resource in the Project Area of Influence. However, where wetlands are being intersected by linear infrastructure such as roads and the pipeline route, there is a potential for the downstream wetland habitat to be affected if proper management controls are not implemented, particularly during construction. Even with appropriate measures in place, erosion of wetlands is expected to take place downstream of pipeline and road crossings, and flooding upstream of crossings. Indirect Project impacts may put pressure on Lake Albert's capacity to deliver this ecosystem service, namely the presence of approximately 22,000 people on the Buhuka Flats and other nearby villages who do not currently have access to running water and sanitation. The effect of which is currently unknown but may extend regionally, should the water quality of Lake Albert and its capacity to supply water purification services be impacted. The overall potential Project impact intensity on this ecosystem service is thus expected to be **high**.

The sensitivity of the ecosystem service to Project impact is **medium**, as although additional pressures on the nutrient loading of Lake Albert from the expected population influx cannot be readily predicted; the ecosystem service is substitutable with the development of appropriate water treatment and sanitation facilities, which will be addressed by the Influx Management Plan for the Project. The overall severity of potential Project impacts on this ecosystem service is thus considered **high**.

The incorporation of appropriate wastewater treatment and suitable sanitation facilities at the Project, as well as the Influx management plan will reduce the intensity of Project impact on this ecosystem service where it is provided by Lake Albert. Appropriate mitigation to maintain wetland functioning in areas that will be intersected by the pipeline will also contribute to reduced impact intensity. However the sensitivity of the ecosystem service remains the same, therefore a post-mitigation impact of **moderate** severity is predicted.



## ECOSYSTEM SERVICES ASSESSMENT

**Table 15: Impacts on Regulating ecosystem services**

Ecosystem Service	Potential Project Impacts	Pre-mitigation			Post-mitigation		
		Intensity	Sensitivity	Severity	Intensity	Sensitivity	Severity
Regulating Air Quality (Type II)	Loss and degradation impacts on wetlands and vegetation of escarpment and Bugoma Central Forest reserve may reduce the capacity of these ecosystems to supply this ES Reduction in supply of this ES may reduce social license to operate due to beneficiary perception that the Project is the cause, despite application of mitigation measures	Medium - 3	High - 4	Major - 12	Negligible - 1	Low - 2	Minor - 2
Regulating Water Flows and Timing	Placing Project infrastructure within and intercepting wetlands, rivers, streams and drainage lines will both reduce the surface area of these land cover types, reducing their ability to regulate water flows	Medium - 3	Medium - 3	Moderate - 9	Low - 2	Medium - 3	Moderate - 6
Water Purification and Waste Treatment	The role that wetlands play in water purification and waste treatment, in the context of limited formal water treatment systems is likely significant Amount of wetland land cover being directly lost to the footprint is minimal in the context of the available resource in the Project Area of Influence, however indirect impacts on wetland functioning that may occur Indirect Project impacts on Lake Albert nutrient assimilation capacity due to population influx and lack of sanitation facilities	High - 4	Medium - 3	Major - 12	Moderate - 2	Medium - 3	Moderate - 6





### 9.8 Ethical and Spiritual Values; Educational and Inspirational Values

These ES are considered together given that they are rooted in the same cultural landscapes and are potentially affected and demanded by the Project in the same ways. These ecosystem services may be impacted by the Project, and the Project also relies on the maintenance of the supply of these ecosystem services in order to prevent potential impacts on its social licence to operate.

#### *Ethical and Spiritual Values, and Inspirational Values as a Type I Priority ES*

Sacred sites and intangible cultural heritage are inextricably linked with the landscapes and natural ecosystems of Lake Albert and the escarpment, and are important in terms of beneficiaries' sense of identity and heritage. The Project will impact these ecosystem services due to changes as a result of loss of areas of natural ecosystems, and the visual presence of the Project itself in these landscapes; both of which are expected to limit the beneficiaries' capability to benefit from this ecosystem services.

The potential Project impact on the supply of these ecosystem services will have a **medium** intensity, as the effect will likely extend to beneficiaries in the Project Area of Influence, and will last for at least the duration of the Project and probably longer than that – it is probable that even if the Project footprint is rehabilitated post-decommissioning, the escarpment road will remain, and the landscape of Lake Albert and the escarpment will have changed irreversibly, and associated intangible cultural heritage such as oral histories of places though expected to continue, will become modified in future generations.

The sensitivity of these ecosystem services is **high** as they are irreplaceable, based as they are on the Lake Albert and escarpment landscapes as they stand and have stood for generations. The overall severity of the potential Project impact on these ecosystem services is thus considered **major**.

The application of mitigation measures can reduce the intensity of Project impacts. However, the sensitivity of the ecosystem service remains high as it is essentially irreplaceable. A post-mitigation impact of **moderate** severity is predicted.

#### *Ethical and Spiritual Values, and Inspirational Values as a Type II Priority ES*

The Project relies on the continued supply of these ecosystem services to maintain its social licence to operate, granted by the local community who gains most from these ecosystem services.

The intensity of potential effects on the Project due to its reliance on the continued supply of this ES to maintain its social license to operate are **medium** – the Project's operational performance could be moderately affected if beneficiary disaffection and social unrest due to loss of integrity of sacred sites begins to impact the Project's social licence to operate.

The sensitivity of the Project to changes in this ES is **high**, as the supply of this ES by the culturally significant landscapes of Lake Albert and the escarpment are not substitutable. The overall severity of potential impacts on the Project due to its dependence on this ES is therefore **major**.

The application of mitigation measures can reduce the intensity of Project impacts. However, the sensitivity of the ecosystem service remains high as it is essentially irreplaceable. A post-mitigation impact of **moderate** severity is predicted.





**Table 16: Impacts on Cultural ecosystem services**

Ethical and spiritual values, and Inspirational values	Potential Impacts	Pre-mitigation			Post-mitigation		
		Intensity	Sensitivity	Severity	Intensity	Sensitivity	Severity
Type I	<ul style="list-style-type: none"> <li>The Project will impact these ES due to changes as a result of loss of areas of natural ecosystems, and the visual presence of the Project itself in these landscapes</li> </ul>	Medium - 3	High - 4	Major - 12	Low -2	High - 4	Moderate - 8
Type II	<ul style="list-style-type: none"> <li>Potential reduction in the value of this ES for beneficiaries may adversely affect the Project's social license to operate</li> </ul>	Medium - 3	High - 4	Major - 12	Low -2	High - 4	Moderate - 8





## ECOSYSTEM SERVICES ASSESSMENT

### 10.0 RECOMMENDED MITIGATION AND MONITORING MEASURES

Mitigation measures provided in the following sections include those from specialist studies that are specific to potential impacts on the supply of ecosystem services, and suggested additional mitigation measures based on the guidance provided by IPIECA/OGP for oil and gas project impacts and dependencies on ES (IPIECA, 2011). The recommended mitigation measures are presented in Table 17.

**Table 17: Mitigation measures for impacts on Priority Ecosystem Services**

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
<p>■ <b>Food Provision – Grazing for Livestock</b></p>				
<p>Economic displacement experienced by impacted herding communities should be addressed in terms of the IFC Performance Standard 5 discussing the involuntary resettlement and compensation practices in project-affected communities through development of an appropriate Resettlement Action Plan (RAP).</p> <p>The RAP may require a specialist livestock assessment and management component to address impacts to livestock.</p>	<p>■ As stipulated in RAP</p>	-	CNOOC	-
<p>A livelihood restoration plan should be developed to specifically formulate mitigation strategies for the loss of grazing land</p>	<p>■ As stipulated in Livelihood Restoration Plan</p>	-		
<p>Project support of sub-county administration strategies to solve regional farming difficulties such as crop failure due to disease and drought (e.g. introduction of modern farming methods, training farmers in post-harvest techniques, and sensitising farmers about land degradation) as part of the Community Development Plan/Corporate Social Responsibility initiatives</p>	<p>■ As per community development plan.</p>	-	CNOOC	-
<p>The Project could support the local economy by sourcing food locally, where feasible</p>	<p>■ As per community development plan.</p>	-	CNOOC	-





## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
<b>■ Food Provision – Capture Fisheries</b>				
An influx management plan will be developed to address appropriate measures to mitigate the expected Project-associated in-migration effects on capture fisheries	■ Monitoring measures described in the plan	As required	CNOOC	-
Enforcement of a complete ban on wildlife harvesting (hunting/ trapping/ fishing) for all Project personnel	■ No personnel and/or contractors allowed beyond footprint of Project	As required	CNOOC and Contractors	■ Inductions for all staff
Inclusion of a construction camp with mess facilities for locally-hired staff	■ -	-	CNOOC	■ -
<b>■ Food Provision – Wild Foods</b>				
Supporting local communities in developing sustainable farming, ecotourism or other activities that provide alternative food sources and income	■ As per Livelihood Restoration Plan	-	CNOOC	■ -
Support scientific studies and monitoring programs aimed at assessing the sustainability of using local resources, as part of Corporate Social Responsibility initiatives	■ -	-	CNOOC	■ -
Enforcement of a complete ban on wildlife harvesting (hunting/ trapping/ fishing) for all project personnel	■ No personnel and/or contractors allowed beyond footprint of Project	-	CNOOC and Contractors	■ Inductions for all staff
Worker and community education programme focussing on the impacts and risks of bush meat hunting (e.g. disease) to be incorporated into the Community Development Plan	■ As per Community Development Plan	-	CNOOC	■ -







## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Inclusion of a construction camp with mess facilities for workers	■ -	-	CNOOC	■ -
<b>■ Biological Raw Materials – construction material for traditional houses</b>				
Inclusion of a construction camp with accommodation facilities for workers in Project plan	■ -	-	CNOOC	■
Support scientific studies and monitoring programs aimed at assessing the sustainability of using local resources for home construction	■ -	-	CNOOC	■ -
<b>■ Biological Raw Materials – aggregates for construction of Project facilities</b>				
Avoid aggregate extraction in areas of natural habitat or in the vicinity of sites of cultural heritage importance; target aggregate extraction for areas already in degraded state such as subsistence cropland within the Project footprint	■ -	-	CNOOC	■ -
Develop a procurement strategy that encourages use of locally-sourced aggregates, but that involves mechanisms for assessing or maintaining the sustainability of the supply.	■ -	-	CNOOC and Contractors	■ -
<b>■ Biomass Fuel – fire wood and charcoal</b>				
Supply of cheap alternatives (e.g. gas) to local markets by CNOOC to be investigated	■ -	-	CNOOC	■ -
Support scientific studies and monitoring programs aimed at assessing the sustainability of using commercially-planted forms of biomass fuel, such as Jatropha	■ As per Livelihood Restoration Plan	-	CNOOC	■ -





## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Enforcement of a complete ban on harvesting of fire wood at for all project personnel	<ul style="list-style-type: none"> <li>■ No personnel and/or contractors allowed beyond footprint of Project</li> </ul>		CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Inductions for all staff</li> </ul>
<b>■ Fresh Water (Type I)</b>				
Reduce water volumes needed by Project activities through treatment and re-use of process water and waste water	<ul style="list-style-type: none"> <li>■</li> </ul>			<ul style="list-style-type: none"> <li>■ -</li> </ul>
Implement appropriate water pollution control measures such as oil interceptors, treatment of wastewaters	<ul style="list-style-type: none"> <li>■ As per Surface Water report</li> </ul>			<ul style="list-style-type: none"> <li>■ -</li> </ul>
Assessment of the natural capacity of Lake Albert to provide waste assimilation services, and insurance through monitoring and analysis that these are not exceeded	<ul style="list-style-type: none"> <li>■ Monitoring of lake water quality once assimilation capacity has been calculated</li> </ul>	As required	CNOOC	<ul style="list-style-type: none"> <li>■ -</li> </ul>
The development of an Influx Management Plan will identify appropriate measures to mitigate the expected increased waste-loading to surface water systems as a result of in-migration due to the presence of the project.	<ul style="list-style-type: none"> <li>■ Monitoring measures described in the plan</li> </ul>	<ul style="list-style-type: none"> <li>■ Monitoring frequency described in the plan</li> </ul>	CNOOC	<ul style="list-style-type: none"> <li>■</li> </ul>
<b>■ Fresh Water (Type II)</b>				
Reduce water volumes needed by Project activities through treatment and re-use of process water and waste water	<ul style="list-style-type: none"> <li>■ -</li> </ul>	-	CNOOC	<ul style="list-style-type: none"> <li>■ -</li> </ul>





## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Contribute to water catchment management in association with other Projects in neighbouring exploration blocks to promote equitable sharing of fresh water resources of Lake Albert	■ -	-	CNOOC	■ -
<b>■ Regulating Air Quality</b>				
Loss of vegetation and wetland ecosystems to the Project footprint and associated indirect effects to be addressed by the mitigation measures recommended in the Biodiversity Impact Assessment and the Surface Water Impact Assessment	■ As per Surface Water and Biodiversity Impact Assessments	As required	CNOOC	■ -
Dedicate a portion of the land used for the project for native forest, and/ or invest in replacing or protecting CO2 sequestration/storage services in the immediate area, as part of Corporate Social Responsibility initiatives	■ -	-	CNOOC	■ -
Assess the relative importance of natural air quality regulatory services within the Project Area of Influence, and design infrastructure to accommodate and enhance such services where feasible.	■ -	-	CNOOC	■ -
Community education programmes on pollution prevention and monitoring schemes. Promotion of CNOOC corporate social responsibility initiatives	■ -	-	CNOOC	■ -
<b>■ Regulating Water Flows and Timing</b>				
Where possible, avoid or enhance natural barriers such as wetlands before investing in man-made replacements.	■ As per Surface Water and Biodiversity Impact Assessments	As required	CNOOC	■ -





## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Mitigation measures outlined in the Surface Water Impact Assessment and Biodiversity Impact Assessment report include the incorporation of engineered design features to ensure that water flows in impacted wetland systems are maintained. Mitigation measures should be applied as recommended.	<ul style="list-style-type: none"> <li>■ As per Surface Water and Biodiversity Impact Assessments</li> <li>■</li> <li>■</li> </ul>	As required	CNOOC	■ -
<b>■ Water Purification and Waste Treatment</b>				
Minimising the amount of wetland being directly lost to the Project footprint will contribute to reduction of potential impacts on the supply of this ES. Appropriate engineered mitigation measures at wetland and riparian crossings along the pipeline route, which maintain surface and subsurface flows and subsequently the integrity of these systems will also contribute to minimisation of potential impacts	<ul style="list-style-type: none"> <li>■ As per Biodiversity Impact Assessment</li> </ul>	As required	CNOOC	■
Assessment of the natural capacity of Lake Albert and Project-affected wetlands to provide water filtration and waste assimilation services, and insurance through monitoring and analysis that these are not exceeded	<ul style="list-style-type: none"> <li>■ Monitoring of lake water quality once assimilation capacity has been calculated</li> </ul>	As required	CNOOC	■
Appropriate sewerage facilities and wastewater treatment systems to be put in place at construction camp and at long-term operational Project facilities	<ul style="list-style-type: none"> <li>■ Monitoring of quality of wastewater discharge</li> </ul>	As required	CNOOC	-
The development of an Influx Management Plan will identify appropriate measures to mitigate the expected increased waste-loading to surface water systems as a result of in-migration due to the presence of the project.	<ul style="list-style-type: none"> <li>■ Monitoring measures described in the plan</li> </ul>	<ul style="list-style-type: none"> <li>■ Monitoring frequency described in the plan</li> </ul>	CNOOC	-

**■ Cultural Heritage Ecosystem Services**





## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
In accordance with IFC PS8 (Cultural Heritage), where the Project may significantly impact on <b>critical cultural heritage</b> that is essential to the identity and/or cultural, ceremonial, or spiritual aspects of beneficiaries lives, priority will be given to the avoidance of such impacts.	<ul style="list-style-type: none"> <li>■ Avoid development in areas identified as spiritual or sacred sites</li> </ul>		CNOOC	-
Where significant project impacts on <b>critical cultural heritage</b> are unavoidable, the client will obtain the free, prior and informed consent (FPIC) of the Affected Communities, as per IFC PS8 and PS1 requirements	<ul style="list-style-type: none"> <li>■ Undertake a process of Informed Consultation and Participation of the affected communities</li> </ul>		CNOOC	<ul style="list-style-type: none"> <li>■</li> </ul>
Protection of the environmental setting for sacred sites close to construction / operation areas	<ul style="list-style-type: none"> <li>■ No personnel and/or contractors allowed beyond footprint of Project</li> <li>■ Designated no-go areas, e.g., sacred sites, ritual sites</li> <li>■ Screening planting around Project facilities to protect views</li> </ul>	As required	CNOOC and Contractors	<ul style="list-style-type: none"> <li>■ Inductions for all staff</li> </ul>
Maintaining community access to sacred sites and facilitating respect for local intangible cultural heritage, tradition and taboo will ensure that the negative socio-cultural effects are effectively managed – regular platforms for community liaison are recommended and provisions for such should be made in the Cultural Heritage Management Plan (CHMP)	<ul style="list-style-type: none"> <li>■ As per CHMP</li> </ul>	As required	CNOOC	<ul style="list-style-type: none"> <li>■ Inductions for all staff</li> </ul>



## ECOSYSTEM SERVICES ASSESSMENT

Mitigation Measures	Monitoring Indicators	Monitoring Frequency	Responsible Entity	Training Necessary
Cultural sensitivity training to be provided to Project staff and incorporated into the site induction process	■ As per CHMP	As required	CNOOC and Contractors	■ Inductions for all staff



### 11.0 CONCLUSION

The Project will affect beneficiaries of priority ecosystem services in two main ways; the physical presence of the Project infrastructure, and population influx associated with the construction and operation of the Project.

The presence of the Project infrastructure will cause land cover changes and associated loss of supply of ecosystem services; it will also change the physical landscape of the area which lends itself to the cultural heritage value of Lake Albert and the escarpment to local communities. Population influx of job-seekers, and people seeking to provide commercial services to the increasing population, will increase demand for ecosystem services, and therefore increase pressure on the ecosystems that supply these services.

Other than the actual direct and indirect effects of Project activities, maintenance of the Project's social licence to operate from affected beneficiaries is critical. Local people perceive that oil exploration projects have affected fish stocks in Lake Albert and that air quality has deteriorated as a result of oil-related commercial activity in the area. It is therefore crucial that the mitigation hierarchy is followed and all efforts to avoid impacts on Lake Albert water quality, air quality and sites of cultural heritage value are made.

Where avoidance of impacts is impossible, application of the recommended mitigation measures is crucial. In particular, worker and community education programmes are key in both maintaining CNOOC's social licence to operate in the area, and educating beneficiaries to promote sustainable use of the ecosystem services that they rely on. Appropriate resettlement action plans, livelihood restoration plans and influx management plans are key mitigation measures to ensure that the beneficiaries that are most reliant on priority ecosystem services within the Project Area of Influence are suitably accommodated.



### 12.0 REFERENCES

WSS Services (U) Ltd, 2012. *Assessment of water use and demand in Lake Albert Basin of Uganda*, Kampala: Republic of Uganda - Ministry of water and environment.

AECOM, 2012. *Report on the Environmental Baseline in Exploration Area 2*, s.l.: unpublished report to Tullow Uganda Operations Pty..

AECOM, 2013. *Report on Biodiversity Conservation Opportunities in the Albertine Rift*, s.l.: Tullow Uganda Operations Pty..

Agea, J. G. et al., 2011. Wild and semi-wild food plants in Bunyoro-Kitara Kingdom, Uganda: cultural significance, local perceptions and social implications of their consumption. *International Journal of Medicinal and Aromatic Plants* , 1(2), pp. 137-152.

Akwetaireho, S., Akugizibwe, T. & Plumptre, A., 2011. *Socioeconomic values of corridor forests in the Albertine Rift forests of the Murchison-Semliki landscape*, Kampala, Uganda: Wildlife Conservation Society.

Ayebare, S. et al., 2013. *Identifying climate resilient corridors for conservation in the Albertine Rift* , s.l.: Unpublished Report by the Wildlife Conservation Society to MacArthur Foundation.

BioNET-EAFRINET, 2011. *East African Network for Taxonomy - Keys and Fact Sheets - Parkinsonia aculeata*. [Online]

Available at:

[http://keys.lucidcentral.org/keys/v3/eafrinet/weeds/key/weeds/Media/Html/Parkinsonia\\_aculeata\\_\(Parkinsonia\).htm](http://keys.lucidcentral.org/keys/v3/eafrinet/weeds/key/weeds/Media/Html/Parkinsonia_aculeata_(Parkinsonia).htm)

[Accessed 01 August 2014].

Center for Aquatic and Invasive Plants, University of Florida, 2014. *Water hyacinth*. [Online]

Available at: <http://plants.ifas.ufl.edu/node/141>

[Accessed 13 02 2015].

CRA, 2006. *Ecosystems, ecosystem services and their linkages to poverty reductions in Uganda. Final Report.*, National Environmental Management Authority (NEMA), Uganda: Centre for Resource Analysis (CRA).

Eco & Partners, 2014. *Ecological Baseline Report for the Dry Season Surveys in the Kingfisher Field Development and Pipeline Route Areas*. s.l.:Unpublished.

Environmental Assessment Consult (U) Ltd., 2013. *ESIA for 2D Seismic Testing Acquisition and Processing in Kingfisher Field Development Area, Bugoma, Hoima & Kikuube Districts*, s.l.: s.n.

European Commission, 2001. *Earth Architecture in Uganda - Project to promote forms of architecture that respect the environment and are accessible to disadvantaged populations in Uganda*, s.l.: CRATerre Editions.

Forest Department, 2002. *National Biomass Study, Technical Report of 1996-2002*, Kampala: Ministry of Water Lands and Environment.

Global Climate Change Alliance, 2012. *Adaption for climate change in Uganda*. [Online]

Available at: <http://www.gcca.eu/national-programmes/africa/gcca-uganda>

[Accessed June 2014].

*Draft Socio-economic Impact Assessment for the proposed Kingfisher Development*, s.l.: s.n. 2014 (a)

*Soil and Land Capability Specialist Study - Impact Assessment Report for Kingfisher Field development Area in Hoima & Kikuube District, Uganda*, s.l.: s.n. 2014 (d)

*Draft Biodiversity Impact Assessment - Environmental and Social Impact Assessment for Kingfisher Field Development Area, Hoima & Kikuube District, Uganda*, s.l.: s.n. 2014 (f)







*Draft Surface Water report - Environmental and Social Impact Assessment for Kingfisher Field Development Area, Hoima & Kikuube District, Uganda, s.l.: s.n. 2014(b)*

*Scoping Report for the Environmental and Social Impact Assessment for Kingfisher Field development Area in Hoima & Kikuube District, Uganda by CNOOC Uganda Ltd., s.l.: s.n. 2014 (c)*

*ESIA for Kingfisher 3A Block, Hoima District, Uganda. Cultural Heritage Report, s.l.: s.n. 2014(e)*

Hanson, C., Ranganathan, J., Iceland, C. & Finisdore, J., 2012. *The Corporate Ecosystem Services Review. Guidelines for Identifying Business Risks and Opportunities Arising from Ecosystem Change*. Version 2.0 ed. s.l.:World Resources Institute.

Hoima District Local Government, 2011. *Hoima District Development Plan 2011/12-2015/16, s.l.: s.n.*

IFC, 2012a. *Performance Standard 6: Biodiversity Conservation and Sustainable Management of Living Natural Resources, s.l.: World Bank Group.*

IFC, 2012b. *Guidance Note 6: Biodiversity Conservation and Sustainable Management of Living Natural Resources, s.l.: World Bank Group.*

International Lake Environmental Committee, n.d. *Lake Albert*. [Online]  
Available at: [http://wldb.ilec.or.jp/data/databook\\_html/afr/afr-11.html](http://wldb.ilec.or.jp/data/databook_html/afr/afr-11.html)

International Lake Environment Committee Foundation, 1999. *World Lake Database*. [Online]  
Available at: [http://wldb.ilec.or.jp/data/databook\\_html/afr/afr-11.html](http://wldb.ilec.or.jp/data/databook_html/afr/afr-11.html)  
[Accessed 03 06 2014].

IPIECA, 2011. *Ecosystem Services Guidance - Biodiversity and Ecosystem Services guide and checklist, s.l.: s.n.*

Katende, A. B., Birnie, A. & Tengnas, B., 1995. *Useful trees and shrubs for Uganda: Identification, propagation and management for agricultural and pastoral communities. Technical Handbook No 10.. s.l.:Regional Soil Conservation Unit, RSCU/SIDA, Kenya.*

Kent, M. & Coker, P., 1992. *Vegetation description and analysis*. 1st ed. Chichester, England: Joh Wiley & Sons.

Landsberg, F. et al., 2013. *Weaving Ecosystem Services into Impact Assessment*. s.l.:World Resources Institute.

Landsberg, F. et al., 2013. *Weaving Ecosystem Services into Impact Assessment*, s.l.: World Resources Institute.

Lowe, S., Browne , M., Boudjelas, S. & De Poorter, M., 2000. *100 of the World's Worst Invasive Alien Species. A selection from the Global Invasive Species Database.*, 12pp: The Invasive Species Specialist Group (ISSG) a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN).

Macfarlane, D. M. et al., 2008. *WET-Health A technique for rapidly assessing wetland health. WRC Report TT 340/08*, Pretoria: Water Research Commission.

McLaughlin, D. M. & Cohen, M. J., 2013. Realizing ecosystem services: wetland hydrologic function along a gradient of ecosystem condition. *Ecological Applications*, Volume 23, pp. 1619-1631.

Nelson, G. C., 2005. Drivers of Ecosystem Change: Summary Chapter. In: R. Hassan, R. Scholes & N. Ash, eds. *Ecosystems and Human Well-being: Current State and Trends*. s.l.:Island Press, pp. 73-76.

NEMA, 1996. *State of the environment Report for Uganda*. Kampala: National Environmental Management Authority.



- NEMA, 2002. *National Biodiversity Strategy and Action Plan*, The Republic of Uganda: National Environment Management Authority (NEMA).
- Olupot, W., McNeilage, A. J. & Plumptre, A. J., 2009. *An analysis of socioeconomics of bushmeat hunting at major hunting sites in Uganda. Working Paper No. 38*, s.l.: Wildlife Conservation Society.
- PANParks, unknown. *Tourism as an Ecosystem Service - Financing tourism in Protected Areas and introduction to PAN Parks Foundation as a case study*. [Online]  
Available at: [http://www.eurosite.org/files/preview\\_of\\_4\\_kun -- tourism as an ecosystem service - case study of pan parks.ppt .pdf](http://www.eurosite.org/files/preview_of_4_kun_-_tourism_as_an_ecosystem_service_-_case_study_of_pan_parks.ppt_.pdf)  
[Accessed 13 02 2015].
- Plumptre , A. et al., 2010. *Biodiversity Surveys of the Bugoma Forest Reserve, Smaller Central Forest Reserves, and Corridor Forests South of Bugoma*, s.l.: Wildlife Conservation Society.
- Plumptre, A. J., 2002. *Extent and Status of the Forests in the Ugandan Albertine Rift* , s.l.: Wildlife Conservation Society.
- Ramsar, 1992. *Ramsar Wetlands - Uganda*. [Online]  
Available at: <http://ramsar.wetlands.org/Portals/15/UGANDA.pdf>  
[Accessed June 2014].
- Scullion, S., 2007. *Inland Fisheries Co-Management in East Africa*. s.l., Lake Victoria Fisheries Organisation.
- Secretariat of the Convention on Biological Diversity, 2006. *Biodiversity in Impact Assessment, Background Document to CBD Decision VIII/28: Voluntary Guidelines on Biodiversity-Inclusive Impact Assessment*, s.l.: Netherlands Commission for Environmental Assessment.
- Smith, M. et al., 2013. Long-term effects of fire frequency and season on herbaceous vegetation in savannas of the Kruger National Park, South Africa.. *Journal of Plant Ecology*, 6(1), pp. 71-83.
- Solomon, S. & George, K., 2012. *Report for the development of Marine Tourism in Uganda - PHase II (Lake Kyoga, Lake Albert and Waterfalls)*, s.l.: The Uganda Tourism Board.
- Taabu-Munyaho, A. et al., 2012. *Report of the Catch Assessment Survey of Lake Albert – Albert Nile Conducted in July 2012. ATAAS Project 5: Monitoring changes in stocks of the major commercial fish species in the five major waterbodies of Uganda (Victoria, Albert, Kyoga, Edward, George, Uganda: National Fisheries Resource Research Institute*.
- TEEB, 2010. *he Economics of Ecosystems and Biodiversity: Mainstreaming the Economics of Nature: A Synthesis of the Approach, Conclusions and Recommendations of TEEB*, s.l.: The Economics of Ecosystems and Biodiversity.
- Tengberg, A. et al., 2012. Cultural ecosystem services provided by landscapes: Assessment of heritage values and identity. *Ecosystem Services*, Volume 2, pp. 14-26.
- Treweek (ed), J., 1999. *Ecological Impact Assessment*. London: Blackwell Scientific.
- Tucker, G., 2005. Biodiversity Evaluation Methods. In: D. Hill, et al. eds. *Handbook of Biodiversity Methods*. Cambridge, UK: Cambridge University Press.
- Twongo, T., 1996. Growing Impact of Water Hyacinth on Nearshore Environments on Lakes Victoria and Kyoga (East Africa). In: I. A. Johnson, ed. *Limnology, Climatology and Paleoclimatology of the East African Lakes*. s.l.:Gordon and Breach Science Publishers SA., pp. 633-637.
- Uganda Nile Discourse Forum, 2013. *Baseline Survey on Lakes Edward and Albert Integrated Fisheries and Water Resources Management Project*, s.l.: Uganda Nile Discourse Forum.
- UNESCO, 2005. *Bark Cloth Making in Uganda*. [Online]  
Available at: [http://www.unesco.org/culture/intangible-heritage/40afr\\_uk.htm](http://www.unesco.org/culture/intangible-heritage/40afr_uk.htm)  
[Accessed 13 08 20114].





Wandera, S. B., 2000. *The fisheries of the Ugandan sector of Lake Albert. A report on the current state of the fishery of the Ugandan sector of Lake Albert.*, 29pp: Prepared for FAO sub-regional representative for Southern and Eastern Africa (SAFD).

Wandera, S. B. & Balirwa, J. S., 2010. Fish species diversity and relative abundance in Lake Albert—Uganda.. *Aquatic Ecosystem Health & Management*, 13(3), p. 284–293.

WHO, 2011. *World Health Organisation Guidelines for Drinking-Water Quality*. Geneva: WHO Press.

Zommers, Z. & MacDonald, D. W., 2012. Protected Areas as Frontiers for Human Migration.. *Conservation Biology*, 26(3), p. 547–556.



# **APPENDIX A**

## **Document Limitations**



### DOCUMENT LIMITATIONS

This Document has been provided by Independent Consultants subject to the following limitations:

- i) This Document has been prepared for the particular purpose outlined in the Consultants' proposal and no responsibility is accepted for the use of this Document, in whole or in part, in other contexts or for any other purpose.
- ii) The scope and the period of the Consultants' Services are as described in Consultants' proposal, and are subject to restrictions and limitations. The Consultant did not perform a complete assessment of all possible conditions or circumstances that may exist at the site referenced in the Document. If a service is not expressly indicated, do not assume it has been provided. If a matter is not addressed, do not assume that any determination has been made by the Consultants in regards to it.
- iii) Conditions may exist which were undetectable given the limited nature of the enquiry the Consultant was retained to undertake with respect to the site. Variations in conditions may occur between investigatory locations, and there may be special conditions pertaining to the site which have not been revealed by the investigation and which have not therefore been taken into account in the Document. Accordingly, additional studies and actions may be required.
- iv) In addition, it is recognised that the passage of time affects the information and assessment provided in this Document. The Consultants' opinions are based upon information that existed at the time of the production of the Document. It is understood that the Services provided allowed Consultant to form no more than an opinion of the actual conditions of the site at the time the site was visited and cannot be used to assess the effect of any subsequent changes in the quality of the site, or its surroundings, or any laws or regulations.
- v) Any assessments made in this Document are based on the conditions indicated from published sources and the investigation described. No warranty is included, either express or implied, that the actual conditions will conform exactly to the assessments contained in this Document.
- vi) Where data supplied by the client or other external sources, including previous site investigation data, have been used, it has been assumed that the information is correct unless otherwise stated. No responsibility is accepted by the Consultants for incomplete or inaccurate data supplied by others.
- vii) The Client acknowledges that the Consultants may have retained sub-consultants affiliated with the Consultants to provide Services for the benefit of the Consultants. The Consultants will be fully responsible to the Client for the Services and work done by all of its sub-consultants and subcontractors. The Client agrees that it will only assert claims against and seek to recover losses, damages or other liabilities from the Consultant and not the Consultants' affiliated companies. To the maximum extent allowed by law, the Client acknowledges and agrees it will not have any legal recourse, and waives any expense, loss, claim, demand, or cause of action, against the Consultant's affiliated companies, and their employees, officers and directors.
- viii) This Document is provided for sole use by the Client and is confidential to it and its professional advisers. No responsibility whatsoever for the contents of this Document will be accepted to any person other than the Client. Any use which a third party makes of this Document, or any reliance on or decisions to be made based on it, is the responsibility of such third parties. the Consultants accept no responsibility for damages, if any, suffered by any third party as a result of decisions made or actions based on this Document.



# APPENDIX B

## Prioritisation of ES according to Project Impact



## ECOSYSTEM SERVICES ASSESSMENT

### PRIORITISATION OF ES ACCORDING TO PROJECT IMPACT

Priority ecosystem services are those services for which the answers to questions 1 and 2 are “Yes” or “Unknown”, **and** “No” or “Unknown” to question 3.

#### Impact prioritisation spreadsheet

Ecosystem Service	Potentially affected beneficiaries	Potentially affected benefits	1. Could the project affect the ability of others to benefit from this ES? (Y/N/?)	2. Is this ES important to beneficiaries' livelihoods, health, safety or culture? (Y/N/?)	3. Do beneficiaries have viable alternative to this ES? (Y/N/?)	Priority ES 1 = Priority 0 = Non-priority
<b>Provisioning</b>						
Food – Subsistence crops	Pipeline route community	Income, livelihoods, food intake	Y	Y	?	1
Food - Grazing for Livestock	Buhuka Flats community	Reduced grazing area due to Project land-take and increased pressure from population influx will reduce grazing availability, which may limit the ability of people to raise livestock for subsistence and livelihood purposes	?	Y	N	1
	Migratory herders	As above	?	Y	N	1
Food – Capture fisheries	Buhuka Flats community	Income, livelihoods, food intake	Y – stakeholder perception that the Project will affect fish stocks, increased demand from population influx	Y	N	1
Food – wild foods	Pipeline route community	Reduced bush meat availability due to reductions in woodland/bush land cover that supports hunted species Reduced vegetation cover may limit bee's ability to produce honey and honey production	?	Y	?	1
Biological raw materials – construction of traditional houses	Buhuka Flats and Pipeline route communities	Ability to construct homes and animal shelters	?	Y	Y	0





## ECOSYSTEM SERVICES ASSESSMENT

Ecosystem Service	Potentially affected beneficiaries	Potentially affected benefits	1. Could the project affect the ability of others to benefit from this ES? (Y/N/?)	2. Is this ES important to beneficiaries' livelihoods, health, safety or culture? (Y/N/?)	3. Do beneficiaries have viable alternative to this ES? (Y/N/?)	Priority ES 1 = Priority 0 = Non-priority
Biological raw materials – extraction of aggregates for Project Construction	The Project Buhuka Flats and Pipeline route communities	Landscape value and spiritual and inspirational values Impacts via land take of ecosystems that may provide priority ecosystem services	?	Y	N	1
Biomass fuel – wood and charcoal	Buhuka Flats and Pipeline route communities	Energy sources for cooking, fish processing, brick making	Y	Y	Y – most fuel sources in Project area of influence have been exhausted and charcoal and fire wood are purchased	1
Fresh water	Buhuka Flats and Pipeline route communities	Availability and quality of fresh water for drinking may be compromised by abstraction from Lake Albert and interception of sources by the pipeline	Y – stakeholder perception that the Project will affect water quality	Y	N	1
Medicinal plants	Buhuka Flats and Pipeline route communities	Availability of traditional medicines	N – areas of forest and grassland loss to footprint in context are minimal	n/a	n/a	0
<b>Regulating</b>						
Air quality	Buhuka Flats and Pipeline route communities	Project effects on ecosystems that provide this ecosystem service are negligible in the context of available unaffected areas in LSA	N	n/a	n/a	0
Water flows and timing	Buhuka Flats community	Wetlands and the unique hydrological system of the Buhuka Flats may be disturbed/interrupted	Y	Y	N	1
Soil stability & erosion control	Buhuka Flats community	Vegetation clearance for construction may reduce the ability of the surrounding soils to withstand erosive forces of wind and floods	N	n/a	n/a	0





## ECOSYSTEM SERVICES ASSESSMENT

Ecosystem Service	Potentially affected beneficiaries	Potentially affected benefits	1. Could the project affect the ability of others to benefit from this ES? (Y/N/?)	2. Is this ES important to beneficiaries' livelihoods, health, safety or culture? (Y/N/?)	3. Do beneficiaries have viable alternative to this ES? (Y/N/?)	Priority ES 1 = Priority 0 = Non-priority
Water purification and waste treatment	Buhuka Flats and Pipeline route communities	Disturbance of wetlands by proposed infrastructure may impact the integrity of wetlands and their ability to provide ES Population influx may increase nutrient loading and pressurise assimilative capacity of Lake Albert	Y	Y	N	1
<b>Cultural</b>						
Recreation and ecotourism	Buhuka Flats community	Tourism associated with oil company staff is currently developing – development may ultimately be restricted by Project presence and visual impact effects	Y	N	n/a	0
Ethical and spiritual values	Buhuka Flats community	Sacred sites and intangible cultural heritage are intrinsically linked with natural ecosystems such as wetlands, rivers, lake and forests and substantially contribute to beneficiaries' sense of identity	Y	Y	N	1
Educational and inspirational	Buhuka Flats community	the Lake Albert and Escarpment landscapes inspire folklore and contribute to beneficiaries' sense of heritage and identity	Y	Y	N	1



# **APPENDIX C**

## **Prioritisation of ES according to Project Demand**



## ECOSYSTEM SERVICES ASSESSMENT

### PRIORITISATION OF ES ACCORDING TO PROJECT DEPENDENCE

Priority ES are those services for which the answers to question 1 is “Yes” or “Unknown”, **and** “No” or “Unknown” to question 2. If the answer to question 1 is no, it is automatically a non-priority ecosystem services. Changes in an ecosystem services can be driven both by causes of ecosystem change external to the Project and by the Project’s own impacts.

#### Priority ecosystem services according to the extent of Project Demand

Ecosystem Service	1. Could this ES change in ways that will affect operational performance (Y/N/?)	2. Does the Project have viable alternatives to this ES (Y/N/?)	Priority ES 1 = Priority 0 = Non-priority
<b>Provisioning</b>			
Food – Subsistence crops	N	n/a	0
Food - Grazing for Livestock	N	n/a	0
Food – Capture fisheries	N	N	0
Food – wild foods	N	n/a	0
Biological raw materials – aggregate extraction for Project facility construction	?	N	1
Biomass fuel – wood and charcoal	N	n/a	0
Fresh water	Y – stakeholders perceive that the Project may impact the quality of water, therefore the Project is reliant on the quality and quantity of freshwater remaining constant throughout its lifetime in order to maintain its social license to operate  Cumulative impact of abstraction by other projects unknown	N	1
Medicinal plants	N	n/a	0
<b>Regulating</b>			
Air quality	? – stakeholders perceive that the Project may impact the air quality, therefore the Project is reliant on the quality of air remaining constant throughout its lifetime in order to maintain the social license to operate	N	1





## ECOSYSTEM SERVICES ASSESSMENT

Ecosystem Service	1. Could this ES change in ways that will affect operational performance (Y/N/?)	2. Does the Project have viable alternatives to this ES (Y/N/?)	Priority ES 1 = Priority 0 = Non-priority
Climate regulation	Y – the expected changes in climatic conditions over the lifetime of the Project could lead to a rise in Lake Albert’s water level, which would result in flooding of the flats, which could affect operational performance	Y – design and engineering mitigation measures take into account flood and climate modelling predictions	0
Water flows and timing	Y – Soils, wetlands and drainage lines influence the timing and magnitude of water runoff, flooding and aquifer recharge. The Project will affect wetlands and drainage lines which may cause flooding on the Buhuka Flats, which could affect operational performance	Y – engineering mitigation measures to manage surface and sub-surface flows in the construction and operation phases of the Project are considered sufficient to reduce potential impacts to negligible significance	0
Soil stability & erosion control	Y – predicted rise in the level of Lake Albert over the Project lifetime could lead to an increase in erosion of the shoreline, thereby reducing the width of the Flats and increasing flood likelihood. Vegetation removal for site clearance could also contribute	N – Engineered measures for the control of erosion arising from vegetation removal are considered sufficient to minimise the impacts of vegetation clearance. The Project, in compliance with the requirements of IFC Performance Standards 1 and 3, has undertaken predictive modelling to ensure that the Project’s operational performance will not be put at risk by rises in Lake Albert’s level	0
Water purification and waste treatment	Y - Nutrient assimilative capacity of the lake may reach thresholds, resulting in eutrophication	Y – Project incorporates a water treatment system which can be used as necessary	0
<b>Cultural</b>			
Recreation and ecotourism	N	n/a	0
Ethical and spiritual values	Y – the Project is reliant on the availability of this ES remaining constant throughout its lifetime in order to maintain its social license to operate	N – there are no alternatives to the presence of the Project in the landscape	1
Educational and inspirational	? – the Project could be reliant on the availability of this ES remaining constant throughout its lifetime in order to maintain its social license to operate	N – there are no alternatives to the presence of the Project in the landscape	1





